



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:00 PM GMT

PDB ID : 3VRY
Title : Crystal structure of HCK complexed with a pyrrolo-pyrimidine inhibitor 4-Amino-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl-cyclopentane
Authors : Kuratani, M.; Tomabechei, Y.; Niwa, H.; Handa, N.; Yokoyama, S.
Deposited on : 2012-04-21
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4309 (2.50-2.46)
Clashscore	102246	5050 (2.50-2.46)
Ramachandran outliers	100387	4961 (2.50-2.46)
Sidechain outliers	100360	4963 (2.50-2.46)
RSRZ outliers	91569	4319 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	<div> <div>3%</div> <div>79%</div> <div>13%</div> <div>6%</div> </div>
1	B	454	<div> <div>4%</div> <div>79%</div> <div>16%</div> <div>• •</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase HCK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	P	S	0	0	0
			3473	2225	582	645	1	20			
1	B	435	Total	C	N	O	P	S	0	0	0
			3518	2250	592	655	1	20			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLY	-	EXPRESSION TAG	UNP P08631
A	79	ALA	-	EXPRESSION TAG	UNP P08631
A	80	MET	-	EXPRESSION TAG	UNP P08631
A	81	GLY	-	EXPRESSION TAG	UNP P08631
A	82	SER	-	EXPRESSION TAG	UNP P08631
A	83	GLY	-	EXPRESSION TAG	UNP P08631
A	84	ILE	-	EXPRESSION TAG	UNP P08631
A	85	ARG	-	EXPRESSION TAG	UNP P08631
A	528	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	529	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	530	ILE	GLN	ENGINEERED MUTATION	UNP P08631
B	78	GLY	-	EXPRESSION TAG	UNP P08631
B	79	ALA	-	EXPRESSION TAG	UNP P08631
B	80	MET	-	EXPRESSION TAG	UNP P08631
B	81	GLY	-	EXPRESSION TAG	UNP P08631
B	82	SER	-	EXPRESSION TAG	UNP P08631
B	83	GLY	-	EXPRESSION TAG	UNP P08631
B	84	ILE	-	EXPRESSION TAG	UNP P08631
B	85	ARG	-	EXPRESSION TAG	UNP P08631
B	528	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	529	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	530	ILE	GLN	ENGINEERED MUTATION	UNP P08631

- Molecule 2 is 4-AMINO-5-(4-PHENOXYPHENYL)-7H-PYRROLO[2,3-D]PYRIMIDIN-7-Y L-CYCLOPENTANE (three-letter code: B43) (formula: C₂₃H₂₂N₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	23	4	1		
2	B	1	Total	C	N	O	0	0
			28	23	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	117	Total	O	0	0
			117	117		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	120	Total	O	0	0
			120	120		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase HCK



• Molecule 1: Tyrosine-protein kinase HCK



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.59Å 73.88Å 178.47Å 90.00° 95.84° 90.00°	Depositor
Resolution (Å)	40.75 – 2.48 40.75 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.0 (40.75-2.48) 99.0 (40.75-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.48Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.226 , 0.281 0.222 , 0.275	Depositor DCC
R_{free} test set	2220 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.6	EDS
Estimated twinning fraction	0.029 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 44395 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7288	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PTR, B43, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3539	0.59	0/4776
1	B	0.43	0/3585	0.56	0/4839
All	All	0.44	0/7124	0.58	0/9615

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3473	0	3444	38	0
1	B	3518	0	3490	42	0
2	A	28	0	22	0	0
2	B	28	0	22	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	117	0	0	3	0
5	B	120	0	0	5	0
All	All	7288	0	6978	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:LYS:O	1:A:241:LYS:NZ	2.14	0.80
1:A:181:THR:OG1	1:A:204:LYS:NZ	2.17	0.76
1:B:485:GLU:O	1:B:486:ASN:ND2	2.16	0.73
1:B:489:GLU:OE1	5:B:794:HOH:O	2.08	0.70
1:B:378:GLU:HG3	1:B:441:ILE:HG12	1.73	0.70
1:A:191:ASP:OD1	1:A:192:TYR:N	2.25	0.69
1:A:489:GLU:HG2	1:B:206:ARG:HH22	1.56	0.69
1:B:380:ARG:NH1	5:B:740:HOH:O	2.21	0.68
1:A:239:CYS:SG	5:A:792:HOH:O	2.42	0.67
1:B:308:LEU:HD11	1:B:330:THR:HG22	1.77	0.67
1:B:250:LYS:H	1:B:250:LYS:HD3	1.58	0.66
1:B:322:LEU:HD22	1:B:402:ILE:HB	1.81	0.62
1:B:362:LYS:NZ	5:B:765:HOH:O	2.29	0.62
1:A:156:ARG:N	1:A:179:GLU:OE2	2.30	0.61
1:B:485:GLU:H	1:B:485:GLU:CD	2.05	0.59
1:A:149:TRP:HZ2	1:A:225:GLN:HG2	1.67	0.59
1:A:385:ARG:HG3	1:A:439:PHE:CD2	2.38	0.59
1:A:343:LYS:NZ	1:A:354:GLU:OE1	2.37	0.58
1:A:221:PHE:HZ	1:A:230:HIS:CD2	2.22	0.57
1:B:388:ARG:HB3	1:B:428:TRP:CD1	2.40	0.57
1:A:239:CYS:O	1:A:240:GLN:NE2	2.39	0.56
1:A:234:GLY:H	1:A:241:LYS:NZ	2.04	0.55
1:B:264:ARG:NH2	1:B:331:LYS:O	2.40	0.55
1:B:365:ASP:O	1:B:369:GLN:HG3	2.07	0.55
1:A:300:GLY:C	1:A:302:MET:HB2	2.29	0.52
1:B:307:PHE:CE2	1:B:334:ILE:HG21	2.44	0.52
1:B:243:SER:OG	1:B:244:VAL:N	2.43	0.52
1:A:252:GLN:HE22	1:A:339:GLU:HB3	1.75	0.51
1:B:448:PHE:CE2	1:B:452:LEU:HD13	2.46	0.51
1:A:225:GLN:NE2	1:A:229:ASP:OD2	2.44	0.50
1:B:494:ILE:HD11	1:B:515:VAL:HG11	1.94	0.50
1:A:90:ALA:HA	1:A:138:VAL:HG12	1.93	0.50
1:A:523:THR:O	1:B:490:GLU:HG2	2.11	0.50
1:B:377:ILE:HG23	1:B:382:TYR:HB3	1.94	0.50
1:A:446:TRP:CE3	1:A:499:TRP:HA	2.47	0.50
1:A:97:ILE:HD13	1:A:253:LYS:HE3	1.93	0.49
1:B:483:ARG:NH1	1:B:487:CYS:O	2.40	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:479:TYR:O	1:B:480:ARG:HD3	2.14	0.48
1:A:178:SER:HB2	1:A:527:PTR:O3P	2.14	0.47
1:A:530:ILE:HD13	1:A:531:PRO:HD2	1.97	0.47
1:B:426:ILE:HD11	1:B:468:ASN:HB3	1.96	0.46
1:B:92:TYR:CD1	1:B:251:PRO:HG3	2.50	0.46
1:B:172:SER:HA	1:B:244:VAL:O	2.16	0.46
1:B:445:VAL:HG11	1:B:506:ARG:HB3	1.97	0.46
1:A:324:LYS:HE3	5:A:716:HOH:O	2.14	0.46
1:B:160:GLU:HA	1:B:200:VAL:HG21	1.97	0.46
1:B:466:MET:HE2	1:B:471:VAL:HG22	1.97	0.46
1:B:206:ARG:HB2	1:B:214:TYR:CE1	2.51	0.45
1:A:310:GLU:O	1:A:314:MET:HG3	2.17	0.45
1:B:119:TRP:CZ2	1:B:254:PRO:HG2	2.50	0.45
1:A:291:LYS:HE2	1:A:340:PHE:CD2	2.52	0.44
1:B:137:TYR:CZ	1:B:251:PRO:HG2	2.52	0.44
1:B:206:ARG:HB2	1:B:214:TYR:CZ	2.53	0.44
1:B:207:THR:HG22	1:B:213:PHE:CE1	2.52	0.44
1:B:379:GLN:NE2	5:B:778:HOH:O	2.51	0.44
1:B:467:SER:OG	1:B:470:GLU:HB2	2.17	0.44
1:A:427:LYS:HE2	1:A:471:VAL:HG21	2.00	0.44
1:A:216:SER:OG	1:A:217:PRO:HD2	2.19	0.43
1:A:307:PHE:CE2	1:A:334:ILE:HG21	2.53	0.43
1:A:279:GLY:HA3	1:A:296:THR:O	2.19	0.43
1:A:489:GLU:HG3	1:B:204:LYS:HG3	2.00	0.43
1:B:461:ILE:HA	1:B:462:PRO:HD2	1.89	0.42
1:A:430:ALA:HA	1:A:431:PRO:HD3	1.88	0.42
1:A:301:SER:N	1:A:302:MET:HB2	2.35	0.42
1:A:169:MET:HG3	1:A:170:LEU:O	2.19	0.42
1:B:169:MET:O	1:B:190:ARG:HG2	2.20	0.42
1:B:411:ILE:HA	1:B:411:ILE:HD12	1.83	0.42
1:A:234:GLY:H	1:A:241:LYS:HZ2	1.68	0.41
1:B:160:GLU:HG2	1:B:200:VAL:HB	2.01	0.41
1:A:203:TYR:OH	5:A:792:HOH:O	2.22	0.41
1:B:446:TRP:CE3	1:B:499:TRP:HA	2.56	0.41
1:A:432:GLU:HG3	1:A:432:GLU:H	1.59	0.41
1:A:441:ILE:HD12	1:A:441:ILE:HA	1.95	0.41
1:A:235:ASN:O	1:A:235:ASN:ND2	2.51	0.41
1:B:201:LYS:HD2	1:B:528:GLU:OE2	2.20	0.41
1:B:207:THR:HG22	1:B:213:PHE:CD1	2.55	0.41
2:B:602:B43:H19	5:B:816:HOH:O	2.21	0.41
1:A:452:LEU:HD23	1:A:495:MET:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	422/454 (93%)	405 (96%)	15 (4%)	2 (0%)	34	53
1	B	430/454 (95%)	411 (96%)	17 (4%)	2 (0%)	34	53
All	All	852/908 (94%)	816 (96%)	32 (4%)	4 (0%)	34	53

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	SER
1	A	303	SER
1	B	302	MET
1	B	410	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/393 (96%)	361 (96%)	15 (4%)	38	63
1	B	381/393 (97%)	364 (96%)	17 (4%)	34	57
All	All	757/786 (96%)	725 (96%)	32 (4%)	36	60

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	ILE
1	A	178	SER
1	A	181	THR
1	A	193	ASP
1	A	218	ARG
1	A	225	GLN
1	A	235	ASN
1	A	240	GLN
1	A	241	LYS
1	A	248	SER
1	A	249	SER
1	A	303	SER
1	A	387	LEU
1	A	432	GLU
1	A	456	VAL
1	B	86	ILE
1	B	92	TYR
1	B	124	SER
1	B	208	LEU
1	B	218	ARG
1	B	250	LYS
1	B	252	GLN
1	B	302	MET
1	B	308	LEU
1	B	328	VAL
1	B	387	LEU
1	B	432	GLU
1	B	456	VAL
1	B	470	GLU
1	B	476	GLU
1	B	486	ASN
1	B	530	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	HIS
1	A	240	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PTR	A	527	1,4	14,16,17	1.22	1 (7%)	18,22,24	0.91	2 (11%)
1	PTR	B	527	1,4	14,16,17	1.05	1 (7%)	18,22,24	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	527	1,4	-	0/9/11/13	0/1/1/1
1	PTR	B	527	1,4	-	0/9/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	527	PTR	OH-CZ	-4.03	1.30	1.40
1	B	527	PTR	OH-CZ	-3.65	1.31	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	527	PTR	O-C-CA	-2.08	120.08	125.49
1	A	527	PTR	O2P-P-OH	2.37	113.73	105.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	527	PTR	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	B43	A	601	-	30,32,32	1.00	1 (3%)	32,45,45	2.98	3 (9%)
2	B43	B	602	-	30,32,32	0.88	1 (3%)	32,45,45	2.81	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B43	A	601	-	-	0/8/19/19	0/5/5/5
2	B43	B	602	-	-	0/8/19/19	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	B43	C16-N2	-3.75	1.32	1.38
2	B	602	B43	C16-N2	-3.25	1.33	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	B43	N3-C23-N4	-11.35	120.20	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	602	B43	N3-C23-N4	-10.77	120.64	128.89
2	A	601	B43	C16-N2-C17	-10.55	116.57	125.44
2	B	602	B43	C16-N2-C17	-10.23	116.84	125.44
2	A	601	B43	C16-C3-C4	-4.30	117.19	125.37
2	B	602	B43	C16-C3-C4	-4.22	117.35	125.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	602	B43	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/454 (94%)	0.15	14 (3%) 50 54	19, 40, 72, 85	0
1	B	434/454 (95%)	0.19	17 (3%) 43 47	24, 41, 71, 92	0
All	All	862/908 (94%)	0.17	31 (3%) 46 51	19, 41, 72, 92	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	213	PHE	5.3
1	B	424	PHE	4.5
1	A	531	PRO	4.3
1	A	214	TYR	3.8
1	B	301	SER	3.7
1	A	142	ASP	3.6
1	B	211	GLY	3.4
1	B	210	ASN	3.4
1	A	223	THR	3.2
1	B	438	SER	2.7
1	B	302	MET	2.7
1	A	222	SER	2.6
1	B	465	GLY	2.5
1	B	469	PRO	2.5
1	B	476	GLU	2.5
1	A	233	LYS	2.5
1	B	304	VAL	2.5
1	B	142	ASP	2.4
1	A	220	THR	2.4
1	B	426	ILE	2.4
1	B	382	TYR	2.4
1	A	216	SER	2.3
1	A	219	SER	2.3
1	A	301	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	305	GLU	2.3
1	B	436	PHE	2.3
1	A	227	LEU	2.2
1	A	465	GLY	2.2
1	B	208	LEU	2.1
1	A	244	VAL	2.1
1	B	409	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	PTR	A	527	16/17	0.96	0.16	-	40,48,56,61	0
1	PTR	B	527	16/17	0.97	0.15	-	22,33,37,37	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	B43	B	602	28/28	0.95	0.18	1.47	22,31,39,41	0
4	CA	A	603	1/1	0.81	0.24	1.35	43,43,43,43	0
2	B43	A	601	28/28	0.96	0.17	0.62	18,26,32,35	0
4	CA	B	601	1/1	0.86	0.19	0.50	57,57,57,57	0
3	CL	B	603	1/1	0.99	0.09	-2.80	43,43,43,43	0
3	CL	A	602	1/1	0.97	0.09	-	41,41,41,41	0

6.5 Other polymers

There are no such residues in this entry.