



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:00 PM GMT

PDB ID : 3VS3  
Title : Crystal structure of HCK complexed with a pyrrolo-pyrimidine inhibitor 7-[trans-4-(4-methylpiperazin-1-yl)cyclohexyl]-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine  
Authors : Kuratani, M.; Tomaebchi, Y.; Handa, N.; Yokoyama, S.  
Deposited on : 2012-04-21  
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	5130 (2.20-2.16)
Clashscore	102246	5965 (2.20-2.16)
Ramachandran outliers	100387	5863 (2.20-2.16)
Sidechain outliers	100360	5864 (2.20-2.16)
RSRZ outliers	91569	5142 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	<div> <div>13%</div> <div>86%</div> <div>9%</div> <div>• •</div> </div>
1	B	454	<div> <div>9%</div> <div>82%</div> <div>12%</div> <div>• 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CA	A	602	-	-	-	X
3	CA	B	602	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7434 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase HCK.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	435	Total	C	N	O	P	S	0	0	0
			3518	2250	592	655	1	20			
1	B	432	Total	C	N	O	P	S	0	0	0
			3504	2243	589	651	1	20			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLY	-	EXPRESSION TAG	UNP P08631
A	79	ALA	-	EXPRESSION TAG	UNP P08631
A	80	MET	-	EXPRESSION TAG	UNP P08631
A	81	GLY	-	EXPRESSION TAG	UNP P08631
A	82	SER	-	EXPRESSION TAG	UNP P08631
A	83	GLY	-	EXPRESSION TAG	UNP P08631
A	84	ILE	-	EXPRESSION TAG	UNP P08631
A	85	ARG	-	EXPRESSION TAG	UNP P08631
A	528	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	529	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	530	ILE	GLN	ENGINEERED MUTATION	UNP P08631
B	78	GLY	-	EXPRESSION TAG	UNP P08631
B	79	ALA	-	EXPRESSION TAG	UNP P08631
B	80	MET	-	EXPRESSION TAG	UNP P08631
B	81	GLY	-	EXPRESSION TAG	UNP P08631
B	82	SER	-	EXPRESSION TAG	UNP P08631
B	83	GLY	-	EXPRESSION TAG	UNP P08631
B	84	ILE	-	EXPRESSION TAG	UNP P08631
B	85	ARG	-	EXPRESSION TAG	UNP P08631
B	528	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	529	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	530	ILE	GLN	ENGINEERED MUTATION	UNP P08631

- Molecule 2 is 7-[TRANS-4-(4-METHYLPIPERAZIN-1-YL)CYCLOHEXYL]-5-(4-PHE NOXYPHENYL)-7H-PYRROLO[2,3-D]PYRIMIDIN-4-AMINE (three-letter code: VSE)

(formula: C<sub>29</sub>H<sub>34</sub>N<sub>6</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 36	C 29	N 6	O 1	0	0
2	B	1	Total 36	C 29	N 6	O 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Ca 1 1	0	0
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Cl 1 1	0	0
4	A	1	Total Cl 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	164	Total 164	O 164	0	0
5	B	172	Total 172	O 172	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Tyrosine-protein kinase HCK



#### • Molecule 1: Tyrosine-protein kinase HCK



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.45Å 73.59Å 180.71Å 90.00° 96.30° 90.00°	Depositor
Resolution (Å)	31.46 – 2.17 31.46 – 2.17	Depositor EDS
% Data completeness (in resolution range)	98.8 (31.46-2.17) 98.8 (31.46-2.17)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.38 (at 2.18Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.226 , 0.279 0.222 , 0.273	Depositor DCC
$R_{free}$ test set	3357 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 58.7	EDS
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 66227 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7434	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL, VSE, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/3585	0.42	0/4839
1	B	0.25	0/3570	0.43	0/4816
All	All	0.24	0/7155	0.43	0/9655

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3518	0	3490	24	0
1	B	3504	0	3481	28	0
2	A	36	0	34	2	0
2	B	36	0	34	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	164	0	0	1	0
5	B	172	0	0	2	0
All	All	7434	0	7039	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:ASN:H	1:A:211:GLY:HA3	1.38	0.86
1:A:110:MET:HG2	1:A:124:SER:HA	1.76	0.68
1:B:191:ASP:OD1	1:B:192:TYR:N	2.30	0.65
1:B:193:ASP:HB3	1:B:196:GLN:HB2	1.78	0.65
1:A:301:SER:N	1:A:302:MET:HA	2.16	0.60
1:A:300:GLY:HA3	1:A:301:SER:HB3	1.84	0.59
1:B:257:LYS:NZ	5:B:852:HOH:O	2.34	0.59
1:A:377:ILE:HG23	1:A:382:TYR:HB3	1.83	0.59
1:B:181:THR:OG1	1:B:204:LYS:NZ	2.35	0.59
1:B:114:GLU:HG2	1:B:121:LYS:HB3	1.85	0.58
1:B:456:VAL:HG11	1:B:487:CYS:HB2	1.87	0.56
1:A:301:SER:O	1:A:301:SER:OG	2.20	0.55
1:A:528:GLU:OE2	5:A:832:HOH:O	2.18	0.55
1:A:210:ASN:N	1:A:211:GLY:HA3	2.16	0.53
1:B:310:GLU:OE2	1:B:409:ARG:NH2	2.41	0.53
1:B:213:PHE:N	1:B:221:PHE:O	2.42	0.53
1:B:110:MET:HB3	1:B:124:SER:HA	1.90	0.52
1:A:318:GLN:OE1	1:A:324:LYS:NZ	2.37	0.51
1:B:307:PHE:CE2	1:B:334:ILE:HG21	2.46	0.51
1:B:307:PHE:HA	1:B:410:VAL:HG21	1.93	0.50
1:B:423:LYS:HZ2	1:B:423:LYS:HB3	1.77	0.49
1:A:474:ALA:HB1	1:A:479:TYR:HB3	1.94	0.49
1:A:151:PHE:HB3	1:A:154:ILE:HG13	1.94	0.49
1:A:489:GLU:OE2	1:B:206:ARG:NH2	2.45	0.49
1:B:299:PRO:O	1:B:301:SER:N	2.47	0.48
1:A:231:TYR:HB2	1:A:242:LEU:HD22	1.96	0.48
1:A:308:LEU:HD21	1:A:330:THR:HG22	1.95	0.48
1:A:467:SER:H	1:A:470:GLU:HB3	1.79	0.47
1:B:377:ILE:HG23	1:B:382:TYR:HB3	1.97	0.47
1:B:170:LEU:O	1:B:243:SER:OG	2.31	0.47
1:A:300:GLY:HA3	1:A:301:SER:CB	2.45	0.46
1:A:151:PHE:CD1	1:A:174:MET:HB2	2.51	0.46
1:A:294:VAL:HG22	1:A:337:ILE:HG13	1.98	0.45
1:B:380:ARG:NH1	5:B:717:HOH:O	2.29	0.45
1:B:303:SER:HB2	1:B:306:ALA:HB3	1.99	0.45
1:B:442:LYS:NZ	1:B:506:ARG:O	2.36	0.44
1:B:232:LYS:HA	1:B:242:LEU:HB2	1.98	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:601:VSE:H36	2:A:601:VSE:H17	1.64	0.43
1:B:218:ARG:NH1	1:B:236:ASP:OD2	2.52	0.43
1:A:502:ARG:HD2	1:A:502:ARG:H	1.84	0.43
1:B:119:TRP:CD1	1:B:257:LYS:HG3	2.54	0.42
1:A:114:GLU:HB2	1:A:121:LYS:HB3	2.02	0.42
1:A:172:SER:HA	1:A:244:VAL:O	2.20	0.42
1:B:264:ARG:NH2	1:B:331:LYS:O	2.48	0.42
1:B:231:TYR:CE1	1:B:236:ASP:HB3	2.55	0.42
1:B:341:MET:HG3	1:B:393:LEU:HD23	2.02	0.41
1:B:257:LYS:H	1:B:257:LYS:HD2	1.85	0.41
1:A:243:SER:OG	1:A:244:VAL:N	2.53	0.41
1:A:378:GLU:HG3	1:A:441:ILE:HG12	2.03	0.41
1:B:225:GLN:HA	1:B:228:VAL:HG22	2.03	0.41
2:A:601:VSE:H1	2:A:601:VSE:CAP	2.34	0.40
1:A:502:ARG:HD3	1:A:505:GLU:HB2	2.04	0.40
1:B:297:MET:HE3	1:B:297:MET:HB3	1.98	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/454 (95%)	415 (96%)	14 (3%)	1 (0%)	52	57
1	B	425/454 (94%)	409 (96%)	14 (3%)	2 (0%)	34	33
All	All	855/908 (94%)	824 (96%)	28 (3%)	3 (0%)	39	40

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	300	GLY
1	B	485	GLU

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	425	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/393 (97%)	370 (97%)	11 (3%)	50	59
1	B	380/393 (97%)	364 (96%)	16 (4%)	36	42
All	All	761/786 (97%)	734 (96%)	27 (4%)	43	50

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	GLU
1	A	142	ASP
1	A	209	ASP
1	A	247	MET
1	A	301	SER
1	A	305	GLU
1	A	387	LEU
1	A	424	PHE
1	A	432	GLU
1	A	502	ARG
1	A	529	GLU
1	B	87	ILE
1	B	141	VAL
1	B	148	GLU
1	B	162	GLN
1	B	198	ASP
1	B	204	LYS
1	B	220	THR
1	B	257	LYS
1	B	302	MET
1	B	304	VAL
1	B	324	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	387	LEU
1	B	410	VAL
1	B	423	LYS
1	B	493	ASN
1	B	530	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	PTR	A	527	1,3	14,16,17	1.26	1 (7%)	18,22,24	0.74	1 (5%)
1	PTR	B	527	1,3	14,16,17	1.15	1 (7%)	18,22,24	0.74	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	527	1,3	-	0/9/11/13	0/1/1/1
1	PTR	B	527	1,3	-	0/9/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	527	PTR	OH-CZ	-4.14	1.30	1.40
1	B	527	PTR	OH-CZ	-4.08	1.30	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	527	PTR	O-C-CA	-2.13	119.95	125.49
1	A	527	PTR	O-C-CA	-2.10	120.02	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	VSE	A	601	-	39,41,41	0.74	1 (2%)	47,58,58	3.14	11 (23%)
2	VSE	B	601	-	39,41,41	0.80	1 (2%)	47,58,58	3.16	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VSE	A	601	-	-	0/12/36/36	0/6/6/6
2	VSE	B	601	-	-	0/12/36/36	0/6/6/6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	VSE	CAH-NAG	-3.34	1.33	1.38
2	A	601	VSE	CAH-NAG	-3.03	1.33	1.38

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	VSE	N3-C2-N1	-10.01	121.23	128.89
2	B	601	VSE	N3-C2-N1	-9.85	121.35	128.89
2	B	601	VSE	CAH-NAG-CAQ	-6.39	120.07	125.44
2	A	601	VSE	CAX-CAY-NAZ	-5.96	104.49	110.79
2	B	601	VSE	CAX-CAY-NAZ	-5.94	104.51	110.79
2	A	601	VSE	CAH-NAG-CAQ	-5.34	120.95	125.44
2	A	601	VSE	CAS-CAT-NAW	-3.63	104.39	112.57
2	B	601	VSE	CAS-CAT-NAW	-3.02	105.76	112.57
2	B	601	VSE	CAH-CAI-CAJ	-2.87	119.92	125.37
2	A	601	VSE	CAH-CAI-CAJ	-2.42	120.78	125.37
2	A	601	VSE	CAY-CAX-NAW	-2.35	105.28	110.79
2	A	601	VSE	CBB-NAW-CAT	2.83	117.91	112.37
2	B	601	VSE	CBB-NAW-CAT	3.26	118.76	112.37
2	A	601	VSE	CBC-NAZ-CAY	4.91	118.29	110.63
2	B	601	VSE	CBC-NAZ-CAY	4.95	118.34	110.63
2	A	601	VSE	CBC-NAZ-CBA	6.14	120.20	110.63
2	B	601	VSE	CBC-NAZ-CBA	6.44	120.67	110.63
2	B	601	VSE	CBA-NAZ-CAY	8.73	120.98	109.53
2	A	601	VSE	CBA-NAZ-CAY	9.12	121.50	109.53
2	A	601	VSE	CBB-NAW-CAX	9.54	126.53	109.23
2	B	601	VSE	CBB-NAW-CAX	9.68	126.78	109.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	VSE	2	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	434/454 (95%)	0.67	59 (13%) 4 5	28, 50, 85, 102	1 (0%)
1	B	431/454 (94%)	0.65	42 (9%) 10 11	25, 49, 88, 114	0
All	All	865/908 (95%)	0.66	101 (11%) 6 7	25, 50, 88, 114	1 (0%)

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	531	PRO	9.1
1	B	213	PHE	7.3
1	B	301	SER	6.6
1	A	209	ASP	6.5
1	B	234	GLY	6.0
1	A	301	SER	6.0
1	A	436	PHE	5.3
1	B	142	ASP	4.9
1	B	214	TYR	4.8
1	A	465	GLY	4.7
1	A	469	PRO	4.7
1	B	195	ARG	4.4
1	B	302	MET	4.4
1	A	305	GLU	4.3
1	B	530	ILE	4.2
1	A	473	ARG	4.2
1	A	477	ARG	3.9
1	A	208	LEU	3.9
1	A	353	ASP	3.8
1	A	448	PHE	3.8
1	A	424	PHE	3.6
1	A	387	LEU	3.5
1	A	435	ASN	3.5
1	A	304	VAL	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	452	LEU	3.4
1	A	211	GLY	3.4
1	A	438	SER	3.3
1	A	452	LEU	3.3
1	A	302	MET	3.3
1	B	233	LYS	3.2
1	B	424	PHE	3.2
1	B	465	GLY	3.1
1	A	192	TYR	3.1
1	B	139	ALA	3.1
1	A	451	LEU	3.0
1	B	207	THR	3.0
1	A	426	ILE	2.9
1	A	449	GLY	2.8
1	A	142	ASP	2.8
1	A	194	PRO	2.8
1	B	412	GLU	2.8
1	A	408	ALA	2.8
1	A	195	ARG	2.7
1	A	470	GLU	2.7
1	B	451	LEU	2.7
1	A	502	ARG	2.7
1	B	99	HIS	2.7
1	A	212	GLY	2.6
1	A	495	MET	2.6
1	B	469	PRO	2.6
1	A	352	SER	2.6
1	B	218	ARG	2.6
1	B	216	SER	2.6
1	A	258	ASP	2.5
1	B	448	PHE	2.5
1	A	450	ILE	2.5
1	A	467	SER	2.4
1	A	294	VAL	2.4
1	B	294	VAL	2.4
1	A	306	ALA	2.4
1	A	407	LEU	2.4
1	B	239	CYS	2.4
1	B	185	TYR	2.4
1	B	389	ALA	2.3
1	A	351	LYS	2.3
1	A	491	LEU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	336	ILE	2.3
1	B	402	ILE	2.3
1	A	357	LYS	2.3
1	B	151	PHE	2.3
1	B	423	LYS	2.3
1	B	409	ARG	2.2
1	A	210	ASN	2.2
1	A	428	TRP	2.2
1	B	248	SER	2.2
1	A	475	LEU	2.2
1	B	144	LEU	2.2
1	B	477	ARG	2.2
1	A	446	TRP	2.2
1	B	470	GLU	2.2
1	A	392	ILE	2.2
1	B	242	LEU	2.2
1	A	388	ARG	2.2
1	B	217	PRO	2.2
1	A	429	THR	2.1
1	A	447	SER	2.1
1	B	220	THR	2.1
1	B	336	ILE	2.1
1	A	300	GLY	2.1
1	A	485	GLU	2.1
1	A	367	SER	2.1
1	A	207	THR	2.1
1	B	504	GLU	2.1
1	B	473	ARG	2.1
1	A	86	ILE	2.1
1	A	504	GLU	2.1
1	B	241	LYS	2.1
1	B	236	ASP	2.0
1	A	385	ARG	2.0
1	A	193	ASP	2.0
1	A	356	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
1	PTR	B	527	16/17	0.96	0.14	-	45,58,64,67	0
1	PTR	A	527	16/17	0.97	0.12	-	24,36,44,46	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	B	602	1/1	0.15	0.37	6.11	74,74,74,74	0
3	CA	A	602	1/1	0.76	0.30	5.86	52,52,52,52	0
2	VSE	B	601	36/36	0.94	0.17	0.55	23,30,55,60	0
2	VSE	A	601	36/36	0.91	0.16	0.27	26,37,56,63	0
4	CL	B	603	1/1	0.98	0.08	-2.72	60,60,60,60	0
4	CL	A	603	1/1	0.95	0.07	-4.01	53,53,53,53	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.