



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:00 PM GMT

PDB ID : 3VTB
Title : Crystal structure of rat vitamin D receptor bound to a partial agonist 25-adenantyl-23-yne-19-norvitamin D ADTK1
Authors : Nakabayashi, M.; Kudo, T.; Tokiwa, H.; Makishima, M.; Yamada, S.; Ikura, T.; Ito, N.
Deposited on : 2012-05-26
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	
2	B	13	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2091 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vitamin D3 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	240	1924	1223	329	360	12	0	0	0

There are 57 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	106	GLY	-	EXPRESSION TAG	UNP P13053
A	107	SER	-	EXPRESSION TAG	UNP P13053
A	108	HIS	-	EXPRESSION TAG	UNP P13053
A	109	MET	-	EXPRESSION TAG	UNP P13053
A	110	GLY	-	EXPRESSION TAG	UNP P13053
A	111	SER	-	EXPRESSION TAG	UNP P13053
A	112	PRO	-	EXPRESSION TAG	UNP P13053
A	113	ASN	-	EXPRESSION TAG	UNP P13053
A	114	SER	-	EXPRESSION TAG	UNP P13053
A	115	PRO	-	EXPRESSION TAG	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	TYR	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	PRO	DELETION	UNP P13053
A	?	-	ARG	DELETION	UNP P13053
A	?	-	PRO	DELETION	UNP P13053
A	?	-	THR	DELETION	UNP P13053
A	?	-	LEU	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	PHE	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	GLY	DELETION	UNP P13053
A	?	-	ASN	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	ASP	DELETION	UNP P13053
A	?	-	LEU	DELETION	UNP P13053
A	?	-	TYR	DELETION	UNP P13053
A	?	-	THR	DELETION	UNP P13053
A	?	-	THR	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	LEU	DELETION	UNP P13053
A	?	-	ASP	DELETION	UNP P13053
A	?	-	MET	DELETION	UNP P13053
A	?	-	MET	DELETION	UNP P13053
A	?	-	GLU	DELETION	UNP P13053
A	?	-	PRO	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	GLY	DELETION	UNP P13053
A	?	-	PHE	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	ASN	DELETION	UNP P13053
A	?	-	LEU	DELETION	UNP P13053
A	?	-	ASP	DELETION	UNP P13053
A	?	-	LEU	DELETION	UNP P13053
A	?	-	ASN	DELETION	UNP P13053
A	?	-	GLY	DELETION	UNP P13053
A	?	-	GLU	DELETION	UNP P13053
A	?	-	ASP	DELETION	UNP P13053
A	?	-	SER	DELETION	UNP P13053
A	?	-	ASP	DELETION	UNP P13053
A	?	-	ASP	DELETION	UNP P13053
A	?	-	PRO	DELETION	UNP P13053

- Molecule 2 is a protein called COACTIVATOR PEPTIDE DRIP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	11	Total	C	N	O	S	0	0	0
			91	59	17	13	2			

- Molecule 3 is (1R,3R,7E,17BETA)-17-[(2R,6S)-6-HYDROXY-6-[(3S,5S,7S)-TRICYCLO[3.3.1.1^{3,7}]DEC-1-YL]HEX-4-YN-2-YL]-2-METHYLIDENE-9,10-SECOESTRA-5,7-DIENE-1,3-DIOL (three-letter code: TKA) (formula: C₃₅H₅₀O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C O	0	0
			38	35 3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	37	Total	O	0	0
			37	37		
4	B	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vitamin D3 receptor



- Molecule 2: COACTIVATOR PEPTIDE DRIP



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	153.56 Å 43.20 Å 42.49 Å 90.00° 95.56° 90.00°	Depositor
Resolution (Å)	25.24 – 2.00 25.24 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.0 (25.24-2.00) 89.8 (25.24-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.92 (at 1.99 Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.246 , 0.286 0.246 , 0.285	Depositor DCC
R_{free} test set	1678 reflections (9.82%)	DCC
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.733	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 62.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 18583 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2091	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TKA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1964	0.53	0/2656
2	B	0.33	0/92	0.50	0/121
All	All	0.34	0/2056	0.53	0/2777

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1924	0	1936	59	0
2	B	91	0	102	6	0
3	A	38	0	50	0	0
4	A	37	0	0	3	0
4	B	1	0	0	0	0
All	All	2091	0	2088	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:403:GLN:HG2	1:A:405:GLU:HG2	1.34	1.09
1:A:301:HIS:HA	1:A:396:GLN:HE22	1.31	0.95
1:A:337:PRO:HG2	1:A:349:GLU:HG3	1.48	0.95
1:A:123:LYS:N	1:A:365:CYS:HG	1.75	0.85
1:A:250:LEU:HD23	1:A:347:LEU:HD13	1.58	0.84
1:A:293:VAL:HG13	1:A:306:ILE:HG21	1.62	0.82
1:A:403:GLN:CG	1:A:405:GLU:HG2	2.16	0.74
1:A:340:PRO:HG2	4:A:627:HOH:O	1.85	0.73
1:A:242:LYS:HD3	2:B:634:LEU:HA	1.74	0.70
1:A:301:HIS:HA	1:A:396:GLN:NE2	2.07	0.66
1:A:223:LEU:HB3	1:A:224:PRO:HD3	1.77	0.65
1:A:374:GLN:HB2	1:A:378:LYS:NZ	2.11	0.65
1:A:405:GLU:HG3	1:A:406:ASN:N	2.12	0.63
1:A:360:GLN:O	1:A:364:ARG:HG2	1.98	0.63
1:A:125:SER:H	1:A:128:GLN:NE2	1.96	0.62
1:A:253:ASP:O	1:A:257:VAL:HG23	1.98	0.62
1:A:260:LYS:NZ	1:A:260:LYS:HB3	2.15	0.62
1:A:123:LYS:HD2	1:A:361:THR:HG21	1.83	0.60
1:A:238:ILE:HG12	1:A:259:LEU:HD13	1.84	0.59
1:A:123:LYS:HD2	1:A:128:GLN:HE22	1.69	0.57
1:A:335:VAL:O	1:A:352:GLN:HB2	2.05	0.56
1:A:151:ARG:HG2	1:A:151:ARG:HH11	1.71	0.56
1:A:281:SER:HB2	1:A:291:TYR:O	2.07	0.54
1:A:301:HIS:CE1	1:A:305:LEU:HD13	2.42	0.54
1:A:226:LEU:HD12	1:A:299:ALA:HB2	1.90	0.53
1:A:301:HIS:ND1	1:A:396:GLN:NE2	2.55	0.52
1:A:125:SER:H	1:A:128:GLN:HE21	1.55	0.52
1:A:127:GLU:O	1:A:131:ILE:HG13	2.10	0.52
1:A:363:ILE:HB	1:A:364:ARG:NH1	2.26	0.51
1:A:416:GLU:OE2	2:B:628:PRO:HD2	2.10	0.51
1:A:260:LYS:HE3	2:B:627:HIS:NE2	2.26	0.51
1:A:374:GLN:HB2	1:A:378:LYS:HZ1	1.75	0.50
1:A:293:VAL:HG23	4:A:622:HOH:O	2.10	0.50
1:A:373:HIS:O	1:A:374:GLN:HB2	2.11	0.50
2:B:625:LYS:HB2	2:B:627:HIS:CE1	2.47	0.49
1:A:344:ASP:CG	1:A:347:LEU:HB2	2.33	0.49
1:A:344:ASP:OD1	1:A:347:LEU:HD12	2.15	0.47
1:A:362:TYR:CE2	1:A:366:ARG:HG3	2.50	0.46
1:A:360:GLN:O	1:A:364:ARG:NH1	2.49	0.46
1:A:123:LYS:N	1:A:365:CYS:SG	2.85	0.45
2:B:626:ASN:O	2:B:628:PRO:HD3	2.17	0.45
1:A:259:LEU:HD11	2:B:634:LEU:HD21	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:TYR:HA	1:A:295:ASP:OD2	2.17	0.45
1:A:123:LYS:CD	1:A:128:GLN:HE22	2.28	0.45
1:A:123:LYS:HG2	1:A:124:LEU:N	2.31	0.44
1:A:277:MET:HA	1:A:280:MET:HE2	1.98	0.44
1:A:269:LEU:HD13	1:A:386:LEU:HD11	1.99	0.43
1:A:277:MET:HA	1:A:280:MET:CE	2.48	0.43
1:A:123:LYS:CG	1:A:124:LEU:N	2.81	0.43
1:A:260:LYS:NZ	1:A:260:LYS:CB	2.81	0.43
1:A:151:ARG:HG2	1:A:151:ARG:NH1	2.31	0.43
1:A:307:GLU:HB3	1:A:308:PRO:CD	2.49	0.43
1:A:335:VAL:HG12	1:A:335:VAL:O	2.19	0.43
1:A:339:ARG:O	1:A:342:VAL:HG23	2.18	0.42
1:A:130:HIS:O	1:A:134:ILE:HG12	2.19	0.42
1:A:382:LYS:HA	1:A:382:LYS:HD3	1.83	0.42
1:A:335:VAL:HG13	1:A:348:VAL:HG13	2.00	0.42
1:A:235:GLN:NE2	4:A:611:HOH:O	2.50	0.42
1:A:260:LYS:HZ3	1:A:260:LYS:HB3	1.85	0.41
1:A:307:GLU:HB3	1:A:308:PRO:HD3	2.03	0.41
1:A:247:PHE:CE2	1:A:255:GLN:HG2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/271 (87%)	224 (95%)	10 (4%)	2 (1%)	24	15
2	B	9/13 (69%)	8 (89%)	0	1 (11%)	0	0
All	All	245/284 (86%)	232 (95%)	10 (4%)	3 (1%)	16	8

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	626	ASN
1	A	219	PRO
1	A	124	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/247 (89%)	214 (97%)	6 (3%)	52	52
2	B	11/13 (85%)	11 (100%)	0	100	100
All	All	231/260 (89%)	225 (97%)	6 (3%)	54	54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	229	LEU
1	A	307	GLU
1	A	311	LYS
1	A	347	LEU
1	A	367	HIS
1	A	402	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	128	GLN
1	A	381	GLN
1	A	396	GLN
1	A	403	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TKA	A	500	-	38,43,43	2.38	22 (57%)	54,66,66	1.49	6 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TKA	A	500	-	-	0/18/92/92	0/3/6/6

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	TKA	C22-C23	-2.65	1.43	1.46
3	A	500	TKA	C30-C29	2.07	1.59	1.54
3	A	500	TKA	C16-C17	2.10	1.59	1.54
3	A	500	TKA	C12-C11	2.16	1.58	1.52
3	A	500	TKA	C04-C05	2.19	1.54	1.50
3	A	500	TKA	C32-C31	2.21	1.58	1.52
3	A	500	TKA	C09-C08	2.23	1.56	1.50
3	A	500	TKA	C28-C02	2.34	1.37	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	TKA	C20-C17	2.46	1.59	1.54
3	A	500	TKA	C37-C29	2.52	1.60	1.54
3	A	500	TKA	C14-C08	2.54	1.57	1.51
3	A	500	TKA	C36-C31	2.83	1.60	1.52
3	A	500	TKA	C36-C35	2.87	1.60	1.52
3	A	500	TKA	C38-C33	2.89	1.60	1.52
3	A	500	TKA	C12-C13	2.94	1.59	1.54
3	A	500	TKA	C10-C05	2.95	1.56	1.50
3	A	500	TKA	C32-C33	3.10	1.60	1.52
3	A	500	TKA	C38-C35	3.16	1.60	1.52
3	A	500	TKA	C01-C02	3.22	1.54	1.50
3	A	500	TKA	C34-C29	4.00	1.64	1.54
3	A	500	TKA	C18-C13	4.56	1.62	1.54
3	A	500	TKA	C03-C02	5.00	1.56	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	TKA	C20-C22-C23	-5.84	105.90	113.68
3	A	500	TKA	C13-C14-C08	-3.96	105.75	112.85
3	A	500	TKA	C14-C13-C17	-2.75	96.88	99.74
3	A	500	TKA	C30-C29-C25	-2.32	104.87	110.31
3	A	500	TKA	C11-C12-C13	-2.05	110.51	113.17
3	A	500	TKA	C18-C13-C14	3.73	116.50	110.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/271 (88%)	0.61	28 (11%) 6 7	32, 51, 70, 88	0
2	B	11/13 (84%)	0.98	3 (27%) 1 1	50, 60, 82, 83	0
All	All	251/284 (88%)	0.62	31 (12%) 5 6	32, 51, 72, 88	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	159	MET	7.7
1	A	218	SER	6.5
1	A	420	ASN	5.9
1	A	402	PHE	5.8
1	A	230	VAL	4.5
1	A	123	LYS	4.0
2	B	625	LYS	3.5
1	A	126	GLU	3.5
1	A	219	PRO	3.3
1	A	264	ILE	3.2
1	A	346	LYS	3.0
1	A	267	ILE	2.9
1	A	266	VAL	2.9
1	A	334	ILE	2.9
1	A	332	ILE	2.7
1	A	368	PRO	2.7
2	B	626	ASN	2.7
1	A	330	MET	2.7
1	A	227	ALA	2.6
1	A	419	GLY	2.6
1	A	248	ARG	2.5
1	A	403	GLN	2.4
1	A	269	LEU	2.3
1	A	328	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	229	LEU	2.3
1	A	331	ALA	2.2
1	A	287	GLN	2.2
2	B	635	LYS	2.2
1	A	157	VAL	2.2
1	A	398	ARG	2.2
1	A	226	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	TKA	A	500	38/38	0.92	0.27	1.76	31,38,44,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.