



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:02 PM GMT

PDB ID : 3VUM
Title : Crystal structure of a cysteine-deficient mutant M7 in MAP kinase JNK1
Authors : Nakaniwa, T.; Kinoshita, T.; Inoue, T.
Deposited on : 2012-07-02
Resolution : 2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	<div> <div>5%</div> <div>56%</div> <div>29%</div> <div>6%</div> <div>6%</div> </div>
2	F	11	<div> <div>55%</div> <div>27%</div> <div>9%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	TRS	A	402	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	347	2806	1802	478	514	12	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	VAL	CYS	ENGINEERED MUTATION	UNP A1L4K2
A	79	VAL	CYS	ENGINEERED MUTATION	UNP A1L4K2
A	116	SER	CYS	ENGINEERED MUTATION	UNP A1L4K2
A	137	VAL	CYS	ENGINEERED MUTATION	UNP A1L4K2
A	163	ALA	CYS	ENGINEERED MUTATION	UNP A1L4K2
A	213	VAL	CYS	ENGINEERED MUTATION	UNP A1L4K2
A	245	SER	CYS	ENGINEERED MUTATION	UNP A1L4K2
A	365	HIS	-	EXPRESSION TAG	UNP A1L4K2
A	366	HIS	-	EXPRESSION TAG	UNP A1L4K2
A	367	HIS	-	EXPRESSION TAG	UNP A1L4K2
A	368	HIS	-	EXPRESSION TAG	UNP A1L4K2
A	369	HIS	-	EXPRESSION TAG	UNP A1L4K2
A	370	HIS	-	EXPRESSION TAG	UNP A1L4K2

- Molecule 2 is a protein called Peptide from C-Jun-amino-terminal kinase-interacting protein 1.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	F	11	95	61	19	15	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	78	Total 78	O 78	0	0
5	F	6	Total 6	O 6	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 8



- Molecule 2: Peptide from C-Jun-amino-terminal kinase-interacting protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	161.88Å 161.88Å 87.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.85 – 2.69 33.85 – 2.69	Depositor EDS
% Data completeness (in resolution range)	99.8 (33.85-2.69) 99.9 (33.85-2.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.64 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.188 , 0.255 0.192 , 0.253	Depositor DCC
R_{free} test set	990 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	54.3	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 19294 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2998	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	3/2870 (0.1%)	1.01	4/3884 (0.1%)
2	F	0.96	0/97	1.29	1/129 (0.8%)
All	All	0.89	3/2967 (0.1%)	1.02	5/4013 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	234	TRP	CD2-CE2	6.49	1.49	1.41
1	A	352	TRP	CD2-CE2	5.85	1.48	1.41
1	A	209	TRP	CD2-CE2	5.11	1.47	1.41

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	168	LEU	CA-CB-CG	10.68	139.85	115.30
2	F	553	ARG	C-N-CD	5.70	140.37	128.40
1	A	41	VAL	CB-CA-C	-5.67	100.63	111.40
1	A	135	MET	CG-SD-CE	-5.62	91.21	100.20
1	A	50	ARG	NE-CZ-NH1	-5.25	117.67	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	339	ASP	Peptide
1	A	49	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2806	0	2815	115	0
2	F	95	0	103	15	0
3	A	5	0	0	0	0
4	A	8	0	12	1	0
5	A	78	0	0	5	0
5	F	6	0	0	1	0
All	All	2998	0	2930	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

All (128) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:553:ARG:HG2	2:F:554:PRO:HD3	1.23	1.11
1:A:233:GLN:HG3	1:A:234:TRP:N	1.68	1.09
1:A:82:HIS:HD2	1:A:84:ASN:H	1.12	0.98
1:A:29:LEU:HB3	1:A:41:VAL:HG11	1.48	0.95
1:A:127:ARG:NH1	2:F:559:THR:O	2.00	0.94
1:A:30:LYS:O	1:A:41:VAL:HG13	1.67	0.94
1:A:233:GLN:HG3	1:A:234:TRP:H	1.30	0.90
2:F:553:ARG:HG2	2:F:554:PRO:CD	2.02	0.89
1:A:169:ASP:O	1:A:170:PHE:CD1	2.28	0.87
1:A:169:ASP:O	1:A:170:PHE:HD1	1.57	0.86
2:F:553:ARG:N	2:F:553:ARG:HE	1.75	0.83
1:A:318:HIS:HD2	1:A:320:TYR:H	1.21	0.83
1:A:170:PHE:HB3	1:A:173:ALA:HB3	1.58	0.82
1:A:172:LEU:HB3	5:A:524:HOH:O	1.79	0.82
2:F:553:ARG:NE	2:F:553:ARG:N	2.26	0.82
1:A:259:TYR:HD2	1:A:259:TYR:C	1.83	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:LYS:H	1:A:156:ASN:HD21	1.29	0.79
1:A:170:PHE:HA	5:A:534:HOH:O	1.81	0.79
1:A:149:HIS:HD2	1:A:151:ASP:H	1.31	0.79
1:A:338:PRO:O	1:A:340:LYS:N	2.16	0.77
1:A:259:TYR:HE2	1:A:263:ARG:HD2	1.47	0.77
1:A:20:PHE:CZ	1:A:41:VAL:HG23	2.19	0.76
1:A:82:HIS:HD2	1:A:84:ASN:N	1.84	0.75
1:A:318:HIS:CD2	1:A:320:TYR:H	2.04	0.75
1:A:259:TYR:CD2	1:A:259:TYR:C	2.58	0.74
1:A:343:ASP:HA	1:A:345:ARG:HD2	1.70	0.74
1:A:72:ARG:HD3	1:A:173:ALA:O	1.86	0.74
2:F:553:ARG:CG	2:F:554:PRO:HD3	2.11	0.73
1:A:170:PHE:CB	1:A:173:ALA:HB3	2.19	0.71
1:A:41:VAL:HG12	1:A:42:ALA:O	1.90	0.71
1:A:8:ASN:O	1:A:9:ASN:HB2	1.91	0.70
1:A:82:HIS:CD2	1:A:84:ASN:H	2.03	0.69
1:A:23:LEU:HD13	1:A:367:HIS:HA	1.74	0.69
1:A:174:ARG:O	1:A:174:ARG:HD3	1.92	0.69
1:A:141:HIS:CE1	1:A:335:PRO:HD3	2.29	0.68
1:A:208:ILE:HD11	1:A:312:VAL:HG12	1.75	0.67
1:A:338:PRO:C	1:A:340:LYS:H	1.98	0.67
1:A:287:ASN:HB2	5:A:537:HOH:O	1.95	0.66
1:A:170:PHE:HB3	1:A:173:ALA:CB	2.25	0.66
1:A:29:LEU:HD22	1:A:43:ALA:HB2	1.78	0.65
1:A:30:LYS:O	1:A:41:VAL:CG1	2.44	0.64
1:A:296:ASP:OD2	1:A:318:HIS:HE1	1.81	0.64
1:A:169:ASP:C	1:A:170:PHE:CD1	2.71	0.64
1:A:259:TYR:CE2	1:A:263:ARG:HD2	2.31	0.63
2:F:553:ARG:CG	2:F:554:PRO:CD	2.72	0.63
1:A:119:ILE:HD13	1:A:217:GLU:HG2	1.82	0.61
1:A:25:ARG:NH2	1:A:48:LEU:HD22	2.15	0.61
2:F:553:ARG:CD	2:F:554:PRO:HD2	2.31	0.61
1:A:247:GLU:CD	1:A:247:GLU:H	2.05	0.60
2:F:553:ARG:HE	2:F:553:ARG:CA	2.16	0.59
1:A:348:THR:HG22	1:A:351:GLU:HG3	1.85	0.59
1:A:76:LEU:HB2	1:A:170:PHE:HD2	1.68	0.58
1:A:56:LYS:HE2	1:A:58:SER:OG	2.04	0.57
1:A:153:LYS:H	1:A:156:ASN:ND2	2.01	0.57
1:A:20:PHE:CZ	1:A:41:VAL:CG2	2.87	0.56
2:F:555:LYS:NZ	5:F:606:HOH:O	2.38	0.56
1:A:139:ILE:HD11	1:A:152:LEU:CD1	2.36	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:553:ARG:CD	2:F:554:PRO:CD	2.84	0.55
1:A:236:LYS:O	1:A:240:GLN:HB2	2.07	0.55
1:A:233:GLN:CG	1:A:234:TRP:N	2.52	0.54
1:A:259:TYR:O	1:A:259:TYR:HD2	1.91	0.54
1:A:168:LEU:O	1:A:169:ASP:HB2	2.07	0.54
1:A:8:ASN:O	1:A:8:ASN:CG	2.46	0.53
1:A:240:GLN:HG3	1:A:274:LEU:CD2	2.38	0.53
1:A:59:ARG:NH1	1:A:102:GLN:OE1	2.41	0.53
1:A:257:ARG:HG3	1:A:261:GLU:HG3	1.91	0.53
1:A:118:VAL:HA	1:A:121:MET:HE2	1.90	0.53
1:A:149:HIS:HD2	1:A:151:ASP:N	2.03	0.53
1:A:41:VAL:HG12	1:A:42:ALA:N	2.24	0.52
1:A:145:ALA:HB3	1:A:147:ILE:HD12	1.92	0.52
1:A:338:PRO:C	1:A:340:LYS:N	2.61	0.51
1:A:82:HIS:CD2	1:A:83:LYS:N	2.80	0.50
1:A:95:GLN:HG2	1:A:100:GLU:O	2.11	0.50
1:A:118:VAL:HA	1:A:121:MET:CE	2.42	0.49
1:A:76:LEU:HD12	1:A:170:PHE:HB2	1.94	0.49
1:A:234:TRP:O	1:A:237:VAL:HB	2.13	0.48
1:A:147:ILE:HD11	1:A:337:ILE:HD11	1.95	0.48
1:A:72:ARG:O	1:A:76:LEU:HG	2.13	0.48
1:A:365:HIS:O	1:A:366:HIS:HB2	2.12	0.48
1:A:149:HIS:CD2	1:A:151:ASP:H	2.21	0.48
1:A:228:THR:O	1:A:229:ASP:HB3	2.12	0.47
1:A:90:ASN:HD22	1:A:91:VAL:N	2.12	0.47
1:A:300:LYS:HB3	1:A:310:ILE:HD13	1.97	0.47
1:A:25:ARG:HH21	1:A:48:LEU:HD22	1.80	0.47
1:A:10:PHE:O	1:A:24:LYS:HE3	2.14	0.47
1:A:174:ARG:C	1:A:174:ARG:HD3	2.35	0.47
1:A:189:ARG:HH12	1:A:231:ILE:HD11	1.79	0.47
1:A:240:GLN:HE21	1:A:274:LEU:HD23	1.80	0.46
1:A:58:SER:O	1:A:59:ARG:C	2.52	0.46
1:A:202:TYR:CD1	1:A:202:TYR:C	2.89	0.46
1:A:341:GLN:HG2	1:A:343:ASP:HB3	1.98	0.46
1:A:160:LYS:HB3	1:A:162:ASP:OD1	2.16	0.46
1:A:245:SER:HB3	4:A:402:TRS:N	2.31	0.45
1:A:132:LEU:HD21	1:A:215:MET:HG3	1.98	0.45
2:F:553:ARG:HD2	2:F:554:PRO:HD2	1.98	0.45
1:A:32:ILE:HD13	1:A:42:ALA:HB2	1.98	0.45
1:A:312:VAL:HG13	5:A:507:HOH:O	2.15	0.45
1:A:198:LEU:HB2	1:A:200:MET:HG3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:553:ARG:NE	2:F:554:PRO:HD2	2.32	0.45
1:A:194:PRO:HA	1:A:197:ILE:HG12	1.98	0.45
1:A:59:ARG:HA	1:A:102:GLN:O	2.17	0.44
1:A:139:ILE:HD11	1:A:152:LEU:HD11	1.97	0.44
1:A:130:TYR:CD2	2:F:557:PRO:HD2	2.52	0.44
1:A:98:LEU:HD12	1:A:353:LYS:HD3	1.99	0.44
1:A:195:GLU:HA	1:A:200:MET:HE3	2.00	0.44
2:F:553:ARG:NE	2:F:553:ARG:CA	2.77	0.44
1:A:364:GLU:O	1:A:365:HIS:HB2	2.17	0.43
1:A:202:TYR:O	1:A:202:TYR:CD1	2.71	0.43
1:A:118:VAL:O	1:A:121:MET:HG2	2.19	0.43
1:A:229:ASP:OD1	1:A:232:ASP:HB2	2.19	0.43
1:A:87:GLY:HA3	1:A:109:GLU:OE2	2.18	0.43
1:A:90:ASN:HD22	1:A:91:VAL:H	1.67	0.42
1:A:280:PHE:HB3	1:A:281:PRO:HD2	2.02	0.42
1:A:8:ASN:O	1:A:9:ASN:CB	2.62	0.42
1:A:358:LYS:O	1:A:362:ASP:HB2	2.19	0.42
1:A:240:GLN:HE21	1:A:274:LEU:CD2	2.34	0.41
1:A:318:HIS:CG	1:A:319:PRO:HD2	2.55	0.41
1:A:156:ASN:HD22	1:A:156:ASN:C	2.23	0.41
1:A:170:PHE:N	1:A:170:PHE:CD1	2.87	0.41
1:A:348:THR:O	1:A:349:ILE:C	2.58	0.41
1:A:231:ILE:HG22	1:A:231:ILE:O	2.21	0.41
1:A:318:HIS:CD2	1:A:319:PRO:HD2	2.56	0.41
1:A:326:ASP:C	1:A:326:ASP:OD1	2.59	0.40
1:A:30:LYS:HD2	1:A:30:LYS:HA	1.85	0.40
1:A:29:LEU:HD22	1:A:43:ALA:CB	2.50	0.40
1:A:318:HIS:HA	1:A:319:PRO:HD3	1.91	0.40
1:A:286:HIS:HB3	5:A:560:HOH:O	2.20	0.40
1:A:363:LEU:HD13	1:A:363:LEU:HA	1.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/370 (93%)	305 (89%)	24 (7%)	14 (4%)	3	7
2	F	9/11 (82%)	9 (100%)	0	0	100	100
All	All	352/381 (92%)	314 (89%)	24 (7%)	14 (4%)	4	8

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	ASP
1	A	263	ARG
1	A	339	ASP
1	A	365	HIS
1	A	366	HIS
1	A	35	GLY
1	A	349	ILE
1	A	233	GLN
1	A	36	ALA
1	A	343	ASP
1	A	9	ASN
1	A	169	ASP
1	A	197	ILE
1	A	227	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/331 (93%)	261 (85%)	47 (15%)	3	8
2	F	11/11 (100%)	9 (82%)	2 (18%)	2	5
All	All	319/342 (93%)	270 (85%)	49 (15%)	3	8

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ASP
1	A	8	ASN
1	A	13	VAL
1	A	37	GLN
1	A	48	LEU
1	A	50	ARG
1	A	63	ASN
1	A	74	LEU
1	A	79	VAL
1	A	90	ASN
1	A	91	VAL
1	A	115	LEU
1	A	116	SER
1	A	117	GLN
1	A	122	GLU
1	A	132	LEU
1	A	142	LEU
1	A	150	ARG
1	A	156	ASN
1	A	168	LEU
1	A	170	PHE
1	A	174	ARG
1	A	189	ARG
1	A	229	ASP
1	A	230	HIS
1	A	233	GLN
1	A	250	LYS
1	A	256	VAL
1	A	257	ARG
1	A	259	TYR
1	A	263	ARG
1	A	265	LYS
1	A	285	GLU
1	A	287	ASN
1	A	289	LEU
1	A	302	LEU
1	A	310	ILE
1	A	322	ASN
1	A	336	LYS
1	A	338	PRO
1	A	341	GLN
1	A	343	ASP
1	A	345	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	349	ILE
1	A	363	LEU
1	A	365	HIS
1	A	366	HIS
2	F	553	ARG
2	F	555	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	37	GLN
1	A	64	GLN
1	A	66	HIS
1	A	82	HIS
1	A	90	ASN
1	A	141	HIS
1	A	149	HIS
1	A	156	ASN
1	A	240	GLN
1	A	286	HIS
1	A	318	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	401	-	4,4,4	0.90	0	6,6,6	0.55	0
4	TRS	A	402	-	7,7,7	1.16	0	9,9,9	2.82	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	401	-	-	0/0/0/0	0/0/0/0
4	TRS	A	402	-	-	0/9/9/9	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	402	TRS	C1-C-N	-2.59	103.37	108.09
4	A	402	TRS	C3-C-N	2.60	112.83	108.09
4	A	402	TRS	O2-C2-C	4.46	120.22	111.18
4	A	402	TRS	O3-C3-C	5.91	123.13	111.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	402	TRS	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/370 (93%)	-0.12	17 (4%) 33 32	28, 48, 106, 151	0
2	F	11/11 (100%)	-0.33	0 100 100	28, 37, 60, 70	0
All	All	358/381 (93%)	-0.12	17 (4%) 35 34	28, 48, 106, 151	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	ASP	6.7
1	A	342	LEU	3.6
1	A	228	THR	3.5
1	A	340	LYS	3.4
1	A	229	ASP	3.4
1	A	367	HIS	3.3
1	A	343	ASP	3.3
1	A	344	GLU	3.3
1	A	227	GLY	3.0
1	A	366	HIS	2.9
1	A	253	GLN	2.8
1	A	339	ASP	2.7
1	A	250	LYS	2.5
1	A	254	PRO	2.3
1	A	230	HIS	2.3
1	A	287	ASN	2.1
1	A	286	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	TRS	A	402	8/8	0.75	0.25	2.80	60,73,84,87	0
3	SO4	A	401	5/5	0.97	0.09	-9.33	60,65,69,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.