



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:02 PM GMT

PDB ID : 3VV9
Title : Crystal structure of cyanide-insensitive alternative oxidase from *Trypanosoma brucei*
Authors : Shiba, T.; Kido, Y.; Sakamoto, K.; Inaoka, D.K.; Tsuge, C.; Tatsumi, R.; Balogun, E.O.; Nara, T.; Aoki, T.; Honma, T.; Tanaka, A.; Inoue, M.; Matsuoka, S.; Saimoto, H.; Moore, A.L.; Harada, S.; Kita, K.
Deposited on : 2012-07-17
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2228 (2.90-2.82)
Clashscore	102246	2499 (2.90-2.82)
Ramachandran outliers	100387	2439 (2.90-2.82)
Sidechain outliers	100360	2442 (2.90-2.82)
RSRZ outliers	91569	2236 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	 2% 37% 37% 7% • 19%
1	B	329	 4% 39% 35% 7% • 19%
1	C	329	 3% 35% 36% 10% • 19%
1	D	329	 4% 38% 34% 9% 19%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OH	B	503	-	-	-	X
3	OH	C	503	-	-	-	X
3	OH	D	503	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alternative oxidase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2170	1390	386	384	10			
1	B	268	Total	C	N	O	S	0	0	0
			2179	1395	387	387	10			
1	C	266	Total	C	N	O	S	0	0	0
			2163	1385	385	383	10			
1	D	266	Total	C	N	O	S	0	0	0
			2163	1385	385	383	10			

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Fe	0	0
			2	2		
2	A	2	Total	Fe	0	0
			2	2		
2	D	2	Total	Fe	0	0
			2	2		
2	C	2	Total	Fe	0	0
			2	2		

- Molecule 3 is HYDROXIDE ION (three-letter code: OH) (formula: HO).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0
3	B	1	Total O 1 1	0	0
3	C	1	Total O 1 1	0	0
3	D	1	Total O 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	14	Total O 14 14	0	0
4	B	8	Total O 8 8	0	0
4	C	7	Total O 7 7	0	0
4	D	2	Total O 2 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alternative oxidase, mitochondrial



- Molecule 1: Alternative oxidase, mitochondrial



- Molecule 1: Alternative oxidase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	261.31Å 63.12Å 136.46Å 90.00° 121.38° 90.00°	Depositor
Resolution (Å)	45.38 – 2.85 45.38 – 2.85	Depositor EDS
% Data completeness (in resolution range)	87.6 (45.38-2.85) 90.4 (45.38-2.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.86Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.192 , 0.247 0.191 , 0.232	Depositor DCC
R_{free} test set	1990 reflections (5.14%)	DCC
Wilson B-factor (Å ²)	86.9	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 79.9	EDS
Estimated twinning fraction	0.524 for H, K, L 0.476 for H+4/2L, -K, -L 0.157 for -h-2*k,l,-k,l	Xtriage
Reported twinning fraction	0.524 for H, K, L 0.476 for H+4/2L, -K, -L	Depositor
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	1 of 40879 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8718	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, OH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2220	0.78	1/3015 (0.0%)
1	B	0.53	0/2229	0.79	3/3028 (0.1%)
1	C	0.47	0/2212	0.73	0/3004
1	D	0.50	0/2212	0.75	0/3004
All	All	0.50	0/8873	0.76	4/12051 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	278	ASP	CB-CG-OD1	5.57	123.31	118.30
1	B	52	LEU	CA-CB-CG	5.46	127.85	115.30
1	A	156	LEU	CA-CB-CG	5.19	127.25	115.30
1	B	108	GLY	N-CA-C	-5.10	100.34	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2170	0	2202	143	0
1	B	2179	0	2207	143	0
1	C	2163	0	2194	113	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2163	0	2194	122	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	14	0	0	3	0
4	B	8	0	0	0	0
4	C	7	0	0	2	0
4	D	2	0	0	0	0
All	All	8718	0	8797	473	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 473 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:163:ARG:O	1:C:167:MET:HG2	1.46	1.13
1:D:295:LYS:NZ	1:D:295:LYS:HB2	1.62	1.12
1:B:297:ASN:HB2	1:B:298:PRO:CD	1.86	1.05
1:A:170:ILE:HG13	1:B:145:MET:HB3	1.05	1.04
1:D:295:LYS:HZ3	1:D:295:LYS:HB2	1.20	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/329 (80%)	203 (77%)	48 (18%)	14 (5%)	2	7
1	B	266/329 (81%)	208 (78%)	50 (19%)	8 (3%)	5	19
1	C	264/329 (80%)	201 (76%)	48 (18%)	15 (6%)	2	5
1	D	264/329 (80%)	206 (78%)	44 (17%)	14 (5%)	2	7
All	All	1059/1316 (80%)	818 (77%)	190 (18%)	51 (5%)	3	9

5 of 51 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	71	GLU
1	B	297	ASN
1	C	110	ILE
1	D	103	SER
1	A	105	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/292 (83%)	206 (85%)	36 (15%)	4	10
1	B	243/292 (83%)	208 (86%)	35 (14%)	4	10
1	C	241/292 (82%)	192 (80%)	49 (20%)	1	3
1	D	241/292 (82%)	204 (85%)	37 (15%)	3	9
All	All	967/1168 (83%)	810 (84%)	157 (16%)	3	7

5 of 157 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	32	VAL
1	C	103	SER
1	D	242	VAL
1	C	44	LEU
1	C	83	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	260	ASN
1	B	285	GLN
1	D	269	HIS
1	B	273	ASN
1	B	274	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are modelled with single atom and 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/329 (81%)	-0.09	6 (2%) 65 61	54, 76, 104, 132	0
1	B	268/329 (81%)	-0.09	13 (4%) 33 27	53, 75, 108, 152	0
1	C	266/329 (80%)	0.10	11 (4%) 41 34	61, 90, 137, 175	0
1	D	266/329 (80%)	0.06	13 (4%) 33 27	54, 81, 123, 157	0
All	All	1067/1316 (81%)	-0.01	43 (4%) 42 34	53, 80, 119, 175	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	109	SER	6.1
1	C	110	ILE	4.8
1	C	235	THR	4.6
1	A	52	LEU	4.6
1	C	175	PRO	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	OH	B	503	1/1	0.97	0.38	8.19	40,40,40,40	0
3	OH	D	503	1/1	0.99	0.31	4.95	79,79,79,79	0
3	OH	C	503	1/1	0.99	0.28	3.93	36,36,36,36	0
2	FE	C	502	1/1	0.93	0.21	1.00	71,71,71,71	0
2	FE	A	502	1/1	0.89	0.19	0.59	70,70,70,70	0
2	FE	B	501	1/1	0.85	0.18	0.59	78,78,78,78	0
2	FE	D	501	1/1	0.97	0.20	0.51	84,84,84,84	0
2	FE	B	502	1/1	0.90	0.17	0.44	64,64,64,64	0
2	FE	C	501	1/1	0.99	0.18	-0.49	79,79,79,79	0
3	OH	A	503	1/1	0.95	0.15	-0.54	43,43,43,43	0
2	FE	D	502	1/1	0.96	0.17	-0.63	59,59,59,59	0
2	FE	A	501	1/1	0.92	0.10	-2.73	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.