



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:03 PM GMT

PDB ID : 3VZB
Title : Crystal structure of Sphingosine Kinase 1
Authors : Min, X.; Walker, N.P.; Wang, Z.
Deposited on : 2012-10-10
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	<div> <div>2%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>
1	B	360	<div> <div>6%</div> <div>77%</div> <div>15%</div> <div>6%</div> </div>
1	C	360	<div> <div>7%</div> <div>83%</div> <div>11%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SQS	A	401	-	-	-	X
2	SQS	B	401	-	-	-	X
4	EDO	A	403	-	-	-	X
4	EDO	A	404	-	-	-	X
4	EDO	A	406	-	-	-	X
4	EDO	A	407	-	-	-	X
4	EDO	A	408	-	-	-	X
4	EDO	B	403	-	-	-	X
4	EDO	B	404	-	-	-	X
4	EDO	B	406	-	-	-	X
4	EDO	B	407	-	-	-	X
4	EDO	C	402	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sphingosine kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	1	0
			2803	1793	498	491	21			
1	B	339	Total	C	N	O	S	0	1	0
			2659	1705	468	466	20			
1	C	344	Total	C	N	O	S	0	1	0
			2700	1728	477	474	21			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	ALA	-	EXPRESSION TAG	UNP Q9NYA1
A	6	MET	-	EXPRESSION TAG	UNP Q9NYA1
A	7	GLY	-	EXPRESSION TAG	UNP Q9NYA1
A	8	SER	-	EXPRESSION TAG	UNP Q9NYA1
B	5	ALA	-	EXPRESSION TAG	UNP Q9NYA1
B	6	MET	-	EXPRESSION TAG	UNP Q9NYA1
B	7	GLY	-	EXPRESSION TAG	UNP Q9NYA1
B	8	SER	-	EXPRESSION TAG	UNP Q9NYA1
C	5	ALA	-	EXPRESSION TAG	UNP Q9NYA1
C	6	MET	-	EXPRESSION TAG	UNP Q9NYA1
C	7	GLY	-	EXPRESSION TAG	UNP Q9NYA1
C	8	SER	-	EXPRESSION TAG	UNP Q9NYA1

- Molecule 2 is (2S,3R,4E)-2-AMINOOCCTADEC-4-ENE-1,3-DIOL (three-letter code: SQS) (formula: C₁₈H₃₇NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			21	18	1	2		
2	B	1	Total	C	N	O	0	0
			21	18	1	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	182	Total	O	0	0
			182	182		
5	B	120	Total	O	0	0
			120	120		
5	C	82	Total	O	0	0
			82	82		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Sphingosine kinase 1



• Molecule 1: Sphingosine kinase 1



• Molecule 1: Sphingosine kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	101.86 Å 226.26 Å 106.49 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.22 – 2.00 48.17 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.22-2.00) 99.7 (48.17-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.213 , 0.262 0.216 , 0.262	Depositor DCC
R_{free} test set	4156 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 82864 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8663	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SQS, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/2871	0.70	2/3896 (0.1%)
1	B	0.44	0/2722	0.69	3/3691 (0.1%)
1	C	0.43	0/2764	0.66	0/3747
All	All	0.44	0/8357	0.68	5/11334 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	294	VAL	CB-CA-C	-5.94	100.12	111.40
1	B	123	TYR	N-CA-CB	5.55	120.59	110.60
1	A	10	VAL	CB-CA-C	-5.44	101.06	111.40
1	B	35	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	B	35	ARG	NE-CZ-NH1	5.16	122.88	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	362	VAL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2803	0	2847	22	0
1	B	2659	0	2686	25	0
1	C	2700	0	2726	15	0
2	A	21	0	37	0	0
2	B	21	0	37	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	1	0
4	A	24	0	36	6	0
4	B	20	0	30	2	0
4	C	16	0	24	0	0
5	A	182	0	0	1	1
5	B	120	0	0	1	0
5	C	82	0	0	0	0
All	All	8663	0	8423	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:SER:O	1:B:122:HIS:O	1.61	1.17
1:B:111:GLY:O	1:B:114:ASN:ND2	2.10	0.85
1:A:362:VAL:HG22	4:A:403:EDO:H11	1.59	0.84
1:B:169:LEU:HD12	1:B:340:VAL:HG22	1.67	0.76
1:A:362:VAL:HG22	4:A:403:EDO:C1	2.19	0.73
1:B:89:ASN:HD21	1:B:164:PHE:H	1.37	0.71
1:B:336:GLY:N	1:B:347:SER:O	2.26	0.68
1:A:68:GLU:O	1:A:71:ARG:HG3	1.96	0.65
1:A:337:VAL:HG13	4:A:406:EDO:H12	1.80	0.64
1:C:207:ARG:HA	1:C:255:GLU:O	2.02	0.60
1:B:203:LEU:HD21	1:B:299:LEU:HD22	1.82	0.60
1:B:120:LEU:HD22	1:B:363:SER:HB3	1.84	0.59
1:C:258:VAL:HG22	1:C:290:VAL:HB	1.84	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:79:SER:HB3	1:C:83:LEU:HD12	1.86	0.57
1:B:142:LEU:HA	1:B:362:VAL:HG21	1.87	0.57
1:A:111:GLY:O	1:A:114:ASN:ND2	2.37	0.56
1:A:294:VAL:HG13	1:A:318:TYR:HB2	1.88	0.55
1:A:258:VAL:HG22	1:A:290:VAL:HB	1.88	0.55
1:A:126:TYR:OH	1:A:144:ARG:NH2	2.41	0.53
1:A:207:ARG:NH1	1:A:333:ASP:OD2	2.43	0.52
1:B:61:ARG:NH1	1:B:86[A]:GLU:OE2	2.42	0.52
1:B:52:MET:SD	1:B:63:LEU:HD13	2.50	0.52
1:B:122:HIS:O	1:B:123:TYR:CG	2.63	0.52
1:C:106:CYS:HB2	1:C:362:VAL:CG1	2.41	0.51
1:A:106:CYS:HB2	1:A:362:VAL:HG13	1.93	0.50
1:A:10:VAL:HG13	1:A:146:LEU:HD23	1.94	0.49
1:C:77:VAL:HG13	1:C:83:LEU:HB3	1.94	0.49
1:C:76:VAL:HB	1:C:106:CYS:HB3	1.95	0.49
1:B:193:THR:HG22	1:B:197:PHE:CE1	2.46	0.49
1:B:22:ASN:HD22	1:B:22:ASN:C	2.16	0.49
1:A:77:VAL:HG13	1:A:83:LEU:HD13	1.95	0.49
1:B:242:GLU:OE1	1:B:242:GLU:N	2.45	0.49
1:A:142:LEU:HA	1:A:362:VAL:HG21	1.94	0.48
1:B:329:LEU:HD23	1:B:338:PHE:HE2	1.78	0.48
1:C:77:VAL:CG1	1:C:83:LEU:HB3	2.44	0.48
1:B:54:THR:O	1:B:54:THR:HG23	2.14	0.47
1:C:79:SER:CB	1:C:83:LEU:HD12	2.45	0.47
1:B:153:LEU:HB3	1:B:165:SER:HB3	1.96	0.46
1:A:328:ARG:HH21	1:A:351:GLN:HE22	1.62	0.46
1:A:153:LEU:HB3	1:A:165:SER:HB3	1.98	0.46
1:B:64:VAL:HG11	1:B:91:LEU:HG	1.98	0.46
1:B:114:ASN:HB2	4:B:407:EDO:O1	2.16	0.45
1:B:55:GLU:O	1:B:56:ARG:CB	2.64	0.45
1:A:362:VAL:O	4:A:403:EDO:H12	2.16	0.45
1:B:95:PRO:HA	1:B:231:GLN:HE22	1.82	0.45
1:C:169:LEU:HD13	1:C:340:VAL:HG22	1.97	0.45
1:C:329:LEU:HD23	1:C:338:PHE:HE2	1.82	0.44
1:C:231:GLN:HA	1:C:231:GLN:OE1	2.18	0.44
1:A:363:SER:HB2	5:A:517:HOH:O	2.18	0.43
1:C:106:CYS:HB2	1:C:362:VAL:HG13	2.00	0.42
1:A:16:ARG:HA	1:A:48:SER:O	2.19	0.42
1:A:272:MET:HB3	1:A:306:MET:HE2	2.02	0.42
1:A:219:GLY:H	4:A:404:EDO:C1	2.33	0.42
1:A:222:THR:HG22	4:A:404:EDO:H22	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:88:VAL:HG21	1:C:166:VAL:HG11	2.01	0.42
1:B:158:ALA:HB2	1:B:349:ALA:HB3	2.01	0.42
1:B:112:SER:OG	1:B:189:GLU:HA	2.20	0.42
1:A:71:ARG:HG3	1:A:71:ARG:H	1.66	0.41
1:C:111:GLY:HA3	3:C:401:SO4:O4	2.20	0.41
1:C:170:ALA:HB1	1:C:174:ILE:HB	2.03	0.41
1:B:363:SER:HB2	5:B:508:HOH:O	2.21	0.40
1:B:149:PRO:O	4:B:406:EDO:H12	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:561:HOH:O	5:A:561:HOH:O[3_654]	1.92	0.28

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/360 (100%)	350 (98%)	7 (2%)	2 (1%)	30	22
1	B	332/360 (92%)	320 (96%)	8 (2%)	4 (1%)	16	8
1	C	339/360 (94%)	327 (96%)	11 (3%)	1 (0%)	46	41
All	All	1030/1080 (95%)	997 (97%)	26 (2%)	7 (1%)	26	19

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	56	ARG
1	B	123	TYR
1	B	79	SER
1	C	79	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	79	SER
1	A	57	ARG
1	B	253	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/299 (100%)	273 (91%)	27 (9%)	12	7
1	B	285/299 (95%)	260 (91%)	25 (9%)	12	7
1	C	289/299 (97%)	262 (91%)	27 (9%)	11	6
All	All	874/897 (97%)	795 (91%)	79 (9%)	12	7

All (79) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	VAL
1	A	22	ASN
1	A	31	LEU
1	A	35[A]	ARG
1	A	35[B]	ARG
1	A	56	ARG
1	A	71	ARG
1	A	76	VAL
1	A	83	LEU
1	A	147	LEU
1	A	152	LEU
1	A	153	LEU
1	A	162	ARG
1	A	169	LEU
1	A	182	GLU
1	A	183	LYS
1	A	194	LEU
1	A	198	LEU
1	A	203	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	210	LEU
1	A	217	ARG
1	A	225	SER
1	A	256	ASP
1	A	258	VAL
1	A	277	MET
1	A	294	VAL
1	A	358	TYR
1	B	22	ASN
1	B	35	ARG
1	B	41	LEU
1	B	57	ARG
1	B	61	ARG
1	B	130	THR
1	B	147	LEU
1	B	152	LEU
1	B	153	LEU
1	B	162	ARG
1	B	169	LEU
1	B	179	LEU
1	B	199	ARG
1	B	200	LEU
1	B	204	ARG
1	B	210	LEU
1	B	241	LEU
1	B	256	ASP
1	B	258	VAL
1	B	277	MET
1	B	294	VAL
1	B	298	MET
1	B	299	LEU
1	B	346	VAL
1	B	362	VAL
1	C	17	VAL
1	C	22	ASN
1	C	35	ARG
1	C	50	THR
1	C	57	ARG
1	C	76	VAL
1	C	105	LEU
1	C	116	LEU
1	C	147	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	152	LEU
1	C	153	LEU
1	C	162	ARG
1	C	186	ARG
1	C	199	ARG
1	C	203	LEU
1	C	204	ARG
1	C	215	VAL
1	C	254	ASP
1	C	256	ASP
1	C	258	VAL
1	C	273	PHE
1	C	277	MET
1	C	291	ARG
1	C	299	LEU
1	C	310	ARG
1	C	346	VAL
1	C	361	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
1	A	351	GLN
1	B	22	ASN
1	B	89	ASN
1	B	231	GLN
1	B	351	GLN
1	C	22	ASN
1	C	32	GLN
1	C	85	HIS
1	C	131	ASN
1	C	230	GLN
1	C	351	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SQS	A	401	-	19,20,20	0.47	0	16,21,21	0.98	2 (12%)
3	SO4	A	402	-	4,4,4	0.35	0	6,6,6	0.07	0
4	EDO	A	403	-	3,3,3	0.26	0	2,2,2	0.82	0
4	EDO	A	404	-	3,3,3	0.42	0	2,2,2	0.21	0
4	EDO	A	405	-	3,3,3	0.42	0	2,2,2	0.44	0
4	EDO	A	406	-	3,3,3	0.41	0	2,2,2	0.29	0
4	EDO	A	407	-	3,3,3	0.39	0	2,2,2	0.80	0
4	EDO	A	408	-	3,3,3	0.49	0	2,2,2	0.32	0
2	SQS	B	401	-	19,20,20	0.62	0	16,21,21	0.76	1 (6%)
3	SO4	B	402	-	4,4,4	0.32	0	6,6,6	0.28	0
4	EDO	B	403	-	3,3,3	0.52	0	2,2,2	0.36	0
4	EDO	B	404	-	3,3,3	0.49	0	2,2,2	0.25	0
4	EDO	B	405	-	3,3,3	0.54	0	2,2,2	0.32	0
4	EDO	B	406	-	3,3,3	0.34	0	2,2,2	0.36	0
4	EDO	B	407	-	3,3,3	0.42	0	2,2,2	0.51	0
3	SO4	C	401	-	4,4,4	0.41	0	6,6,6	0.56	0
4	EDO	C	402	-	3,3,3	0.51	0	2,2,2	0.25	0
4	EDO	C	403	-	3,3,3	0.46	0	2,2,2	0.30	0
4	EDO	C	404	-	3,3,3	0.41	0	2,2,2	0.64	0
4	EDO	C	405	-	3,3,3	0.51	0	2,2,2	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SQS	A	401	-	-	0/21/21/21	0/0/0/0
3	SO4	A	402	-	-	0/0/0/0	0/0/0/0
4	EDO	A	403	-	-	0/1/1/1	0/0/0/0
4	EDO	A	404	-	-	0/1/1/1	0/0/0/0
4	EDO	A	405	-	-	0/1/1/1	0/0/0/0
4	EDO	A	406	-	-	0/1/1/1	0/0/0/0
4	EDO	A	407	-	-	0/1/1/1	0/0/0/0
4	EDO	A	408	-	-	0/1/1/1	0/0/0/0
2	SQS	B	401	-	-	0/21/21/21	0/0/0/0
3	SO4	B	402	-	-	0/0/0/0	0/0/0/0
4	EDO	B	403	-	-	0/1/1/1	0/0/0/0
4	EDO	B	404	-	-	0/1/1/1	0/0/0/0
4	EDO	B	405	-	-	0/1/1/1	0/0/0/0
4	EDO	B	406	-	-	0/1/1/1	0/0/0/0
4	EDO	B	407	-	-	0/1/1/1	0/0/0/0
3	SO4	C	401	-	-	0/0/0/0	0/0/0/0
4	EDO	C	402	-	-	0/1/1/1	0/0/0/0
4	EDO	C	403	-	-	0/1/1/1	0/0/0/0
4	EDO	C	404	-	-	0/1/1/1	0/0/0/0
4	EDO	C	405	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	SQS	C6-C5-C4	-2.08	118.91	125.14
2	B	401	SQS	O3-C3-C2	2.14	111.06	107.48
2	A	401	SQS	O3-C3-C2	2.78	112.13	107.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	EDO	3	0
4	A	404	EDO	2	0
4	A	406	EDO	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	406	EDO	1	0
4	B	407	EDO	1	0
3	C	401	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/360 (100%)	0.04	6 (1%) 73 73	21, 31, 54, 70	0
1	B	339/360 (94%)	0.58	22 (6%) 22 23	20, 35, 71, 96	0
1	C	344/360 (95%)	0.64	25 (7%) 18 19	29, 42, 61, 85	0
All	All	1043/1080 (96%)	0.41	53 (5%) 32 33	20, 37, 63, 96	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	348	GLU	5.7
1	C	208	GLY	5.1
1	B	182	GLU	5.0
1	B	231	GLN	4.3
1	A	224	ALA	3.8
1	B	188	GLY	3.8
1	C	249	TRP	3.6
1	B	57	ARG	3.6
1	C	247	SER	3.4
1	B	158	ALA	3.3
1	B	185	ARG	3.2
1	C	318	TYR	3.2
1	C	14	PRO	3.1
1	C	315	GLU	3.0
1	C	241	LEU	2.9
1	C	240	PRO	2.9
1	C	209	ARG	2.8
1	B	306	MET	2.8
1	A	228	VAL	2.7
1	C	245	VAL	2.7
1	B	203	LEU	2.7
1	C	200	LEU	2.7
1	B	245	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	243	GLU	2.6
1	C	256	ASP	2.6
1	B	160	GLY	2.5
1	C	236	ALA	2.5
1	C	239	VAL	2.5
1	C	348	GLU	2.5
1	C	230	GLN	2.5
1	A	56	ARG	2.5
1	C	294	VAL	2.4
1	B	336	GLY	2.4
1	C	242	GLU	2.4
1	B	209	ARG	2.3
1	A	57	ARG	2.3
1	B	251	VAL	2.3
1	B	337	VAL	2.3
1	A	230	GLN	2.3
1	C	292	ALA	2.3
1	A	334	GLY	2.2
1	B	184	TYR	2.2
1	C	186	ARG	2.2
1	C	289	TYR	2.2
1	C	337	VAL	2.2
1	B	190	MET	2.1
1	C	332	LYS	2.1
1	B	242	GLU	2.1
1	B	207	ARG	2.1
1	C	205	THR	2.0
1	C	331	PRO	2.0
1	B	161	LEU	2.0
1	B	277	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EDO	A	403	4/4	0.89	0.32	10.38	36,44,47,49	0
4	EDO	C	402	4/4	0.73	0.28	7.12	48,51,53,58	0
2	SQS	B	401	21/21	0.60	0.38	6.96	56,62,71,73	0
4	EDO	B	403	4/4	0.81	0.22	6.35	40,44,46,50	0
4	EDO	A	404	4/4	0.93	0.34	4.52	45,48,50,52	0
4	EDO	A	406	4/4	0.91	0.20	4.34	35,38,39,39	0
2	SQS	A	401	21/21	0.83	0.19	3.51	47,51,76,82	0
4	EDO	A	408	4/4	0.84	0.15	3.44	50,51,51,51	0
4	EDO	B	404	4/4	0.90	0.28	3.29	59,60,61,61	0
4	EDO	A	407	4/4	0.90	0.16	2.83	38,39,40,41	0
4	EDO	B	407	4/4	0.89	0.26	2.75	55,55,55,56	0
4	EDO	B	406	4/4	0.91	0.20	2.08	42,45,46,49	0
4	EDO	B	405	4/4	0.87	0.17	1.56	51,52,52,52	0
4	EDO	C	403	4/4	0.74	0.25	1.46	51,55,55,56	0
4	EDO	A	405	4/4	0.92	0.14	0.59	43,43,45,46	0
3	SO4	C	401	5/5	0.97	0.12	-0.59	47,48,49,50	0
3	SO4	B	402	5/5	0.92	0.14	-0.69	60,61,64,71	0
3	SO4	A	402	5/5	0.94	0.12	-0.82	70,74,74,76	0
4	EDO	C	404	4/4	0.90	0.13	-1.03	42,43,43,44	0
4	EDO	C	405	4/4	0.75	0.18	-	49,50,50,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.