



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:14 PM GMT

PDB ID : 3W1U
Title : Structure of Trypanosoma cruzi dihydroorotate dehydrogenase in complex with MII-3-111
Authors : Inaoka, D.K.; Iida, M.; Tabuchi, T.; Lee, N.; Matsuoka, S.; Shiba, T.; Sakamoto, K.; Suzuki, S.; Balogun, E.O.; Nara, T.; Aoki, T.; Inoue, M.; Honma, T.; Tanaka, A.; Harada, S.; Kita, K.
Deposited on : 2012-11-21
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	<div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div>
1	B	313	<div> <div></div> <div>86%</div> <div>13%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	402	-	-	-	X
3	GOL	A	405	-	-	-	X
3	GOL	A	406	-	-	-	X
3	GOL	B	403	-	-	-	X
3	GOL	B	404	-	-	-	X
3	GOL	B	405	-	-	-	X
3	GOL	B	406	-	-	X	X
3	GOL	B	407	-	-	-	X
3	GOL	B	408	-	-	-	X
3	GOL	B	409	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 5504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydroorotate dehydrogenase (fumarate).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	0	0
			2388	1530	396	444	18			
1	B	313	Total	C	N	O	S	0	0	0
			2388	1530	396	444	18			

- Molecule 2 is 5-[2-(4-METHYLPHENYL)ETHYL]-2,6-DIOXO-1,2,3,6-TETRAHYDROPYRIMIDINE-4-CARBOXYLIC ACID (three-letter code: LRO) (formula: C₁₄H₁₄N₂O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			20	14	2	4		
2	B	1	Total	C	N	O	0	0
			20	14	2	4		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
4	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 5 is COBALT HEXAMMINE(III) (three-letter code: NCO) (formula: $\text{CoH}_{18}\text{N}_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	Co	N	0	0
			7	1	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	280	Total 280	O 280	0	0
6	B	261	Total 261	O 261	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase (fumarate)



- Molecule 1: Dihydroorotate dehydrogenase (fumarate)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.92Å 71.74Å 129.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.65 – 1.85 26.65 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (26.65-1.85) 99.9 (26.65-1.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.34 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.151 , 0.185 0.153 , 0.187	Depositor DCC
R_{free} test set	2764 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	2 of 54638 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5504	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 27.17 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.3058e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, GOL, LRO, NCO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.17	7/2440 (0.3%)	1.02	5/3305 (0.2%)
1	B	1.14	5/2440 (0.2%)	0.99	3/3305 (0.1%)
All	All	1.16	12/4880 (0.2%)	1.01	8/6610 (0.1%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	303	GLU	CD-OE2	-8.43	1.16	1.25
1	A	302	GLU	CD-OE1	-8.13	1.16	1.25
1	B	303	GLU	CD-OE2	-5.94	1.19	1.25
1	A	131	PRO	N-CD	5.80	1.55	1.47
1	A	144	GLU	CG-CD	5.75	1.60	1.51
1	B	49	PRO	N-CD	5.61	1.55	1.47
1	A	36	SER	CB-OG	-5.47	1.35	1.42
1	B	54	PRO	N-CD	5.43	1.55	1.47
1	B	105	GLU	CD-OE1	-5.43	1.19	1.25
1	A	217	PHE	CG-CD2	-5.38	1.30	1.38
1	A	205	GLU	CD-OE2	5.31	1.31	1.25
1	B	68	SER	CB-OG	5.16	1.49	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	LYS	CD-CE-NZ	8.58	131.43	111.70
1	A	88	ASP	CB-CG-OD1	5.88	123.59	118.30
1	A	133	VAL	C-N-CD	5.61	140.19	128.40
1	B	77	ASP	CB-CG-OD1	5.47	123.23	118.30
1	B	183	GLU	OE1-CD-OE2	-5.46	116.75	123.30
1	A	112	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	B	175	ASP	CB-CG-OD1	5.22	123.00	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	287	ASP	CB-CG-OD1	5.17	122.96	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2388	0	2391	16	0
1	B	2388	0	2391	30	0
2	A	20	0	13	1	0
2	B	20	0	13	3	0
3	A	30	0	40	2	0
3	B	48	0	64	12	0
4	A	31	0	19	0	0
4	B	31	0	19	1	0
5	A	7	0	0	0	0
6	A	280	0	0	6	0
6	B	261	0	0	6	0
All	All	5504	0	4950	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:132:ASN:HB2	6:B:582:HOH:O	1.64	0.97
1:B:9:HIS:CE1	3:B:406:GOL:H11	2.11	0.85
1:B:138:GLN:HG2	6:B:751:HOH:O	1.80	0.81
1:B:234:ASN:HD21	1:B:238:ARG:HE	1.30	0.80
1:A:59:MET:HE3	6:A:616:HOH:O	1.86	0.73
1:B:96:LEU:CD2	1:B:98:ILE:HD11	2.21	0.69
2:B:401:LRO:H13	3:B:403:GOL:H31	1.73	0.69
1:B:9:HIS:ND1	3:B:406:GOL:H11	2.09	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:LEU:O	1:A:9:HIS:HD2	1.78	0.66
1:A:234:ASN:HD21	1:A:238:ARG:HE	1.44	0.66
2:B:401:LRO:CAG	3:B:403:GOL:H31	2.28	0.63
1:B:93:PRO:HG3	3:B:406:GOL:H31	1.81	0.62
1:A:195:SER:HB3	6:A:763:HOH:O	2.02	0.60
1:B:82:TYR:OH	1:B:87:HIS:HD2	1.84	0.60
1:B:136:LYS:HD3	6:B:756:HOH:O	2.03	0.58
1:B:59:MET:HE3	6:B:611:HOH:O	2.04	0.58
1:A:82:TYR:OH	1:A:87:HIS:HD2	1.88	0.57
1:B:96:LEU:CD2	1:B:98:ILE:CD1	2.84	0.56
1:A:241:PRO:HA	3:A:404:GOL:H31	1.87	0.55
1:B:209:VAL:O	3:B:408:GOL:H31	2.06	0.55
1:A:111:ARG:NH1	6:A:546:HOH:O	2.39	0.55
1:A:130:CYS:SG	6:A:548:HOH:O	2.37	0.53
1:B:89:TYR:HD2	3:B:406:GOL:H12	1.73	0.52
1:B:160:PRO:HB3	1:B:188:LYS:HG3	1.93	0.51
1:A:6:LEU:HD12	1:A:6:LEU:C	2.30	0.51
1:A:120:LYS:HE3	6:A:676:HOH:O	2.12	0.50
1:B:7:LEU:O	1:B:9:HIS:HD2	1.96	0.49
2:B:401:LRO:H12	6:B:753:HOH:O	2.14	0.47
1:B:234:ASN:ND2	1:B:238:ARG:HE	2.05	0.47
1:B:96:LEU:HD21	1:B:98:ILE:HD11	1.95	0.47
1:A:87:HIS:HE1	1:A:92:LYS:O	1.99	0.45
1:B:237:TYR:CE2	3:B:404:GOL:H31	2.51	0.45
1:B:87:HIS:HE1	1:B:92:LYS:O	2.00	0.45
1:B:20:GLY:HA3	4:B:410:FMN:N5	2.32	0.44
1:B:237:TYR:CZ	3:B:404:GOL:H2	2.53	0.44
1:B:71:LEU:N	1:B:72:PRO:CD	2.81	0.43
1:A:59:MET:CE	6:A:616:HOH:O	2.57	0.43
1:B:162:GLY:HA3	1:B:189:PHE:CE1	2.53	0.43
1:B:241:PRO:HA	3:B:404:GOL:H12	2.00	0.43
1:B:96:LEU:HD22	1:B:98:ILE:HD11	1.99	0.43
1:A:97:SER:HA	1:A:125:GLU:O	2.19	0.42
1:B:5:ASN:ND2	6:B:706:HOH:O	2.51	0.42
1:B:226:LEU:HB3	1:B:227:PRO:HD3	2.01	0.41
1:A:217:PHE:CE1	3:B:402:GOL:H12	2.55	0.41
1:A:234:ASN:ND2	1:A:238:ARG:HE	2.14	0.41
1:B:162:GLY:HA3	1:B:189:PHE:CZ	2.54	0.41
1:B:97:SER:HA	1:B:125:GLU:O	2.20	0.41
2:A:401:LRO:H9	3:A:405:GOL:H12	2.01	0.41
1:B:9:HIS:CE1	3:B:406:GOL:C1	2.94	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:ILE:HA	1:A:219:GLY:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/313 (99%)	301 (97%)	10 (3%)	0	100	100
1	B	311/313 (99%)	303 (97%)	8 (3%)	0	100	100
All	All	622/626 (99%)	604 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/258 (100%)	256 (99%)	2 (1%)	86	82
1	B	258/258 (100%)	251 (97%)	7 (3%)	52	34
All	All	516/516 (100%)	507 (98%)	9 (2%)	68	54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	131	PRO
1	B	0	MET
1	B	6	LEU
1	B	81	LYS
1	B	130	CYS
1	B	136	LYS
1	B	138	GLN
1	B	312	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	9	HIS
1	A	87	HIS
1	A	234	ASN
1	A	275	GLN
1	B	9	HIS
1	B	87	HIS
1	B	234	ASN
1	B	275	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	LRO	A	401	-	16,21,21	3.12	8 (50%)	16,29,29	3.14	7 (43%)
3	GOL	A	402	-	5,5,5	0.73	0	5,5,5	0.99	0
3	GOL	A	403	-	5,5,5	0.33	0	5,5,5	0.98	0
3	GOL	A	404	-	5,5,5	0.35	0	5,5,5	0.65	0
3	GOL	A	405	-	5,5,5	0.98	0	5,5,5	1.20	0
3	GOL	A	406	-	5,5,5	0.69	0	5,5,5	0.47	0
4	FMN	A	407	-	32,33,33	1.48	7 (21%)	34,50,50	1.94	4 (11%)
5	NCO	A	408	-	6,6,6	1.85	3 (50%)	0,15,15	0.00	-
2	LRO	B	401	-	16,21,21	2.34	6 (37%)	16,29,29	2.42	3 (18%)
3	GOL	B	402	-	5,5,5	0.69	0	5,5,5	0.38	0
3	GOL	B	403	-	5,5,5	0.71	0	5,5,5	0.81	0
3	GOL	B	404	-	5,5,5	0.73	0	5,5,5	0.74	0
3	GOL	B	405	-	5,5,5	1.03	0	5,5,5	0.74	0
3	GOL	B	406	-	5,5,5	0.42	0	5,5,5	2.03	2 (40%)
3	GOL	B	407	-	5,5,5	0.61	0	5,5,5	0.41	0
3	GOL	B	408	-	5,5,5	0.42	0	5,5,5	0.75	0
3	GOL	B	409	-	5,5,5	1.05	1 (20%)	5,5,5	0.99	0
4	FMN	B	410	-	32,33,33	1.47	7 (21%)	34,50,50	2.47	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LRO	A	401	-	-	0/5/9/9	0/2/2/2
3	GOL	A	402	-	-	0/4/4/4	0/0/0/0
3	GOL	A	403	-	-	0/4/4/4	0/0/0/0
3	GOL	A	404	-	-	0/4/4/4	0/0/0/0
3	GOL	A	405	-	-	0/4/4/4	0/0/0/0
3	GOL	A	406	-	-	0/4/4/4	0/0/0/0
4	FMN	A	407	-	-	0/18/18/18	0/3/3/3
5	NCO	A	408	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LRO	B	401	-	-	0/5/9/9	0/2/2/2
3	GOL	B	402	-	-	0/4/4/4	0/0/0/0
3	GOL	B	403	-	-	0/4/4/4	0/0/0/0
3	GOL	B	404	-	-	0/4/4/4	0/0/0/0
3	GOL	B	405	-	-	0/4/4/4	0/0/0/0
3	GOL	B	406	-	-	0/4/4/4	0/0/0/0
3	GOL	B	407	-	-	0/4/4/4	0/0/0/0
3	GOL	B	408	-	-	0/4/4/4	0/0/0/0
3	GOL	B	409	-	-	0/4/4/4	0/0/0/0
4	FMN	B	410	-	-	0/18/18/18	0/3/3/3

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	LRO	CAK-CAR	-8.48	1.40	1.52
2	B	401	LRO	CAJ-CAP	-3.79	1.40	1.51
2	B	401	LRO	CAK-CAR	-3.69	1.47	1.52
4	B	410	FMN	C6-C5A	-2.70	1.37	1.41
4	A	407	FMN	C1'-N10	-2.65	1.45	1.48
2	A	401	LRO	CAJ-CAP	-2.65	1.44	1.51
4	B	410	FMN	C1'-N10	-2.14	1.46	1.48
2	B	401	LRO	CAA-CAO	-2.01	1.43	1.51
4	A	407	FMN	C4-C4A	2.02	1.45	1.41
5	A	408	NCO	CO-N4	2.04	2.04	1.96
2	A	401	LRO	CAG-CAO	2.09	1.44	1.38
4	B	410	FMN	C4-C4A	2.11	1.45	1.41
2	B	401	LRO	CAI-CAG	2.12	1.43	1.38
4	A	407	FMN	C9A-N10	2.18	1.41	1.38
3	B	409	GOL	O2-C2	2.18	1.49	1.43
4	B	410	FMN	O3'-C3'	2.31	1.48	1.43
5	A	408	NCO	CO-N2	2.35	2.05	1.96
2	A	401	LRO	CAH-CAF	2.43	1.43	1.38
4	B	410	FMN	C9A-C5A	2.49	1.47	1.42
4	B	410	FMN	C4A-C10	2.55	1.45	1.40
4	A	407	FMN	C5A-N5	2.62	1.39	1.35
4	A	407	FMN	C9A-C5A	2.70	1.48	1.42
4	A	407	FMN	C5'-C4'	2.70	1.55	1.51
5	A	408	NCO	CO-N6	2.83	2.07	1.96
2	A	401	LRO	CAI-CAG	2.92	1.44	1.38
4	B	410	FMN	C5A-N5	3.04	1.40	1.35
2	A	401	LRO	CAS-NAL	3.22	1.44	1.38
2	A	401	LRO	CAQ-NAL	3.50	1.40	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	407	FMN	C4A-C10	3.64	1.47	1.40
2	B	401	LRO	CAT-NAM	4.18	1.40	1.33
2	B	401	LRO	CAQ-NAL	4.75	1.42	1.34
2	A	401	LRO	CAT-NAM	4.98	1.42	1.33

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	LRO	NAM-CAS-NAL	-5.32	118.73	127.69
4	B	410	FMN	C4A-C4-N3	-5.02	116.96	123.52
2	B	401	LRO	NAM-CAS-NAL	-4.99	119.29	127.69
4	A	407	FMN	C4A-C4-N3	-4.69	117.39	123.52
2	A	401	LRO	CAH-CAF-CAO	-3.25	116.84	121.36
3	B	406	GOL	C3-C2-C1	-2.96	98.59	111.06
4	A	407	FMN	N3-C2-N1	-2.70	123.15	127.69
2	A	401	LRO	CAG-CAI-CAP	-2.65	117.30	121.02
4	B	410	FMN	C4-C4A-C10	-2.46	118.36	119.94
4	B	410	FMN	C4A-C10-N10	-2.45	118.74	120.52
4	B	410	FMN	C9A-C5A-N5	-2.26	118.50	122.18
4	B	410	FMN	N3-C2-N1	-2.22	123.95	127.69
4	B	410	FMN	C6-C5A-N5	2.02	121.43	118.92
2	A	401	LRO	CAI-CAG-CAO	2.48	124.80	121.36
3	B	406	GOL	O3-C3-C2	2.52	122.73	109.97
2	A	401	LRO	CAS-NAL-CAQ	2.64	118.33	115.68
4	A	407	FMN	O3P-P-O1P	3.13	120.84	110.63
4	B	410	FMN	C4A-N5-C5A	3.75	121.14	116.72
2	A	401	LRO	CAF-CAH-CAP	4.01	126.67	121.02
2	B	401	LRO	CAS-NAL-CAQ	4.81	120.50	115.68
4	B	410	FMN	C1'-N10-C9A	5.95	125.73	118.83
2	B	401	LRO	CAT-NAM-CAS	6.12	120.26	115.16
4	A	407	FMN	C4-N3-C2	7.83	121.69	115.16
2	A	401	LRO	CAT-NAM-CAS	8.40	122.16	115.16
4	B	410	FMN	C4-N3-C2	9.12	122.76	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

10 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	LRO	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	GOL	1	0
3	A	405	GOL	1	0
2	B	401	LRO	3	0
3	B	402	GOL	1	0
3	B	403	GOL	2	0
3	B	404	GOL	3	0
3	B	406	GOL	5	0
3	B	408	GOL	1	0
4	B	410	FMN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/313 (100%)	-0.38	4 (1%) 79 79	8, 13, 25, 46	0
1	B	313/313 (100%)	-0.32	4 (1%) 79 79	8, 14, 32, 49	0
All	All	626/626 (100%)	-0.35	8 (1%) 79 79	8, 14, 29, 49	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	130	CYS	4.1
1	B	131	PRO	3.4
1	A	312	GLU	3.2
1	B	312	GLU	3.1
1	B	133	VAL	2.6
1	B	132	ASN	2.3
1	A	119	GLU	2.2
1	A	131	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	B	408	6/6	0.79	0.34	20.40	40,47,52,52	0
3	GOL	B	406	6/6	0.82	0.41	14.83	35,42,45,51	0
3	GOL	B	404	6/6	0.82	0.26	13.33	37,37,39,39	0
3	GOL	B	407	6/6	0.86	0.18	8.37	30,35,36,38	0
3	GOL	A	402	6/6	0.80	0.16	8.06	20,24,25,26	0
3	GOL	A	405	6/6	0.80	0.28	7.20	24,32,43,44	0
3	GOL	B	409	6/6	0.82	0.19	6.30	28,35,40,43	0
3	GOL	B	405	6/6	0.59	0.24	5.40	43,47,48,51	0
3	GOL	A	406	6/6	0.86	0.26	4.59	27,35,38,45	0
3	GOL	B	403	6/6	0.86	0.17	3.76	22,35,38,41	0
3	GOL	B	402	6/6	0.96	0.08	0.71	20,22,22,27	0
2	LRO	A	401	20/20	0.94	0.11	0.42	11,14,34,34	0
5	NCO	A	408	7/7	0.99	0.07	0.11	12,12,13,13	0
4	FMN	A	407	31/31	0.98	0.09	-0.02	7,9,13,15	0
4	FMN	B	410	31/31	0.98	0.09	-0.16	8,9,10,12	0
2	LRO	B	401	20/20	0.95	0.10	-0.21	10,13,32,37	0
3	GOL	A	403	6/6	0.97	0.05	-4.32	19,20,21,22	0
3	GOL	A	404	6/6	0.87	0.20	-	43,47,49,50	0

6.5 Other polymers

There are no such residues in this entry.