



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:10 AM GMT

PDB ID : 2W62  
Title : Saccharomyces cerevisiae Gas2p in complex with laminaripentaose  
Authors : Schuettelkopf, A.W.; Hurtado-Guerrero, R.; Van Aalten, D.M.F.  
Deposited on : 2008-12-16  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BGC	A	1495	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BGC	A	1498	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GLYCOLIPID-ANCHORED SURFACE PROTEIN 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	6	0
			3499	2234	558	684	23			

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	5	Total	C	O	0	0
			56	30	26		
2	A	5	Total	C	O	0	0
			56	30	26		

- Molecule 3 is 1,4-BUTANEDIOL (three-letter code: BU1) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	313	Total 313	O 313	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.04Å 70.84Å 149.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.00 – 1.85 19.95 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (23.00-1.85) 99.5 (19.95-1.85)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.11 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.4.0069	Depositor
R, $R_{free}$	0.183 , 0.210 0.187 , 0.211	Depositor DCC
$R_{free}$ test set	555 reflections (1.23%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 45861 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3930	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, BU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/3601	0.62	0/4869

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3338	29	0
2	A	112	0	96	1	0
3	A	6	0	10	0	0
4	A	313	0	0	2	0
All	All	3930	0	3444	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:MET:HE2	1:A:310:PHE:H	1.37	0.88
1:A:29:SER:OG	1:A:31[B]:GLU:HG2	1.78	0.82
1:A:220:MET:HG2	4:A:2168:HOH:O	1.91	0.71
1:A:42:LYS:HE3	1:A:348:ILE:HD11	1.72	0.70
1:A:392:CYS:O	1:A:396:ILE:HG12	1.93	0.69
1:A:433:THR:OG1	1:A:435:GLU:HG3	1.94	0.67
1:A:235:LYS:HE3	4:A:2176:HOH:O	2.01	0.60
1:A:191:LYS:HD2	1:A:228:TYR:CE2	2.39	0.57
1:A:308:MET:HE1	1:A:310:PHE:HD2	1.70	0.56
1:A:149:VAL:O	1:A:153:GLU:HG2	2.07	0.55
1:A:395:GLU:O	1:A:478:GLU:HG2	2.12	0.49
1:A:308:MET:CE	1:A:310:PHE:H	2.18	0.48
1:A:282:ARG:HA	1:A:283:PRO:C	2.34	0.48
1:A:139[B]:SER:OG	2:A:1492:BGC:H2	2.14	0.47
1:A:29:SER:HG	1:A:31[B]:GLU:HG2	1.74	0.47
1:A:60:GLN:H	1:A:308:MET:CE	2.28	0.47
1:A:400:GLU:HB2	1:A:474:TYR:CE1	2.51	0.46
1:A:60:GLN:H	1:A:308:MET:HE1	1.81	0.46
1:A:478:GLU:HA	1:A:481:GLN:HG2	1.97	0.45
1:A:415:PHE:CZ	1:A:448:LEU:HG	2.51	0.45
1:A:152:PHE:CE2	1:A:156:LYS:HE2	2.52	0.44
1:A:246:TRP:CG	1:A:288:GLU:HB3	2.52	0.44
1:A:392:CYS:HA	1:A:395:GLU:HG3	2.00	0.43
1:A:74:ALA:C	1:A:75:PHE:HD1	2.22	0.42
1:A:308:MET:HE1	1:A:310:PHE:CD2	2.53	0.41
1:A:191:LYS:CD	1:A:228:TYR:CE2	3.05	0.40
1:A:308:MET:HE2	1:A:310:PHE:N	2.20	0.40
1:A:31[B]:GLU:HG2	1:A:31[B]:GLU:H	1.70	0.40
1:A:389:LYS:O	1:A:392:CYS:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	433/555 (78%)	429 (99%)	3 (1%)	1 (0%)	52 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	177	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	388/485 (80%)	380 (98%)	8 (2%)	61 45

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	LEU
1	A	135	GLU
1	A	168	LEU
1	A	312	GLU
1	A	363	GLU
1	A	395	GLU
1	A	471	LYS
1	A	489	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BGC	A	1490	2	12,12,12	1.07	1 (8%)	17,17,17	0.92	2 (11%)
2	BGC	A	1491	2	11,11,12	0.96	0	14,15,17	1.10	2 (14%)
2	BGC	A	1492	2	11,11,12	0.82	0	14,15,17	0.99	1 (7%)
2	BGC	A	1493	2	11,11,12	0.96	0	14,15,17	1.78	1 (7%)
2	BGC	A	1494	2	11,11,12	0.45	0	14,15,17	1.16	2 (14%)
2	BGC	A	1495	2	12,12,12	0.89	0	17,17,17	0.60	0
2	BGC	A	1496	2	11,11,12	0.67	0	14,15,17	1.24	1 (7%)
2	BGC	A	1497	2	11,11,12	0.97	0	14,15,17	1.09	0
2	BGC	A	1498	2	11,11,12	0.85	1 (9%)	14,15,17	0.70	0
2	BGC	A	1499	2	11,11,12	1.42	1 (9%)	14,15,17	1.83	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	1490	2	-	0/2/22/22	0/1/1/1
2	BGC	A	1491	2	-	0/2/19/22	0/1/1/1
2	BGC	A	1492	2	-	0/2/19/22	0/1/1/1
2	BGC	A	1493	2	-	0/2/19/22	0/1/1/1
2	BGC	A	1494	2	-	0/2/19/22	0/1/1/1
2	BGC	A	1495	2	-	0/2/22/22	0/1/1/1
2	BGC	A	1496	2	-	0/2/19/22	0/1/1/1
2	BGC	A	1497	2	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	1498	2	-	0/2/19/22	0/1/1/1
2	BGC	A	1499	2	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1498	BGC	C2-C3	2.19	1.55	1.52
2	A	1490	BGC	O5-C1	2.46	1.47	1.43
2	A	1499	BGC	C2-C3	3.77	1.57	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1493	BGC	O3-C3-C2	-6.24	98.73	110.00
2	A	1496	BGC	O3-C3-C2	-3.39	103.86	110.00
2	A	1499	BGC	O2-C2-C3	-3.02	104.04	110.12
2	A	1491	BGC	C1-C2-C3	-2.65	106.40	109.54
2	A	1491	BGC	O3-C3-C4	-2.30	105.15	110.34
2	A	1492	BGC	O5-C1-C2	-2.24	107.22	110.86
2	A	1490	BGC	O3-C3-C4	-2.15	105.50	110.34
2	A	1499	BGC	O5-C1-C2	-2.04	107.55	110.86
2	A	1490	BGC	O5-C5-C4	-2.03	105.86	109.68
2	A	1494	BGC	C3-C4-C5	2.17	113.97	110.20
2	A	1494	BGC	C1-O5-C5	2.21	115.05	112.25
2	A	1499	BGC	C1-O5-C5	3.22	116.33	112.25
2	A	1499	BGC	O2-C2-C1	3.24	115.69	109.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1492	BGC	1	0

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	BU1	A	1500	-	5,5,5	0.64	0	4,4,4	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BU1	A	1500	-	-	0/3/3/3	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	436/555 (78%)	1.28	90 (20%) <b>1</b> <b>1</b>	31, 40, 55, 73	5 (1%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	414	TYR	8.6
1	A	364	SER	8.0
1	A	75	PHE	7.4
1	A	482	PRO	7.1
1	A	64	SER	7.1
1	A	489	CYS	7.1
1	A	404	PHE	7.0
1	A	73	GLY	6.8
1	A	363	GLU	6.1
1	A	355	THR	5.5
1	A	295	ASN	5.4
1	A	409	GLY	5.1
1	A	459	ILE	4.8
1	A	411	TYR	4.7
1	A	468	LEU	4.6
1	A	465	HIS	4.6
1	A	177	VAL	4.3
1	A	403	PRO	4.3
1	A	292	LEU	4.2
1	A	318	VAL	4.1
1	A	455	LEU	4.0
1	A	74	ALA	3.9
1	A	461	ALA	3.8
1	A	458	LYS	3.7
1	A	63	ARG	3.7
1	A	412	GLU	3.7
1	A	324	ASN	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	481	GLN	3.6
1	A	313	GLU	3.6
1	A	187	SER	3.6
1	A	387	ARG	3.5
1	A	168	LEU	3.5
1	A	377	ALA	3.5
1	A	396	ILE	3.4
1	A	410	LYS	3.4
1	A	462	ASN	3.3
1	A	169	LEU	3.3
1	A	76	GLU	3.2
1	A	225	LEU	3.2
1	A	314	ASN	3.2
1	A	323	ASP	3.2
1	A	190	VAL	3.1
1	A	109	ILE	3.0
1	A	401	ILE	2.9
1	A	352	GLU	2.9
1	A	471	LYS	2.8
1	A	36	ILE	2.7
1	A	201	ILE	2.7
1	A	129	VAL	2.7
1	A	271	VAL	2.7
1	A	119	MET	2.7
1	A	413	GLU	2.7
1	A	366	GLU	2.6
1	A	178	THR	2.6
1	A	338	LYS	2.6
1	A	397	LEU	2.6
1	A	336	LEU	2.6
1	A	460	GLY	2.6
1	A	473	VAL	2.6
1	A	306	ALA	2.5
1	A	469	ASN	2.5
1	A	415	PHE	2.4
1	A	106	VAL	2.4
1	A	354	LEU	2.4
1	A	186	ALA	2.4
1	A	228	TYR	2.4
1	A	259	GLU	2.4
1	A	189	PHE	2.4
1	A	452	LEU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	276	PHE	2.4
1	A	365	VAL	2.3
1	A	188	PRO	2.3
1	A	393	LEU	2.3
1	A	77	THR	2.3
1	A	395	GLU	2.3
1	A	301	TRP	2.3
1	A	265	GLU	2.3
1	A	249	TYR	2.3
1	A	29	SER	2.2
1	A	159	ILE	2.2
1	A	417	TYR	2.2
1	A	467	PRO	2.2
1	A	349	THR	2.2
1	A	62	GLN	2.2
1	A	375	TRP	2.2
1	A	257	TYR	2.2
1	A	184	THR	2.2
1	A	273	PHE	2.1
1	A	209	ILE	2.1
1	A	108	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	A	1495	12/12	0.87	0.33	7.05	41,46,49,50	0
2	BGC	A	1498	11/12	0.94	0.21	2.46	38,43,44,45	0
2	BGC	A	1497	11/12	0.94	0.18	1.72	39,41,42,42	0
2	BGC	A	1491	11/12	0.91	0.14	-0.43	39,41,43,44	0
2	BGC	A	1492	11/12	0.92	0.09	-1.50	36,38,39,39	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BGC	A	1490	12/12	0.93	0.10	-1.70	41,42,47,47	0
2	BGC	A	1493	11/12	0.92	0.16	-	39,42,43,46	0
2	BGC	A	1499	11/12	0.80	0.35	-	43,48,52,56	0
2	BGC	A	1496	11/12	0.91	0.29	-	40,40,41,41	0
2	BGC	A	1494	11/12	0.78	0.45	-	54,59,62,65	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BU1	A	1500	6/6	0.78	0.20	1.54	48,52,53,58	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.