



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:49 PM GMT

PDB ID : 4WBO  
Title : Bovine G Protein Coupled Receptor Kinase 1 in Complex with Amlexanox  
Authors : Homan, K.T.; Tesmer, J.J.G.  
Deposited on : 2014-09-03  
Resolution : 2.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2676 (2.84-2.80)
Clashscore	102246	3124 (2.84-2.80)
Ramachandran outliers	100387	3072 (2.84-2.80)
Sidechain outliers	100360	3074 (2.84-2.80)
RSRZ outliers	91569	2690 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	541	<div> <div>6%</div> <div>87%</div> <div>8%</div> </div>
1	B	541	<div> <div>5%</div> <div>88%</div> <div>9%</div> </div>
1	C	541	<div> <div>8%</div> <div>86%</div> <div>11%</div> </div>
1	D	541	<div> <div>9%</div> <div>88%</div> <div>9%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhodopsin kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	497	Total	C	N	O	S	0	0	0
			3988	2559	689	721	19			
1	B	495	Total	C	N	O	S	0	0	0
			3976	2550	687	720	19			
1	C	481	Total	C	N	O	S	0	0	0
			3865	2481	664	701	19			
1	D	491	Total	C	N	O	S	0	0	0
			3957	2541	681	716	19			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	536	HIS	-	expression tag	UNP P28327
A	537	HIS	-	expression tag	UNP P28327
A	538	HIS	-	expression tag	UNP P28327
A	539	HIS	-	expression tag	UNP P28327
A	540	HIS	-	expression tag	UNP P28327
A	541	HIS	-	expression tag	UNP P28327
B	536	HIS	-	expression tag	UNP P28327
B	537	HIS	-	expression tag	UNP P28327
B	538	HIS	-	expression tag	UNP P28327
B	539	HIS	-	expression tag	UNP P28327
B	540	HIS	-	expression tag	UNP P28327
B	541	HIS	-	expression tag	UNP P28327
C	536	HIS	-	expression tag	UNP P28327
C	537	HIS	-	expression tag	UNP P28327
C	538	HIS	-	expression tag	UNP P28327
C	539	HIS	-	expression tag	UNP P28327
C	540	HIS	-	expression tag	UNP P28327
C	541	HIS	-	expression tag	UNP P28327
D	536	HIS	-	expression tag	UNP P28327
D	537	HIS	-	expression tag	UNP P28327
D	538	HIS	-	expression tag	UNP P28327

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	539	HIS	-	expression tag	UNP P28327
D	540	HIS	-	expression tag	UNP P28327
D	541	HIS	-	expression tag	UNP P28327

- Molecule 2 is 2-amino-7-(1-methylethyl)-5-oxo-5H-chromeno[2,3-b]pyridine-3-carboxylic acid (three-letter code: ANW) (formula: C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			22	16	2	4		
2	B	1	Total	C	N	O	0	0
			22	16	2	4		
2	C	1	Total	C	N	O	0	0
			22	16	2	4		
2	D	1	Total	C	N	O	0	0
			22	16	2	4		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	19	Total	O	0	0
			19	19		
4	B	14	Total	O	0	0
			14	14		
4	C	12	Total	O	0	0
			12	12		
4	D	9	Total	O	0	0
			9	9		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Rhodopsin kinase




#### • Molecule 1: Rhodopsin kinase





#### • Molecule 1: Rhodopsin kinase





● Molecule 1: Rhodopsin kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.11Å 119.17Å 174.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.89 – 2.81 24.89 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.4 (24.89-2.81) 99.6 (24.89-2.81)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.241 , 0.267 0.241 , 0.267	Depositor DCC
$R_{free}$ test set	3023 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	75.1	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 36.8	EDS
Estimated twinning fraction	0.013 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 60016 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	15932	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ANW, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/4083	0.51	1/5507 (0.0%)
1	B	0.31	0/4070	0.51	0/5490
1	C	0.31	0/3957	0.52	3/5337 (0.1%)
1	D	0.30	0/4049	0.49	0/5460
All	All	0.31	0/16159	0.51	4/21794 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	495	PHE	CB-CG-CD2	-6.07	116.55	120.80
1	C	495	PHE	CB-CG-CD1	5.98	124.99	120.80
1	C	31	ARG	NE-CZ-NH1	5.11	122.86	120.30
1	A	458	ARG	N-CA-CB	-5.07	101.48	110.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	393	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3988	0	3961	10	0
1	B	3976	0	3943	6	0
1	C	3865	0	3825	8	0
1	D	3957	0	3930	6	0
2	A	22	0	13	1	0
2	B	22	0	13	0	0
2	C	22	0	13	2	0
2	D	22	0	13	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	19	0	0	0	0
4	B	14	0	0	0	0
4	C	12	0	0	0	0
4	D	9	0	0	0	0
All	All	15932	0	15711	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:478:ALA:HB2	2:D:601:ANW:HACA	1.75	0.67
1:A:191:ARG:HA	1:C:102:LEU:HD11	1.84	0.59
1:A:233:MET:HE2	1:A:237:LYS:HG3	1.86	0.58
1:D:198:PHE:O	1:D:199:GLY:O	2.23	0.57
1:A:481:ILE:HG23	1:A:481:ILE:O	2.05	0.56
1:B:344:THR:HG23	1:B:345:LYS:HG3	1.88	0.56
1:A:77:ARG:HG3	1:A:86:LEU:HD21	1.88	0.53
1:B:224:LYS:HA	1:B:229:TYR:HE1	1.74	0.53
1:A:233:MET:CE	1:A:237:LYS:HG3	2.39	0.52
1:D:242:VAL:HG13	1:D:304:HIS:CD2	2.45	0.52
1:C:242:VAL:HG13	1:C:304:HIS:CD2	2.45	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:VAL:HG13	1:B:304:HIS:CD2	2.45	0.51
1:B:116:GLU:OE1	1:B:118:GLN:N	2.43	0.50
1:C:495:PHE:N	1:C:495:PHE:CD1	2.80	0.50
1:D:478:ALA:CB	2:D:601:ANW:HAEB	2.42	0.49
1:A:191:ARG:HA	1:C:102:LEU:CD1	2.44	0.48
1:C:193:LEU:HD11	1:C:203:ALA:HB2	1.96	0.48
1:A:513:ILE:HB	1:A:514:PRO:HD3	1.97	0.46
1:D:513:ILE:HB	1:D:514:PRO:HD3	1.99	0.44
2:C:601:ANW:NAB	2:C:601:ANW:OAU	2.51	0.43
1:D:478:ALA:HB2	2:D:601:ANW:HAEB	2.00	0.43
1:C:513:ILE:HB	1:C:514:PRO:HD3	2.00	0.43
2:A:601:ANW:NAB	2:A:601:ANW:OAU	2.51	0.43
1:B:224:LYS:NZ	1:B:494:ALA:O	2.52	0.42
1:C:268:ASN:HA	2:C:601:ANW:HAJ	2.00	0.42
1:B:513:ILE:HB	1:B:514:PRO:HD3	2.02	0.42
1:A:225:LYS:HB2	1:A:225:LYS:HE3	1.89	0.41
1:A:233:MET:HE3	1:A:236:LYS:HE3	2.02	0.41
1:C:223:LEU:O	1:C:225:LYS:O	2.38	0.40
1:A:491:LYS:HD3	1:A:491:LYS:N	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	493/541 (91%)	475 (96%)	16 (3%)	2 (0%)	39	73
1	B	491/541 (91%)	476 (97%)	15 (3%)	0	100	100
1	C	477/541 (88%)	461 (97%)	16 (3%)	0	100	100
1	D	485/541 (90%)	471 (97%)	13 (3%)	1 (0%)	52	83
All	All	1946/2164 (90%)	1883 (97%)	60 (3%)	3 (0%)	52	83

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	199	GLY
1	A	280	ASP
1	A	30	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/447 (93%)	397 (96%)	17 (4%)	37	71
1	B	413/447 (92%)	404 (98%)	9 (2%)	60	88
1	C	401/447 (90%)	393 (98%)	8 (2%)	63	89
1	D	413/447 (92%)	405 (98%)	8 (2%)	65	90
All	All	1641/1788 (92%)	1599 (97%)	42 (3%)	54	85

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	ASP
1	A	105	GLN
1	A	191	ARG
1	A	206	MET
1	A	222	ARG
1	A	225	LYS
1	A	236	LYS
1	A	279	VAL
1	A	282	ASP
1	A	283	ASN
1	A	390	ARG
1	A	396	VAL
1	A	464	MET
1	A	480	ASN
1	A	491	LYS
1	A	493	VAL
1	A	497	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	99	ASP
1	B	105	GLN
1	B	206	MET
1	B	236	LYS
1	B	390	ARG
1	B	395	LYS
1	B	396	VAL
1	B	482	GLN
1	B	493	VAL
1	C	99	ASP
1	C	193	LEU
1	C	206	MET
1	C	236	LYS
1	C	390	ARG
1	C	396	VAL
1	C	399	LYS
1	C	495	PHE
1	D	99	ASP
1	D	206	MET
1	D	236	LYS
1	D	390	ARG
1	D	396	VAL
1	D	474	ARG
1	D	480	ASN
1	D	504	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	480	ASN
1	B	304	HIS
1	C	304	HIS
1	D	480	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	ANW	A	601	-	20,24,24	1.70	2 (10%)	23,36,36	1.57	5 (21%)
2	ANW	B	601	-	20,24,24	1.76	3 (15%)	23,36,36	1.44	5 (21%)
2	ANW	C	601	-	20,24,24	1.80	3 (15%)	23,36,36	1.77	5 (21%)
2	ANW	D	601	-	20,24,24	1.76	3 (15%)	23,36,36	1.58	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANW	A	601	-	-	0/4/8/8	0/3/3/3
2	ANW	B	601	-	-	0/4/8/8	0/3/3/3
2	ANW	C	601	-	-	0/4/8/8	0/3/3/3
2	ANW	D	601	-	-	0/4/8/8	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	601	ANW	CAN-CAM	-2.63	1.39	1.43
2	D	601	ANW	CAN-CAM	-2.42	1.39	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	ANW	CAN-CAM	-2.37	1.39	1.43
2	B	601	ANW	CAN-CAM	-2.25	1.39	1.43
2	D	601	ANW	CAM-NAA	2.06	1.35	1.32
2	B	601	ANW	CAG-CAF	2.19	1.40	1.37
2	C	601	ANW	CAM-NAA	2.31	1.36	1.32
2	D	601	ANW	CAM-NAB	5.84	1.48	1.34
2	A	601	ANW	CAM-NAB	5.92	1.48	1.34
2	C	601	ANW	CAM-NAB	6.09	1.49	1.34
2	B	601	ANW	CAM-NAB	6.13	1.49	1.34

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	601	ANW	CAJ-CAI-CAH	-3.24	119.59	123.11
2	A	601	ANW	CAG-CAF-CAD	-3.16	117.18	121.67
2	D	601	ANW	CAJ-CAI-CAH	-2.90	119.96	123.11
2	C	601	ANW	CAG-CAF-CAD	-2.89	117.57	121.67
2	A	601	ANW	CAJ-CAI-CAH	-2.36	120.54	123.11
2	B	601	ANW	CAJ-CAI-CAH	-2.29	120.62	123.11
2	B	601	ANW	CAF-CAG-CAH	-2.13	120.09	121.82
2	B	601	ANW	OAS-CAI-CAJ	2.15	118.77	116.10
2	B	601	ANW	CAG-CAH-CAI	2.31	119.19	116.42
2	A	601	ANW	CAG-CAH-CAI	2.53	119.44	116.42
2	A	601	ANW	OAS-CAI-CAJ	2.68	119.43	116.10
2	D	601	ANW	CAG-CAH-CAI	2.80	119.76	116.42
2	C	601	ANW	CAG-CAH-CAI	2.95	119.95	116.42
2	D	601	ANW	OAS-CAI-CAJ	3.16	120.02	116.10
2	D	601	ANW	CAM-NAA-CAL	3.23	120.69	117.56
2	C	601	ANW	OAS-CAI-CAJ	3.26	120.15	116.10
2	A	601	ANW	CAM-NAA-CAL	3.51	120.95	117.56
2	B	601	ANW	CAM-NAA-CAL	4.02	121.45	117.56
2	C	601	ANW	CAM-NAA-CAL	4.02	121.45	117.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	ANW	1	0
2	C	601	ANW	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	601	ANW	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	497/541 (91%)	0.21	33 (6%) 22 12	46, 80, 140, 174	0
1	B	495/541 (91%)	0.26	27 (5%) 29 18	46, 84, 161, 232	0
1	C	481/541 (88%)	0.44	45 (9%) 11 5	54, 89, 156, 232	0
1	D	491/541 (90%)	0.47	49 (9%) 9 4	43, 97, 157, 190	0
All	All	1964/2164 (90%)	0.34	154 (7%) 16 8	43, 88, 154, 232	0

All (154) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	197	GLY	7.9
1	D	478	ALA	7.5
1	D	279	VAL	5.8
1	C	471	PRO	5.5
1	C	496	GLU	5.3
1	C	499	ASP	5.3
1	D	477	TYR	5.2
1	C	472	ASP	5.1
1	A	282	ASP	5.0
1	C	191	ARG	4.8
1	C	504	GLN	4.7
1	D	281	GLU	4.6
1	D	473	SER	4.5
1	B	498	ALA	4.5
1	A	475	THR	4.4
1	D	474	ARG	4.4
1	A	473	SER	4.3
1	D	480	ASN	4.2
1	D	440	ASP	4.2
1	D	283	ASN	4.1
1	D	372	TYR	4.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	C	527	ASP	4.0
1	A	476	VAL	3.9
1	D	414	ASP	3.9
1	B	474	ARG	3.9
1	D	479	LYS	3.8
1	D	376	GLY	3.8
1	D	282	ASP	3.8
1	C	497	LYS	3.8
1	A	140	ASP	3.8
1	B	33	ARG	3.7
1	C	140	ASP	3.7
1	B	480	ASN	3.6
1	D	392	ARG	3.6
1	A	198	PHE	3.6
1	B	230	GLN	3.6
1	D	476	VAL	3.5
1	A	532	ARG	3.5
1	C	226	ARG	3.5
1	D	416	PHE	3.3
1	A	283	ASN	3.3
1	A	30	SER	3.3
1	B	138	ALA	3.3
1	D	274	TYR	3.3
1	A	375	LEU	3.2
1	B	497	LYS	3.2
1	B	225	LYS	3.2
1	D	384	ALA	3.2
1	A	480	ASN	3.2
1	A	479	LYS	3.2
1	C	503	PHE	3.1
1	D	482	GLN	3.1
1	D	530	VAL	3.1
1	B	227	LYS	3.1
1	A	29	ALA	3.1
1	C	501	GLU	3.0
1	A	474	ARG	3.0
1	D	197	GLY	3.0
1	D	278	ASN	3.0
1	A	478	ALA	3.0
1	B	30	SER	3.0
1	D	280	ASP	2.9
1	C	342	GLY	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	141	GLY	2.8
1	C	233	MET	2.8
1	C	192	VAL	2.8
1	B	495	PHE	2.8
1	A	395	LYS	2.8
1	C	498	ALA	2.7
1	D	369	SER	2.7
1	B	479	LYS	2.7
1	C	224	LYS	2.7
1	D	403	GLN	2.7
1	D	377	VAL	2.7
1	A	481	ILE	2.7
1	D	135	ARG	2.7
1	D	407	GLU	2.7
1	D	374	ALA	2.7
1	B	492	GLY	2.7
1	A	477	TYR	2.7
1	C	230	GLN	2.6
1	D	430	LYS	2.6
1	D	373	PHE	2.6
1	D	30	SER	2.6
1	D	198	PHE	2.6
1	D	100	ASP	2.6
1	D	104	PRO	2.6
1	A	281	GLU	2.6
1	A	379	LEU	2.5
1	C	500	THR	2.5
1	C	33	ARG	2.5
1	A	378	THR	2.5
1	C	117	PRO	2.5
1	C	197	GLY	2.5
1	C	502	PHE	2.5
1	A	278	ASN	2.4
1	D	285	GLY	2.4
1	D	454	ASP	2.4
1	B	369	SER	2.4
1	D	295	THR	2.4
1	A	382	MET	2.4
1	B	32	ASP	2.4
1	C	223	LEU	2.4
1	C	261	CYS	2.4
1	D	475	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	533	PRO	2.3
1	C	531	TRP	2.3
1	A	496	GLU	2.3
1	D	33	ARG	2.3
1	B	341	ALA	2.3
1	C	375	LEU	2.3
1	A	274	TYR	2.3
1	C	341	ALA	2.3
1	C	215	CYS	2.3
1	B	192	VAL	2.3
1	C	252	TYR	2.3
1	D	413	PRO	2.3
1	D	531	TRP	2.3
1	D	375	LEU	2.3
1	B	501	GLU	2.3
1	B	527	ASP	2.2
1	A	498	ALA	2.2
1	B	414	ASP	2.2
1	D	443	CYS	2.2
1	C	506	PHE	2.2
1	A	376	GLY	2.2
1	A	492	GLY	2.2
1	C	260	LEU	2.2
1	A	225	LYS	2.2
1	B	342	GLY	2.2
1	C	221	LYS	2.2
1	B	228	GLY	2.2
1	C	228	GLY	2.2
1	D	408	GLN	2.2
1	C	495	PHE	2.1
1	A	280	ASP	2.1
1	B	190	PHE	2.1
1	C	118	GLN	2.1
1	A	377	VAL	2.1
1	B	118	GLN	2.1
1	C	257	LYS	2.1
1	D	341	ALA	2.1
1	C	32	ASP	2.1
1	D	398	ASN	2.1
1	B	476	VAL	2.1
1	C	190	PHE	2.0
1	C	396	VAL	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	289	PRO	2.0
1	C	371	ASP	2.0
1	C	530	VAL	2.0
1	B	401	LEU	2.0
1	C	138	ALA	2.0
1	A	433	GLU	2.0
1	C	120	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ANW	C	601	22/22	0.78	0.29	0.34	87,95,104,105	0
2	ANW	A	601	22/22	0.88	0.27	0.33	61,72,88,91	0
2	ANW	D	601	22/22	0.91	0.21	-0.40	60,71,89,95	0
2	ANW	B	601	22/22	0.90	0.20	-0.61	73,83,98,104	0
3	CL	A	602	1/1	0.97	0.23	-	60,60,60,60	0
3	CL	D	602	1/1	0.96	0.22	-	64,64,64,64	0
3	CL	B	602	1/1	0.98	0.10	-	63,63,63,63	0
3	CL	C	602	1/1	0.99	0.06	-	60,60,60,60	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.