



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:03 PM GMT

PDB ID : 1WC1
Title : Soluble adenylyl cyclase CyaC from *S. platensis* in complex with Rp- ATPal-phaS
Authors : Steegborn, C.; Litvin, T.N.; Levin, L.R.; Buck, J.; Wu, H.
Deposited on : 2004-11-06
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2910 (1.96-1.92)
Clashscore	102246	3095 (1.96-1.92)
Ramachandran outliers	100387	3062 (1.96-1.92)
Sidechain outliers	100360	3062 (1.96-1.92)
RSRZ outliers	91569	2915 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	<div> <div>5%</div> <div> <div></div> <div>75%</div> <div>12%</div> <div>13%</div> </div> </div>
1	B	226	<div> <div>4%</div> <div> <div></div> <div>68%</div> <div>19%</div> <div>13%</div> </div> </div>
1	C	226	<div> <div>10%</div> <div> <div></div> <div>63%</div> <div>22%</div> <div>15%</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADENYLATE CYCLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	15	0	1
			1513	950	266	285	12			
1	B	196	Total	C	N	O	S	11	0	1
			1505	944	265	284	12			
1	C	193	Total	C	N	O	S	15	0	1
			1477	930	258	278	11			

- Molecule 2 is ADENOSINE-5'-RP-ALPHA-THIO-TRIPHOSPHATE (three-letter code: TAT) (formula: C₁₀H₁₆N₅O₁₂P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		
2	C	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total 2	Mg 2	0	0
3	A	2	Total 2	Mg 2	0	0
3	C	2	Total 2	Mg 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	156	Total 156	O 156	0	0
4	B	127	Total 127	O 127	0	0
4	C	102	Total 102	O 102	0	0

- Molecule 1: ADENYLATE CYCLASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	54.23 Å 78.90 Å 283.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.97 – 1.93 39.45 – 1.89	Depositor EDS
% Data completeness (in resolution range)	87.8 (14.97-1.93) 95.2 (39.45-1.89)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.89 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.198 , 0.229 0.202 , 0.230	Depositor DCC
R_{free} test set	2791 reflections (6.66%)	DCC
Wilson B-factor (Å ²)	21.4	Xtriage
Anisotropy	0.375	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 48935 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4979	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TAT, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1537	0.56	0/2080
1	B	0.34	0/1529	0.57	0/2069
1	C	0.31	0/1500	0.52	0/2030
All	All	0.33	0/4566	0.55	0/6179

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1513	0	1522	22	0
1	B	1505	0	1511	37	0
1	C	1477	0	1487	41	0
2	A	31	0	14	0	0
2	B	31	0	14	0	0
2	C	31	0	14	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	156	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	127	0	0	3	0
4	C	102	0	0	3	0
All	All	4979	0	4562	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1005:ARG:N	1:B:1006:PRO:HD2	1.51	1.17
1:B:1146:ASN:HD21	1:B:1150:ARG:HE	1.16	0.90
1:B:1005:ARG:N	1:B:1006:PRO:CD	2.40	0.84
1:B:1028:LEU:HD13	1:B:1111:VAL:HG21	1.66	0.77
1:C:1016:SER:OG	1:C:1063:ILE:HG22	1.85	0.75
1:B:1130:PHE:HA	1:C:1038:ASN:HD21	1.52	0.75
1:C:1029:GLN:HG2	1:C:1032:GLY:H	1.56	0.70
1:C:1183:LEU:HB2	1:C:1186:ILE:HD11	1.72	0.69
1:B:1146:ASN:ND2	1:B:1150:ARG:HE	1.90	0.69
1:B:1070:PRO:HG2	1:B:1071:GLU:OE2	1.95	0.67
1:B:1132:SER:HB3	1:C:1042:GLY:HA2	1.77	0.66
1:C:1165:MET:O	1:C:1168:GLN:HG2	1.96	0.66
1:C:1046:ARG:O	1:C:1050:GLU:HG3	1.96	0.65
1:B:1080:ARG:NH2	1:B:1081:ARG:HE	1.94	0.65
1:B:1042:GLY:HA2	1:C:1132:SER:HB3	1.79	0.64
1:A:1029:GLN:HG2	1:A:1032:GLY:H	1.63	0.64
1:B:1024:MET:HG2	1:B:1111:VAL:HG13	1.80	0.62
1:C:1098:GLN:O	1:C:1102:GLU:HG2	2.00	0.62
1:A:1031:GLN:O	1:A:1035:GLU:HG3	2.01	0.60
1:C:1176:ILE:HD11	1:C:1196:ASN:HB2	1.83	0.60
1:C:1120:ILE:HB	1:C:1162:VAL:HG12	1.83	0.60
1:C:1101:GLN:HG3	1:C:1106:VAL:HG23	1.84	0.60
1:C:1021:PHE:HA	1:C:1024:MET:SD	2.42	0.59
1:B:1146:ASN:O	1:B:1150:ARG:HG2	2.03	0.59
1:A:1146:ASN:ND2	1:A:1186:ILE:HD11	2.18	0.59
1:B:1175:ILE:HD12	1:B:1178:ARG:NH2	2.18	0.58
1:A:1165:MET:HE1	1:B:1164:ALA:HB1	1.86	0.58
1:C:1029:GLN:O	1:C:1033:VAL:HG23	2.04	0.57
1:C:1093:LEU:HD21	1:C:1115:ARG:HA	1.86	0.57
1:B:1173:GLU:CD	1:B:1173:GLU:H	2.08	0.57
1:C:1019:VAL:HG22	1:C:1117:ARG:NH1	2.20	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1186:ILE:HD12	1:A:1186:ILE:N	2.22	0.55
1:C:1014:LEU:HD23	1:C:1014:LEU:C	2.28	0.54
1:B:1186:ILE:HD12	1:B:1186:ILE:N	2.23	0.54
1:C:1133:GLN:CD	1:C:1133:GLN:H	2.11	0.54
1:B:1116:PHE:H	1:B:1158:ASN:ND2	2.04	0.54
1:B:1184:LYS:HG3	4:B:2105:HOH:O	2.07	0.54
1:A:1183:LEU:HB2	1:A:1186:ILE:HD13	1.91	0.53
4:B:2126:HOH:O	1:C:1146:ASN:HB2	2.08	0.53
1:B:1131:GLY:H	1:C:1038:ASN:ND2	2.07	0.52
1:A:1173:GLU:CD	1:A:1173:GLU:H	2.12	0.52
1:A:1149:ALA:O	1:A:1153:GLU:HG3	2.09	0.52
1:A:1146:ASN:O	1:A:1150:ARG:HG2	2.10	0.51
1:B:1149:ALA:O	1:B:1153:GLU:HG3	2.10	0.51
1:A:1094:GLU:HG2	4:A:2068:HOH:O	2.11	0.51
1:C:1183:LEU:CD1	1:C:1190:VAL:HG11	2.41	0.51
1:A:1109:ASN:CG	1:A:1110:GLU:H	2.13	0.50
1:A:1130:PHE:CE2	1:A:1141:ILE:HD12	2.45	0.50
1:C:1093:LEU:HD22	1:C:1116:PHE:CD2	2.47	0.50
1:B:1091:VAL:HG21	1:B:1199:MET:CE	2.42	0.50
1:B:1035:GLU:CD	1:B:1103:ARG:NH2	2.65	0.49
1:A:1046:ARG:O	1:A:1050:GLU:HG3	2.12	0.49
1:C:1059:VAL:HG11	4:C:2100:HOH:O	2.12	0.49
1:B:1014:LEU:HD23	1:B:1014:LEU:C	2.32	0.49
1:C:1176:ILE:N	1:C:1176:ILE:HD12	2.28	0.49
1:C:1143:PRO:O	1:C:1147:ILE:HG13	2.13	0.49
1:B:1188:GLU:CD	1:B:1188:GLU:H	2.16	0.49
1:A:1029:GLN:N	1:A:1029:GLN:OE1	2.47	0.48
1:B:1187:ASP:HB2	4:C:2011:HOH:O	2.13	0.48
1:A:1189:PRO:HG3	4:A:2132:HOH:O	2.14	0.48
1:B:1051:ASN:O	1:B:1081:ARG:HD2	2.14	0.47
1:A:1028:LEU:HG	1:A:1111:VAL:HG21	1.95	0.47
1:B:1081:ARG:HH11	1:B:1081:ARG:HG3	1.79	0.47
1:A:1014:LEU:C	1:A:1014:LEU:HD23	2.35	0.47
1:B:1028:LEU:CD1	1:B:1111:VAL:HG21	2.41	0.47
1:A:1130:PHE:CZ	1:A:1141:ILE:HD12	2.50	0.46
1:B:1133:GLN:OE1	1:B:1133:GLN:O	2.34	0.46
1:C:1021:PHE:O	1:C:1024:MET:HG2	2.16	0.46
1:B:1070:PRO:HG2	1:B:1071:GLU:CD	2.36	0.45
1:A:1011:ILE:HD13	1:A:1127:VAL:HG21	1.99	0.45
1:C:1196:ASN:OD1	1:C:1198:ASN:N	2.45	0.45
1:C:1177:LYS:HE3	1:C:1194:VAL:HG21	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1176:ILE:CD1	1:C:1196:ASN:HB2	2.47	0.45
1:C:1036:LEU:HD23	1:C:1036:LEU:C	2.37	0.45
1:B:1116:PHE:H	1:B:1158:ASN:HD22	1.65	0.44
1:C:1004:LEU:HG	1:C:1004:LEU:O	2.17	0.44
1:C:1101:GLN:CG	1:C:1106:VAL:HG23	2.46	0.43
1:C:1085:THR:O	1:C:1089:MET:HG3	2.17	0.43
1:C:1117:ARG:HD3	1:C:1155:THR:O	2.18	0.43
1:B:1038:ASN:OD1	1:C:1130:PHE:HA	2.19	0.43
1:A:1005:ARG:HG2	1:A:1005:ARG:NH1	2.35	0.42
1:B:1043:GLU:HG3	1:B:1096:LEU:HD21	2.02	0.42
1:B:1146:ASN:ND2	2:C:1500:TAT:H16	2.34	0.42
1:C:1176:ILE:CD1	1:C:1176:ILE:N	2.83	0.42
1:A:1127:VAL:HG12	1:A:1138:PHE:CZ	2.55	0.42
1:B:1085:THR:O	1:B:1089:MET:HG3	2.20	0.42
1:C:1115:ARG:HD2	1:C:1157:PRO:HB3	2.01	0.41
1:C:1165:MET:HB2	4:C:2082:HOH:O	2.19	0.41
1:C:1115:ARG:HB2	1:C:1158:ASN:OD1	2.21	0.41
1:C:1146:ASN:O	1:C:1150:ARG:HG2	2.21	0.41
1:B:1122:GLN:NE2	4:B:2059:HOH:O	2.53	0.41
1:A:1036:LEU:HD23	1:A:1036:LEU:C	2.42	0.41
1:C:1133:GLN:CD	1:C:1133:GLN:N	2.75	0.40
1:B:1087:ARG:HD3	1:B:1199:MET:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/226 (86%)	189 (97%)	6 (3%)	0	100	100
1	B	194/226 (86%)	188 (97%)	5 (3%)	1 (0%)	34	20
1	C	189/226 (84%)	185 (98%)	3 (2%)	1 (0%)	34	20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	578/678 (85%)	562 (97%)	14 (2%)	2 (0%)	46	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1006	PRO
1	C	1198	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/189 (86%)	162 (99%)	1 (1%)	90	89
1	B	162/189 (86%)	160 (99%)	2 (1%)	78	74
1	C	159/189 (84%)	157 (99%)	2 (1%)	76	71
All	All	484/567 (85%)	479 (99%)	5 (1%)	82	79

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1031	GLN
1	B	1152	GLN
1	B	1165	MET
1	C	1005	ARG
1	C	1152	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1098	GLN
1	A	1109	ASN
1	A	1122	GLN
1	A	1152	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1158	ASN
1	B	1026	ASN
1	B	1122	GLN
1	B	1133	GLN
1	B	1146	ASN
1	B	1158	ASN
1	B	1168	GLN
1	B	1198	ASN
1	C	1038	ASN
1	C	1052	GLN
1	C	1168	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TAT	A	1500	3	22,33,33	1.62	3 (13%)	29,52,52	1.04	2 (6%)
2	TAT	B	1500	3	22,33,33	1.68	3 (13%)	29,52,52	1.12	2 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TAT	C	1500	3	22,33,33	1.65	5 (22%)	29,52,52	1.13	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TAT	A	1500	3	-	0/13/38/38	0/3/3/3
2	TAT	B	1500	3	-	0/13/38/38	0/3/3/3
2	TAT	C	1500	3	-	0/13/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1500	TAT	C3'-C2'	-2.51	1.46	1.53
2	A	1500	TAT	C3'-C2'	-2.39	1.46	1.53
2	C	1500	TAT	C3'-C2'	-2.37	1.46	1.53
2	C	1500	TAT	C5-N7	-2.10	1.32	1.39
2	C	1500	TAT	O4'-C4'	2.11	1.49	1.45
2	A	1500	TAT	C2-N1	3.87	1.41	1.33
2	B	1500	TAT	C2-N1	3.97	1.41	1.33
2	C	1500	TAT	C2-N1	4.06	1.41	1.33
2	C	1500	TAT	O4'-C1'	4.28	1.46	1.41
2	B	1500	TAT	O4'-C1'	4.29	1.46	1.41
2	A	1500	TAT	O4'-C1'	4.43	1.46	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1500	TAT	PB-O3A-PA	-2.73	123.84	133.44
2	C	1500	TAT	PB-O3A-PA	-2.37	125.10	133.44
2	A	1500	TAT	PB-O3A-PA	-2.23	125.60	133.44
2	A	1500	TAT	C4-C5-N7	2.65	111.91	109.48
2	C	1500	TAT	C4-C5-N7	2.67	111.94	109.48
2	B	1500	TAT	C4-C5-N7	2.68	111.94	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1500	TAT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/226 (87%)	0.35	12 (6%) 25 32	12, 23, 56, 67	3 (1%)
1	B	196/226 (86%)	0.25	8 (4%) 41 50	15, 25, 47, 63	3 (1%)
1	C	193/226 (85%)	0.52	22 (11%) 7 10	16, 30, 67, 72	4 (2%)
All	All	586/678 (86%)	0.37	42 (7%) 18 26	12, 26, 57, 72	10 (1%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1199	MET	9.0
1	A	1200	LEU	6.8
1	A	1199	MET	5.7
1	C	1023	ARG	5.3
1	A	1109	ASN	5.3
1	C	1105	LEU	5.0
1	A	1028	LEU	4.9
1	C	1112	PRO	4.6
1	C	1186	ILE	4.4
1	A	1104	GLY	4.2
1	C	1004	LEU	4.1
1	C	1104	GLY	4.1
1	C	1111	VAL	4.0
1	C	1026	ASN	3.8
1	C	1107	GLY	3.5
1	C	1106	VAL	3.5
1	A	1102	GLU	3.4
1	C	1187	ASP	3.4
1	C	1025	SER	3.3
1	B	1199	MET	3.2
1	B	1109	ASN	3.1
1	A	1108	ARG	3.0
1	A	1031	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1005	ARG	2.9
1	A	1110	GLU	2.9
1	C	1102	GLU	2.8
1	A	1107	GLY	2.8
1	C	1101	GLN	2.8
1	C	1172	ASP	2.7
1	A	1105	LEU	2.7
1	B	1023	ARG	2.6
1	C	1198	ASN	2.5
1	B	1110	GLU	2.5
1	C	1188	GLU	2.4
1	B	1046	ARG	2.4
1	C	1028	LEU	2.3
1	C	1098	GLN	2.3
1	B	1187	ASP	2.3
1	C	1134	GLU	2.2
1	A	1188	GLU	2.2
1	B	1184	LYS	2.1
1	B	1188	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	TAT	A	1500	31/31	0.96	0.14	0.31	16,24,31,32	0
2	TAT	B	1500	31/31	0.96	0.09	-0.38	21,27,34,36	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	TAT	C	1500	31/31	0.93	0.10	-0.48	24,37,47,48	0
3	MG	A	1502	1/1	0.99	0.09	-1.08	16,16,16,16	0
3	MG	C	1502	1/1	0.91	0.06	-2.00	33,33,33,33	0
3	MG	B	1502	1/1	0.96	0.06	-2.85	18,18,18,18	0
3	MG	C	1501	1/1	0.98	0.06	-	39,39,39,39	0
3	MG	B	1501	1/1	0.99	0.07	-	22,22,22,22	0
3	MG	A	1501	1/1	0.98	0.08	-	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.