



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:50 PM GMT

PDB ID : 4WEA  
Title : Structure and receptor binding prefereneces of recombinant human A(H3N2) virus hemagglutinins  
Authors : Yang, H.; Carney, P.J.; Chang, J.C.; Guo, Z.; Villanueva, J.M.; Stevens, J.  
Deposited on : 2014-09-09  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	<div> <div>7%</div> <div>84%</div> <div>11%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	602	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	607	X	-	-	-
2	NAG	A	608	-	-	-	X
4	SIA	A	610	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4187 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	491	3881	2424	689	750	18	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	ALA	-	expression tag	UNP R9U684
A	-3	ASP	-	expression tag	UNP R9U684
A	-2	LEU	-	expression tag	UNP R9U684
A	-1	GLY	-	expression tag	UNP R9U684
A	0	SER	-	expression tag	UNP R9U684
A	504	SER	-	expression tag	UNP R9U684
A	505	GLY	-	expression tag	UNP R9U684
A	506	ARG	-	expression tag	UNP R9U684
A	507	LEU	-	expression tag	UNP R9U684
A	508	VAL	-	expression tag	UNP R9U684
A	509	PRO	-	expression tag	UNP R9U684
A	510	ARG	-	expression tag	UNP R9U684

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is SUGAR (O-SIALIC ACID) (three-letter code: SIA) (formula:  $C_{11}H_{19}NO_9$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			20	11	1	8		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	160	Total	O	0	0
			160	160		



- Molecule 1: Hemagglutinin



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.73Å 100.73Å 382.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.04 – 2.20 43.34 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (35.04-2.20) 96.9 (43.34-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.88 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.190 , 0.223 0.187 , 0.222	Depositor DCC
$R_{free}$ test set	1869 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.6	Xtriage
Anisotropy	0.539	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 38438 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4187	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/3961	0.52	0/5362

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3881	0	3753	42	0
2	A	98	0	91	2	0
3	A	28	0	25	1	0
4	A	20	0	17	1	0
5	A	160	0	0	7	0
All	All	4187	0	3886	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:ARG:NH2	5:A:704:HOH:O	2.21	0.73
1:A:119:GLU:OE1	1:A:261:ARG:NH1	2.21	0.73
3:A:605:NAG:H82	2:A:607:NAG:H61	1.72	0.71
1:A:245:ILE:HD11	1:A:251:LEU:HD21	1.72	0.71
1:A:127:TRP:HB2	1:A:132:GLN:HE21	1.56	0.70
1:A:104:ASP:OD2	5:A:701:HOH:O	2.09	0.69
1:A:348:ASP:OD2	5:A:702:HOH:O	2.12	0.67
1:A:137:SER:OG	5:A:855:HOH:O	2.12	0.66
1:A:45:ASN:O	5:A:803:HOH:O	2.15	0.64
1:A:97:CYS:O	1:A:224:ARG:NH1	2.31	0.64
1:A:394:GLN:OE1	5:A:703:HOH:O	2.15	0.63
1:A:180:TRP:HE1	1:A:235:THR:HG22	1.64	0.63
1:A:127:TRP:CZ3	1:A:166:VAL:HG11	2.38	0.58
1:A:180:TRP:HE1	1:A:235:THR:CG2	2.18	0.57
1:A:152:ASN:N	1:A:253:ALA:O	2.37	0.56
1:A:173:GLN:N	1:A:173:GLN:OE1	2.31	0.56
1:A:220:ARG:HD2	1:A:229:ARG:HG3	1.90	0.54
1:A:127:TRP:HB2	1:A:132:GLN:NE2	2.23	0.53
1:A:102:VAL:HG22	1:A:232:ILE:HB	1.91	0.52
1:A:245:ILE:HD11	1:A:251:LEU:CD2	2.42	0.49
1:A:84:TRP:HZ3	1:A:118:LEU:HG	1.77	0.49
1:A:348:ASP:N	1:A:348:ASP:OD1	2.45	0.49
1:A:161:TYR:HE2	1:A:249:GLY:HA2	1.77	0.49
1:A:95:SER:O	1:A:224:ARG:NH2	2.46	0.48
1:A:161:TYR:HE2	1:A:249:GLY:N	2.12	0.47
1:A:161:TYR:CE2	1:A:249:GLY:HA2	2.48	0.47
1:A:141:ARG:O	1:A:143:SER:N	2.45	0.47
1:A:52:CYS:HB3	1:A:277:CYS:O	2.15	0.46
1:A:185:PRO:HD2	1:A:217:ILE:HD13	1.96	0.46
1:A:146:SER:OG	1:A:147:PHE:N	2.48	0.46
1:A:99:PRO:HG2	1:A:226:ILE:HB	1.97	0.45
1:A:169:PRO:HB3	1:A:242:ILE:HG12	1.99	0.43
1:A:186:GLY:HA2	1:A:218:GLY:O	2.18	0.43
1:A:141:ARG:NH1	1:A:146:SER:OG	2.51	0.43
1:A:161:TYR:HE2	1:A:249:GLY:CA	2.32	0.43
1:A:266:SER:OG	1:A:267:ILE:N	2.51	0.42
1:A:135:THR:O	4:A:610:SIA:H4	2.20	0.42
1:A:355:HIS:HE1	5:A:712:HOH:O	2.02	0.42
1:A:199:SER:OG	1:A:200:GLY:N	2.54	0.41
1:A:67:ILE:HG13	1:A:105:TYR:CE1	2.56	0.41
1:A:135:THR:HG23	1:A:145:ASN:HB3	2.03	0.40
1:A:34:ILE:HD11	1:A:321:ARG:NE	2.36	0.40

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:GLN:HB2	2:A:604:NAG:H82	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	487/515 (95%)	463 (95%)	21 (4%)	3 (1%)	30 29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	142	ARG
1	A	199	SER
1	A	196	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	428/448 (96%)	420 (98%)	8 (2%)	65 77

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	135	THR
1	A	197	GLN
1	A	204	VAL
1	A	235	THR
1	A	245	ILE
1	A	454	GLN
1	A	483	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	ASN
1	A	132	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	605	1,3	14,14,15	0.46	0	15,19,21	0.58	0
3	NAG	A	606	3	14,14,15	0.25	0	15,19,21	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	605	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	606	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	605	NAG	1	0

## 5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	A	601	1	14,14,15	0.30	0	15,19,21	0.43	0
2	NAG	A	602	1	14,14,15	0.20	0	15,19,21	0.75	1 (6%)
2	NAG	A	603	1	14,14,15	0.20	0	15,19,21	0.32	0
2	NAG	A	604	1	14,14,15	0.33	0	15,19,21	0.32	0
2	NAG	A	607	1	14,14,15	0.52	0	15,19,21	0.63	0
2	NAG	A	608	1	14,14,15	0.40	0	15,19,21	0.64	1 (6%)
2	NAG	A	609	1	14,14,15	0.38	0	15,19,21	0.35	0
4	SIA	A	610	-	16,20,21	0.24	0	18,28,31	1.31	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1	-	0/6/23/26	0/1/1/1
2	NAG	A	602	1	-	0/6/23/26	0/1/1/1
2	NAG	A	603	1	-	0/6/23/26	0/1/1/1
2	NAG	A	604	1	-	0/6/23/26	0/1/1/1
2	NAG	A	607	1	1/1/5/7	0/6/23/26	0/1/1/1
2	NAG	A	608	1	-	0/6/23/26	0/1/1/1
2	NAG	A	609	1	-	0/6/23/26	0/1/1/1
4	SIA	A	610	-	-	0/14/34/38	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	608	NAG	C1-O5-C5	2.36	115.24	112.25
2	A	602	NAG	C1-O5-C5	2.47	115.39	112.25
4	A	610	SIA	O6-C6-C5	2.83	113.12	108.48
4	A	610	SIA	C3-C4-C5	2.84	114.64	111.47

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	607	NAG	C1

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	604	NAG	1	0
2	A	607	NAG	1	0
4	A	610	SIA	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	491/515 (95%)	0.27	37 (7%) 17 17	28, 57, 117, 153	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	ALA	8.4
1	A	158	ASN	5.0
1	A	96	ASN	4.8
1	A	225	ASN	4.6
1	A	193	PHE	4.6
1	A	163	ALA	4.3
1	A	130	VAL	4.2
1	A	162	PRO	4.0
1	A	64	CYS	3.9
1	A	142	ARG	3.9
1	A	223	ILE	3.9
1	A	159	PHE	3.6
1	A	8	ASN	3.6
1	A	226	ILE	3.6
1	A	186	GLY	3.5
1	A	224	ARG	3.4
1	A	197	GLN	3.2
1	A	167	THR	3.2
1	A	161	TYR	3.1
1	A	192	ILE	3.1
1	A	129	GLY	3.1
1	A	145	ASN	3.0
1	A	173	GLN	2.9
1	A	242	ILE	2.7
1	A	141	ARG	2.6
1	A	160	LYS	2.5
1	A	207	LYS	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	98	TYR	2.4
1	A	157	LEU	2.4
1	A	22	ASN	2.2
1	A	239	PRO	2.2
1	A	386	GLY	2.2
1	A	144	ASN	2.1
1	A	94	TYR	2.0
1	A	140	ILE	2.0
1	A	194	LEU	2.0
1	A	137	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	606	14/15	0.86	0.24	-	93,98,102,102	0
3	NAG	A	605	14/15	0.95	0.19	-	78,90,99,100	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	602	14/15	0.79	0.27	6.69	63,72,76,78	0
4	SIA	A	610	20/21	0.41	0.56	4.18	135,146,151,152	0
2	NAG	A	608	14/15	0.88	0.20	2.18	51,63,71,72	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	607	14/15	0.80	0.33	1.63	114,124,133,133	0
2	NAG	A	601	14/15	0.83	0.56	-	91,104,112,118	0
2	NAG	A	609	14/15	0.65	0.43	-	85,98,105,107	0
2	NAG	A	603	14/15	0.89	0.35	-	74,90,95,99	0
2	NAG	A	604	14/15	0.78	0.30	-	88,99,106,110	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.