



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:11 PM GMT

PDB ID : 3WFC
Title : Reduced and carbonmonoxide-bound cytochrome c-dependent nitric oxide reductase (cNOR) from Pseudomonas aeruginosa in complex with antibody fragment
Authors : Sato, N.; Ishii, S.; Hino, T.; Sugimoto, H.; Fukumori, Y.; Shiro, Y.; Tosha, T.
Deposited on : 2013-07-18
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	213	<div> <div>3%</div> <div>78% 17% .</div> </div>
2	H	225	<div> <div>3%</div> <div>76% 20% .</div> </div>
3	B	465	<div> <div>6%</div> <div>74% 18% 5% .</div> </div>
4	C	146	<div> <div>3%</div> <div>84% 9% . .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	CMO	B	804	-	-	X	-
8	10M	B	806	-	-	-	X
8	10M	B	807	-	-	-	X

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 8541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody fab fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	213	Total	C	N	O	S	0	0	0
			1669	1047	277	338	7			

- Molecule 2 is a protein called antibody fab fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	225	Total	C	N	O	S	0	0	0
			1692	1065	280	338	9			

- Molecule 3 is a protein called Nitric oxide reductase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	449	Total	C	N	O	S	0	0	0
			3576	2416	563	572	25			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ARG	DELETION	UNP Q59647

- Molecule 4 is a protein called Nitric oxide reductase subunit C.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	C	142	Total	C	N	O	S	0	0	0
			1123	720	195	202	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	100	LYS	ASN	CONFLICT	UNP Q59646

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (for-

mula: C₃₄H₃₂FeN₄O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 6 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Fe	0	0
			1	1		

- Molecule 7 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			2	1	1		
7	B	1	Total	C	O	0	0
			2	1	1		

- Molecule 8 is SUGAR (DECYL 4-O-ALPHA-D-GLUCOPYRANOSYL-1-THIO-BETA-D-GLUCOPYRANOSIDE) (three-letter code: 10M) (formula: C₂₂H₄₂O₁₀S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	O	S	0	0
			33	22	10	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	O	S	0	0
			33	22	10	1		

- Molecule 9 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total	Ca	0	0
			1	1		

- Molecule 10 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	L	90	Total	O	0	0
			90	90		
11	H	86	Total	O	0	0
			86	86		
11	B	52	Total	O	0	0
			52	52		
11	C	52	Total	O	0	0
			52	52		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody fab fragment light chain




- Molecule 2: antibody fab fragment heavy chain



- Molecule 3: Nitric oxide reductase subunit B



- Molecule 4: Nitric oxide reductase subunit C



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.67Å 107.07Å 196.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.37 – 2.50 33.35 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.6 (33.37-2.50) 98.7 (33.35-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.191 , 0.232 0.191 , 0.233	Depositor DCC
R_{free} test set	3348 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	47.4	Xtriage
Anisotropy	0.392	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 65789 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8541	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, 10M, CA, FE, HEC, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.76	0/1709	0.86	2/2317 (0.1%)
2	H	0.75	0/1735	0.89	4/2367 (0.2%)
3	B	0.63	0/3693	0.74	1/5039 (0.0%)
4	C	0.60	0/1153	0.77	1/1559 (0.1%)
All	All	0.68	0/8290	0.80	8/11282 (0.1%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	H	224	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	L	4	MET	CB-CG-SD	-5.70	95.30	112.40
2	H	224	ARG	NE-CZ-NH2	-5.70	117.45	120.30
2	H	206	CYS	CA-CB-SG	-5.56	103.99	114.00
3	B	198	ASP	CB-CG-OD1	5.40	123.16	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1669	0	1606	34	0
2	H	1692	0	1647	30	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	3576	0	3619	87	0
4	C	1123	0	1092	25	0
5	B	86	0	60	10	0
6	B	1	0	0	0	0
7	B	4	0	0	3	0
8	B	66	0	84	4	0
9	B	1	0	0	0	0
10	C	43	0	30	1	0
11	B	52	0	0	1	0
11	C	52	0	0	1	0
11	H	86	0	0	2	0
11	L	90	0	0	1	0
All	All	8541	0	8138	167	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:107:LYS:HB2	11:H:319:HOH:O	1.63	0.99
3:B:121:ALA:HA	3:B:132:MET:HE1	1.47	0.96
3:B:211:GLU:OE2	7:B:805:CMO:C	2.13	0.95
8:B:806:10M:H17	8:B:807:10M:O8	1.69	0.93
2:H:48:ILE:HG21	2:H:81:MET:HE1	1.51	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	211/213 (99%)	200 (95%)	10 (5%)	1 (0%)	34	55
2	H	223/225 (99%)	211 (95%)	9 (4%)	3 (1%)	15	26
3	B	447/465 (96%)	426 (95%)	21 (5%)	0	100	100
4	C	140/146 (96%)	136 (97%)	4 (3%)	0	100	100
All	All	1021/1049 (97%)	973 (95%)	44 (4%)	4 (0%)	39	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	141	ASP
1	L	211	ARG
2	H	183	SER
2	H	144	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	189/189 (100%)	173 (92%)	16 (8%)	13	25
2	H	192/192 (100%)	168 (88%)	24 (12%)	6	10
3	B	360/371 (97%)	319 (89%)	41 (11%)	7	13
4	C	116/120 (97%)	109 (94%)	7 (6%)	24	43
All	All	857/872 (98%)	769 (90%)	88 (10%)	9	17

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	219	LYS
3	B	115	VAL
4	C	24	LEU
3	B	30	GLN
3	B	92	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such

sidechains are listed below:

Mol	Chain	Res	Type
3	B	54	ASN
3	B	126	ASN
4	C	102	GLN
3	B	62	ASN
3	B	139	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	HEM	B	801	9,3	30,50,50	2.48	8 (26%)	24,82,82	3.27	15 (62%)
5	HEM	B	802	9,3,7	30,50,50	2.28	4 (13%)	24,82,82	3.53	11 (45%)
7	CMO	B	804	5	0,1,1	0.00	-	0,0,0	0.00	-
7	CMO	B	805	6	0,1,1	0.00	-	0,0,0	0.00	-
8	10M	B	806	-	34,34,34	1.13	3 (8%)	43,45,45	1.34	6 (13%)
8	10M	B	807	-	34,34,34	1.59	7 (20%)	43,45,45	1.24	6 (13%)
10	HEC	C	201	4	24,50,50	2.51	5 (20%)	19,82,82	2.83	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEM	B	801	9,3	-	0/10/54/54	0/0/8/8
5	HEM	B	802	9,3,7	-	0/10/54/54	0/0/8/8
7	CMO	B	804	5	-	0/0/0/0	0/0/0/0
7	CMO	B	805	6	-	0/0/0/0	0/0/0/0
8	10M	B	806	-	-	0/19/59/59	0/2/2/2
8	10M	B	807	-	-	0/19/59/59	0/2/2/2
10	HEC	C	201	4	-	0/6/54/54	0/0/8/8

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	802	HEM	C3B-C4B	-9.07	1.43	1.51
5	B	801	HEM	C3B-C4B	-9.07	1.43	1.51
10	C	201	HEC	C3C-C2C	-7.39	1.33	1.40
10	C	201	HEC	C3B-C2B	-5.93	1.34	1.40
5	B	801	HEM	C3D-C4D	-5.35	1.44	1.51

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	802	HEM	C3B-CAB-CBB	-9.85	109.35	124.46
10	C	201	HEC	CBB-CAB-C3B	-9.24	106.82	127.35
5	B	801	HEM	C3C-CAC-CBC	-8.92	110.78	124.46
5	B	802	HEM	CBA-CAA-C2A	-4.61	104.27	112.53
5	B	801	HEM	CMA-C3A-C4A	-4.39	121.10	128.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	801	HEM	5	0
5	B	802	HEM	5	0
7	B	804	CMO	2	0
7	B	805	CMO	1	0
8	B	806	10M	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	807	10M	4	0
10	C	201	HEC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	213/213 (100%)	-0.42	2 (0%) 85 88	33, 44, 62, 108	0
2	H	225/225 (100%)	-0.29	6 (2%) 58 62	33, 43, 72, 134	0
3	B	449/465 (96%)	-0.00	27 (6%) 25 28	37, 58, 96, 124	0
4	C	142/146 (97%)	-0.01	4 (2%) 56 61	35, 56, 80, 99	0
All	All	1029/1049 (98%)	-0.15	39 (3%) 44 49	33, 52, 87, 134	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	381	ARG	5.7
3	B	299	ARG	5.3
2	H	143	THR	5.3
2	H	225	GLY	4.1
1	L	213	GLU	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
8	10M	B	807	33/33	0.82	0.41	8.69	73,91,107,108	0
8	10M	B	806	33/33	0.93	0.25	3.54	65,74,87,89	0
10	HEC	C	201	43/43	0.99	0.16	0.54	35,37,39,41	0
5	HEM	B	801	43/43	0.97	0.18	0.36	37,40,45,49	0
5	HEM	B	802	43/43	0.98	0.21	0.30	36,40,43,44	0
7	CMO	B	805	2/2	0.99	0.21	0.10	39,39,39,40	0
7	CMO	B	804	2/2	0.99	0.19	-0.88	40,40,40,41	0
9	CA	B	808	1/1	1.00	0.20	-1.42	46,46,46,46	0
6	FE	B	803	1/1	0.98	0.12	-	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.