



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 2, 2016 – 12:00 AM GMT

PDB ID : 7WGA
Title : 2.2 ANGSTROMS RESOLUTION STRUCTURE ANALYSIS OF TWO REFINED N-ACETYLNEURAMINYLLACTOSE-WHEAT GERM AGGLUTININ ISOLECTIN COMPLEXES
Authors : Wright, C.S.
Deposited on : 1990-04-03
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	171	
1	B	171	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2530 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WHEAT GERM LECTIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	171	Total	C	N	O	S	0	0	0
			1150	667	211	238	34			
1	B	171	Total	C	N	O	S	0	0	0
			1163	678	213	238	34			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	104	Total	O	0	0
			104	104		
2	B	113	Total	O	0	0
			113	113		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: WHEAT GERM LECTIN



• Molecule 1: WHEAT GERM LECTIN



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	51.22Å 73.60Å 91.42Å 90.00° 98.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.172 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2530	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.20	3/1162 (0.3%)	1.78	19/1556 (1.2%)
1	B	1.17	4/1177 (0.3%)	1.79	23/1576 (1.5%)
All	All	1.18	7/2339 (0.3%)	1.78	42/3132 (1.3%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	115	GLU	CD-OE2	9.11	1.35	1.25
1	A	5	GLU	CD-OE2	7.71	1.34	1.25
1	B	60	CYS	CB-SG	-6.61	1.71	1.82
1	B	72	GLU	CD-OE2	6.01	1.32	1.25
1	B	11	GLU	CD-OE2	5.52	1.31	1.25
1	B	5	GLU	CD-OE2	5.43	1.31	1.25
1	A	24	CYS	CB-SG	-5.08	1.73	1.81

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	CYS	CA-CB-SG	10.15	132.28	114.00
1	A	164	CYS	CA-CB-SG	7.81	128.06	114.00
1	A	139	ARG	NE-CZ-NH2	-7.67	116.47	120.30
1	B	84	ARG	NE-CZ-NH1	-7.53	116.53	120.30
1	A	129	ASP	CB-CG-OD1	7.14	124.73	118.30
1	B	68	GLY	O-C-N	7.13	134.11	122.70
1	A	170	ASP	CB-CG-OD2	-6.86	112.12	118.30
1	B	45	ARG	CD-NE-CZ	6.86	133.20	123.60
1	A	129	ASP	CB-CG-OD2	-6.81	112.17	118.30
1	B	2	ARG	NE-CZ-NH1	6.49	123.54	120.30
1	B	10	MET	CA-CB-CG	6.41	124.19	113.30
1	B	20	GLN	CA-CB-CG	-6.23	99.70	113.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	PHE	CB-CG-CD2	-6.03	116.58	120.80
1	B	45	ARG	NE-CZ-NH1	6.03	123.31	120.30
1	B	21	TYR	CB-CG-CD1	5.92	124.55	121.00
1	B	20	GLN	OE1-CD-NE2	5.73	135.08	121.90
1	B	86	ASP	CB-CG-OD1	5.73	123.45	118.30
1	A	147	CYS	CA-CB-SG	-5.72	103.71	114.00
1	A	109	PHE	N-CA-CB	5.71	120.87	110.60
1	A	45	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	B	45	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	B	83	CYS	CA-CB-SG	5.59	124.06	114.00
1	B	155	ILE	N-CA-CB	5.59	123.65	110.80
1	A	20	GLN	OE1-CD-NE2	5.57	134.72	121.90
1	B	21	TYR	CB-CG-CD2	-5.56	117.67	121.00
1	B	139	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	A	79	GLN	OE1-CD-NE2	5.44	134.42	121.90
1	A	102	LEU	CA-CB-CG	5.44	127.81	115.30
1	B	73	TYR	CB-CG-CD1	-5.42	117.75	121.00
1	B	69	PHE	N-CA-CB	5.29	120.12	110.60
1	B	152	SER	N-CA-CB	5.26	118.39	110.50
1	A	3	CYS	CA-CB-SG	5.24	123.43	114.00
1	A	84	ARG	CD-NE-CZ	-5.19	116.33	123.60
1	B	170	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	A	69	PHE	N-CA-CB	5.13	119.83	110.60
1	A	16	LEU	CB-CA-C	5.11	119.90	110.20
1	A	55	CYS	N-CA-CB	5.10	119.78	110.60
1	B	73	TYR	CA-CB-CG	-5.09	103.73	113.40
1	B	132	CYS	C-N-CA	5.06	132.92	122.30
1	A	102	LEU	CB-CG-CD2	5.05	119.58	111.00
1	B	41	TRP	CA-CB-CG	-5.04	104.13	113.70
1	B	69	PHE	C-N-CA	5.01	132.82	122.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1150	0	969	25	0
1	B	1163	0	988	33	1
2	A	104	0	0	1	0
2	B	113	0	0	9	0
All	All	2530	0	1957	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ASN:HA	1:A:44:LYS:HZ2	1.18	1.05
1:B:149:LYS:HE2	1:B:150:TRP:CE2	2.19	0.77
1:A:7:GLY:O	1:A:8:SER:HB2	1.86	0.76
1:B:44:LYS:NZ	2:B:180:HOH:O	2.19	0.74
1:B:150:TRP:CD1	2:B:253:HOH:O	2.47	0.67
1:B:124:GLY:O	1:B:126:CYS:N	2.24	0.67
1:B:92:GLN:NE2	1:B:106:GLN:O	2.29	0.65
1:B:150:TRP:HD1	2:B:253:HOH:O	1.78	0.63
1:B:88:LYS:HG3	1:B:109:PHE:CD1	2.35	0.61
1:A:31:CYS:HB3	1:A:41:TRP:CD1	2.37	0.60
1:B:129:ASP:HA	2:B:239:HOH:O	2.03	0.59
1:A:44:LYS:NZ	1:A:56:THR:OG1	2.23	0.58
1:B:7:GLY:O	1:B:8:SER:HB2	2.05	0.57
1:B:7:GLY:O	1:B:10:MET:HB2	2.06	0.55
1:A:104:CYS:HB3	1:A:122:GLN:HB2	1.89	0.55
1:B:98:CYS:HB3	1:B:102:LEU:HB2	1.88	0.55
1:A:139:ARG:NH2	2:A:221:HOH:O	2.28	0.54
1:B:138:GLY:HA3	2:B:173:HOH:O	2.08	0.53
1:B:146:CYS:SG	1:B:168:GLY:O	2.67	0.53
1:B:88:LYS:HG3	1:B:109:PHE:HD1	1.74	0.52
1:A:63:GLN:HB3	1:A:77:GLY:HA3	1.90	0.52
1:B:105:SER:O	1:B:122:GLN:NE2	2.32	0.52
1:B:77:GLY:O	1:B:78:CYS:C	2.49	0.52
1:B:60:CYS:HA	1:B:80:GLY:O	2.10	0.51
1:A:20:GLN:HB3	1:A:34:GLY:HA3	1.93	0.50
1:B:88:LYS:HA	1:B:108:GLY:O	2.11	0.50
1:A:147:CYS:O	1:A:164:CYS:HA	2.12	0.49
1:A:146:CYS:HA	1:A:167:GLY:HA3	1.94	0.48
1:A:89:CYS:O	1:A:108:GLY:HA2	2.14	0.48
1:A:128:THR:OG1	1:A:130:LYS:HE3	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:LEU:HD12	1:A:26:MET:CE	2.44	0.47
1:A:2:ARG:HA	1:A:2:ARG:HD2	1.66	0.47
1:A:16:LEU:HD12	1:A:26:MET:HE3	1.97	0.47
1:B:54:THR:HG21	1:B:100:ASN:OD1	2.16	0.46
1:A:12:CYS:HB3	1:A:13:PRO:HD2	1.97	0.46
1:A:88:LYS:HA	1:A:108:GLY:O	2.15	0.46
1:B:37:ASN:HA	1:B:44:LYS:NZ	2.31	0.45
1:B:46:CYS:CA	1:B:50:ALA:HB3	2.46	0.45
1:B:130:LYS:HB3	1:B:130:LYS:HE2	1.66	0.45
1:A:18:CYS:HB2	1:A:37:ASN:ND2	2.32	0.45
1:A:18:CYS:HB2	1:A:37:ASN:HD21	1.82	0.44
1:A:136:ALA:O	1:A:138:GLY:N	2.51	0.44
1:B:88:LYS:HG3	1:B:109:PHE:CE1	2.52	0.44
1:A:160:CYS:O	1:A:170:ASP:N	2.36	0.44
1:A:136:ALA:C	1:A:138:GLY:H	2.20	0.43
1:B:63:GLN:HB3	1:B:77:GLY:HA3	2.01	0.43
1:B:1:PCA:N	2:B:198:HOH:O	2.52	0.43
1:B:37:ASN:HA	1:B:44:LYS:HZ2	1.82	0.43
1:B:106:GLN:N	2:B:220:HOH:O	2.25	0.42
1:A:45:ARG:HD3	1:A:66:TYR:CE1	2.55	0.42
1:B:44:LYS:HB3	2:B:265:HOH:O	2.19	0.42
1:B:149:LYS:HE2	1:B:150:TRP:CZ2	2.53	0.41
1:B:56:THR:O	1:B:57:ASN:HB2	2.19	0.41
1:B:16:LEU:HA	1:B:16:LEU:HD12	1.85	0.41
1:B:45:ARG:HA	1:B:45:ARG:HD2	1.85	0.41
1:A:2:ARG:HG3	1:A:2:ARG:NH1	2.35	0.41
1:A:136:ALA:C	1:A:138:GLY:N	2.75	0.40
1:B:24:CYS:HB2	2:B:219:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:21:TYR:OH	1:B:86:ASP:OD2[3_455]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/171 (99%)	152 (90%)	13 (8%)	4 (2%)	7	2
1	B	169/171 (99%)	147 (87%)	18 (11%)	4 (2%)	7	2
All	All	338/342 (99%)	299 (88%)	31 (9%)	8 (2%)	7	2

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	125	ALA
1	B	75	GLY
1	B	96	LYS
1	A	125	ALA
1	A	137	GLY
1	B	39	ALA
1	A	119	GLY
1	A	32	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/119 (98%)	107 (92%)	9 (8%)	16	10
1	B	118/119 (99%)	107 (91%)	11 (9%)	11	6
All	All	234/238 (98%)	214 (92%)	20 (8%)	13	8

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	GLU
1	A	10	MET
1	A	16	LEU
1	A	37	ASN
1	A	79	GLN
1	A	88	LYS
1	A	96	LYS
1	A	139	ARG
1	A	140	VAL
1	B	5	GLU
1	B	16	LEU
1	B	19	SER
1	B	37	ASN
1	B	49	GLN
1	B	54	THR
1	B	79	GLN
1	B	96	LYS
1	B	99	PRO
1	B	114	SER
1	B	149	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	14	ASN
1	A	37	ASN
1	A	59	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	1	1	7,8,9	0.82	0	9,10,12	1.45	1 (11%)
1	PCA	B	1	1	7,8,9	0.82	1 (14%)	9,10,12	2.26	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1
1	PCA	B	1	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1	PCA	O-C	2.04	1.29	1.19

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	O-C-CA	-4.03	114.79	125.44
1	B	1	PCA	O-C-CA	-3.06	117.35	125.44
1	B	1	PCA	CB-CA-C	6.00	120.96	112.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	1	PCA	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.