



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:08 PM GMT

PDB ID : 1WKW
Title : Crystal structure of the ternary complex of eIF4E-m7GpppA-4EBP1 peptide
Authors : Tomoo, K.; Matsushita, Y.; Fujisaki, H.; Shen, X.; Miyagawa, H.; Kitamura, K.; Miura, K.; Ishida, T.
Deposited on : 2004-06-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	
2	B	20	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1875 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Eukaryotic translation initiation factor 4E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	0	0
			1571	1000	277	288	6			

- Molecule 2 is a protein called Eukaryotic translation initiation factor 4E binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	20	Total	C	N	O	S	0	0	0
			164	102	31	29	2			

- Molecule 3 is P1-7-METHYLGUANOSINE-P3-ADENOSINE-5',5'-TRIPHOSPHATE (three-letter code: GTA) (formula: $C_{21}H_{30}N_{10}O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			51	21	10	17	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	87	Total 87	O 87	0	0
4	B	2	Total 2	O 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Eukaryotic translation initiation factor 4E



- Molecule 2: Eukaryotic translation initiation factor 4E binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	87.77Å 87.77Å 38.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 62.06 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.10) 74.1 (62.06-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 2.10Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.218 , 0.270 0.232 , 0.289	Depositor DCC
R_{free} test set	1330 reflections (10.32%)	DCC
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.541	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 25.6	EDS
Estimated twinning fraction	0.057 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 12894 reflections	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	1875	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1611	0.64	0/2180
2	B	0.46	0/167	0.78	0/220
All	All	0.40	0/1778	0.66	0/2400

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1571	0	1535	79	0
2	B	164	0	165	31	0
3	A	51	0	26	0	0
4	A	87	0	0	9	0
4	B	2	0	0	0	0
All	All	1875	0	1726	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:GLN:HE22	2:B:51:ARG:HB3	1.07	1.10
1:A:40:GLN:NE2	2:B:51:ARG:HB3	1.74	0.99
1:A:200:HIS:O	1:A:203:THR:HG23	1.63	0.97
1:A:40:GLN:HE21	1:A:40:GLN:HA	1.31	0.93
1:A:108:LYS:HD3	1:A:109:ARG:HG3	1.58	0.86
2:B:63:ARG:HH11	2:B:63:ARG:HB2	1.42	0.84
1:A:59:ASN:HA	4:A:1051:HOH:O	1.76	0.84
2:B:61:GLU:C	2:B:63:ARG:H	1.83	0.82
2:B:63:ARG:HH11	2:B:63:ARG:CB	1.98	0.76
1:A:40:GLN:HE22	2:B:51:ARG:CB	1.92	0.76
1:A:50:ASN:HD21	1:A:52:LYS:HG3	1.49	0.76
1:A:203:THR:HG22	4:A:1086:HOH:O	1.87	0.73
1:A:30:ASN:HD22	1:A:30:ASN:C	1.93	0.71
1:A:40:GLN:NE2	2:B:51:ARG:HE	1.88	0.71
1:A:206:LYS:HD3	1:A:207:SER:N	2.07	0.70
1:A:35:ILE:HD12	1:A:36:LYS:O	1.94	0.68
1:A:30:ASN:HD22	1:A:31:PRO:N	1.92	0.68
1:A:73:TRP:HB3	2:B:64:ASN:ND2	2.08	0.67
1:A:204:ALA:HB1	1:A:209:SER:OG	1.99	0.63
1:A:108:LYS:CD	1:A:109:ARG:HG3	2.29	0.62
2:B:56:ARG:HG2	2:B:56:ARG:NH1	2.16	0.60
1:A:52:LYS:HB3	1:A:52:LYS:NZ	2.15	0.60
1:A:210:THR:HB	1:A:212:LYS:NZ	2.17	0.60
1:A:50:ASN:ND2	1:A:52:LYS:HG3	2.16	0.59
2:B:52:ILE:CD1	2:B:52:ILE:H	2.13	0.59
1:A:73:TRP:CD2	2:B:64:ASN:HB2	2.37	0.59
2:B:52:ILE:N	2:B:52:ILE:HD13	2.17	0.59
2:B:63:ARG:NH1	2:B:63:ARG:CB	2.65	0.58
2:B:56:ARG:HG2	2:B:56:ARG:HH11	1.69	0.57
1:A:57:GLN:HB2	4:A:1057:HOH:O	2.03	0.57
1:A:207:SER:HA	4:A:1060:HOH:O	2.04	0.56
1:A:49:LYS:NZ	1:A:86:MET:SD	2.77	0.56
2:B:52:ILE:HD13	2:B:52:ILE:H	1.68	0.56
1:A:40:GLN:HE22	2:B:51:ARG:HE	1.52	0.56
1:A:94:PHE:HE1	1:A:153:VAL:HG12	1.71	0.56
2:B:56:ARG:HG3	2:B:57:LYS:N	2.19	0.56
2:B:57:LYS:O	2:B:61:GLU:HB2	2.06	0.56
1:A:206:LYS:C	1:A:206:LYS:HD3	2.27	0.56
1:A:47:PHE:HB2	1:A:63:ILE:HD11	1.88	0.56
1:A:55:THR:HG22	1:A:57:GLN:H	1.71	0.55
1:A:206:LYS:NZ	1:A:207:SER:HB2	2.20	0.55
1:A:172:ASN:O	1:A:176:VAL:HG23	2.07	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:ALA:HB1	1:A:209:SER:HG	1.71	0.54
2:B:63:ARG:C	2:B:64:ASN:OD1	2.46	0.53
1:A:52:LYS:HB3	1:A:52:LYS:HZ2	1.74	0.53
1:A:96:ASP:HB3	4:A:1069:HOH:O	2.08	0.52
1:A:40:GLN:HE21	1:A:40:GLN:CA	2.08	0.52
2:B:61:GLU:O	2:B:63:ARG:N	2.41	0.52
1:A:210:THR:HB	1:A:212:LYS:HZ3	1.75	0.52
1:A:210:THR:OG1	1:A:212:LYS:HE2	2.09	0.52
1:A:176:VAL:HG11	1:A:215:PHE:CD1	2.45	0.52
1:A:40:GLN:NE2	1:A:40:GLN:HA	2.13	0.51
1:A:49:LYS:O	1:A:59:ASN:HB3	2.11	0.51
1:A:131:LEU:O	1:A:135:LEU:HG	2.10	0.50
2:B:61:GLU:C	2:B:63:ARG:N	2.53	0.50
1:A:91:TYR:HB2	1:A:154:VAL:HG12	1.94	0.50
1:A:120:GLN:H	1:A:120:GLN:CD	2.13	0.50
1:A:34:TYR:CE1	1:A:35:ILE:HG22	2.46	0.50
1:A:170:CYS:SG	1:A:199:SER:HB2	2.52	0.49
1:A:31:PRO:HA	1:A:35:ILE:HG23	1.94	0.49
1:A:132:GLU:CD	2:B:56:ARG:HH21	2.16	0.49
1:A:91:TYR:HB2	1:A:154:VAL:CG1	2.43	0.49
1:A:32:GLU:O	1:A:32:GLU:OE2	2.32	0.48
1:A:30:ASN:C	1:A:30:ASN:ND2	2.66	0.48
1:A:132:GLU:OE1	2:B:56:ARG:NH2	2.45	0.48
1:A:206:LYS:HB3	1:A:209:SER:HB3	1.95	0.48
1:A:206:LYS:HD3	1:A:207:SER:CB	2.44	0.47
1:A:206:LYS:HD3	1:A:207:SER:HB2	1.96	0.47
1:A:207:SER:O	1:A:209:SER:N	2.47	0.47
1:A:51:ASP:OD1	1:A:53:SER:HB2	2.15	0.47
1:A:38:PRO:O	2:B:52:ILE:HG21	2.15	0.47
1:A:206:LYS:CD	1:A:207:SER:N	2.77	0.46
2:B:63:ARG:O	2:B:64:ASN:HB2	2.16	0.46
1:A:40:GLN:HB2	1:A:140:GLU:HG3	1.97	0.45
1:A:142:PHE:HB2	1:A:146:SER:OG	2.17	0.45
1:A:36:LYS:O	1:A:38:PRO:HD3	2.17	0.44
1:A:73:TRP:HB3	2:B:64:ASN:HD22	1.81	0.43
1:A:204:ALA:CB	1:A:209:SER:HG	2.31	0.43
2:B:56:ARG:HH11	2:B:56:ARG:CG	2.32	0.43
1:A:208:GLY:C	1:A:210:THR:H	2.21	0.43
2:B:55:ASP:OD2	2:B:57:LYS:HG2	2.18	0.42
1:A:141:SER:OG	1:A:186:ARG:NH2	2.46	0.42
1:A:108:LYS:HD2	1:A:109:ARG:HE	1.83	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:63:ARG:O	2:B:64:ASN:CB	2.67	0.42
1:A:35:ILE:HD12	4:A:1027:HOH:O	2.20	0.42
1:A:206:LYS:HZ2	1:A:207:SER:HB2	1.83	0.42
1:A:130:TRP:CE3	1:A:154:VAL:HB	2.54	0.42
1:A:30:ASN:HD22	1:A:31:PRO:CD	2.32	0.42
1:A:66:PHE:HA	4:A:1046:HOH:O	2.20	0.42
1:A:28:VAL:HG22	4:A:1079:HOH:O	2.20	0.41
1:A:47:PHE:HE2	1:A:49:LYS:HB2	1.86	0.41
1:A:206:LYS:CD	1:A:207:SER:HB2	2.51	0.41
1:A:77:ASN:ND2	2:B:65:SER:OG	2.53	0.41
1:A:61:ARG:HA	1:A:61:ARG:HD3	1.87	0.41
1:A:125:ASP:O	1:A:128:ARG:HG2	2.21	0.41
2:B:63:ARG:NH1	2:B:63:ARG:HB3	2.34	0.40
1:A:55:THR:HG22	1:A:57:GLN:N	2.35	0.40
1:A:108:LYS:CE	4:A:1062:HOH:O	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/191 (99%)	169 (89%)	17 (9%)	3 (2%)	12	6
2	B	18/20 (90%)	9 (50%)	6 (33%)	3 (17%)	0	0
All	All	207/211 (98%)	178 (86%)	23 (11%)	6 (3%)	6	2

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	62	CYS
1	A	52	LYS
1	A	209	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	210	THR
2	B	50	THR
2	B	65	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/170 (100%)	160 (94%)	10 (6%)	24	20
2	B	18/18 (100%)	15 (83%)	3 (17%)	3	1
All	All	188/188 (100%)	175 (93%)	13 (7%)	19	15

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	35	ILE
1	A	40	GLN
1	A	51	ASP
1	A	82	SER
1	A	91	TYR
1	A	148	ASP
1	A	153	VAL
1	A	198	GLN
1	A	205	THR
2	B	52	ILE
2	B	55	ASP
2	B	63	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	40	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	57	GLN
1	A	77	ASN
1	A	84	ASN
1	A	172	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GTA	A	1000	-	42,56,56	3.64	18 (42%)	50,88,88	3.83	17 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTA	A	1000	-	-	0/24/64/64	0/6/6/6

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1000	GTA	C5B-C4B	-5.23	1.34	1.51
3	A	1000	GTA	P3-O32	-2.38	1.44	1.54
3	A	1000	GTA	O2B-C2B	2.52	1.49	1.43
3	A	1000	GTA	C5C-N7C	2.61	1.48	1.39
3	A	1000	GTA	C2-N2	3.40	1.41	1.34
3	A	1000	GTA	C8-N7	3.90	1.40	1.33
3	A	1000	GTA	C2C-N3C	4.07	1.39	1.32
3	A	1000	GTA	C5C-C4C	4.21	1.50	1.40
3	A	1000	GTA	O4A-C1A	4.21	1.46	1.41
3	A	1000	GTA	O33-C5B	4.39	1.62	1.44
3	A	1000	GTA	C6-N1	4.65	1.41	1.33
3	A	1000	GTA	O4B-C1B	4.73	1.47	1.41
3	A	1000	GTA	C2C-N1C	5.81	1.44	1.33
3	A	1000	GTA	C8-N9	7.00	1.46	1.33
3	A	1000	GTA	C2-N1	7.64	1.48	1.35
3	A	1000	GTA	C6-C5	7.93	1.57	1.41
3	A	1000	GTA	C4-N3	8.01	1.48	1.35
3	A	1000	GTA	C4C-N3C	8.44	1.48	1.35

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1000	GTA	N3C-C2C-N1C	-13.15	118.82	128.89
3	A	1000	GTA	C4C-C5C-N7C	-11.21	99.17	109.48
3	A	1000	GTA	C1B-N9C-C4C	-6.55	117.05	126.94
3	A	1000	GTA	N3-C2-N1	-5.96	118.37	127.44
3	A	1000	GTA	C5-C6-N1	-5.10	116.61	123.59
3	A	1000	GTA	O33-P3-O31	-4.60	91.76	109.62
3	A	1000	GTA	P2-O13-P1	-3.72	122.29	132.73
3	A	1000	GTA	C6-C5-C4	-3.06	117.24	120.90
3	A	1000	GTA	O4B-C1B-N9C	-2.48	102.91	108.10
3	A	1000	GTA	O15-C5A-C4A	-2.28	100.70	109.12
3	A	1000	GTA	O4B-C4B-C3B	-2.26	100.60	105.15
3	A	1000	GTA	N2-C2-N1	2.63	121.55	117.20
3	A	1000	GTA	C5A-C4A-C3A	3.40	128.72	115.21
3	A	1000	GTA	O12-P1-O13	4.45	125.27	105.09
3	A	1000	GTA	C2C-N1C-C6C	4.55	126.90	118.77
3	A	1000	GTA	C6-N1-C2	5.96	124.22	115.94
3	A	1000	GTA	O23-P3-O33	11.70	133.98	102.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/191 (100%)	-0.03	4 (2%) 67 72	7, 14, 33, 40	0
2	B	20/20 (100%)	1.57	6 (30%) 1 1	15, 29, 47, 47	0
All	All	211/211 (100%)	0.12	10 (4%) 35 44	7, 14, 38, 47	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	66	PRO	5.9
2	B	47	PRO	5.5
1	A	208	GLY	3.5
2	B	64	ASN	3.2
2	B	48	GLY	2.9
1	A	210	THR	2.9
1	A	209	SER	2.6
2	B	49	GLY	2.4
2	B	50	THR	2.3
1	A	205	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GTA	A	1000	51/51	0.96	0.12	-0.45	1,24,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.