



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 13, 2016 – 05:39 PM EDT

PDB ID : 4WMK
Title : Crystal structure of mouse Xyloside xylosyltransferase 1 complexed with manganese, product ligand and UDP (Product complex II)
Authors : Yu, H.; Li, H.
Deposited on : 2014-10-09
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4546 (2.10-2.06)
Clashscore	102246	5101 (2.10-2.06)
Ramachandran outliers	100387	5048 (2.10-2.06)
Sidechain outliers	100360	5049 (2.10-2.06)
RSRZ outliers	91569	4556 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div> <div>2%</div> <div>80%</div> <div>16%</div> <div>• •</div> </div>
2	D	50	<div> <div>20%</div> <div>62%</div> <div>6%</div> <div>•</div> <div>30%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2849 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xyloside xylosyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	0	0
			2440	1582	417	426	15			

- Molecule 2 is a protein called Coagulation factor IX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	35	Total	C	N	O	S	0	0	0
			263	160	42	55	6			


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	43	MET	-	initiating methionine	UNP P00740
D	44	ASP	-	expression tag	UNP P00740
D	45	ILE	-	expression tag	UNP P00740
D	85	LEU	-	expression tag	UNP P00740
D	86	GLU	-	expression tag	UNP P00740
D	87	HIS	-	expression tag	UNP P00740
D	88	HIS	-	expression tag	UNP P00740
D	89	HIS	-	expression tag	UNP P00740
D	90	HIS	-	expression tag	UNP P00740
D	91	HIS	-	expression tag	UNP P00740
D	92	HIS	-	expression tag	UNP P00740

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 5 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O S	0	0
			5	4 1		

- Molecule 6 is alpha-D-xylopyranosyl-(1->3)-alpha-D-xylopyranosyl-(1->3)-beta-D-glucopyranose (three-letter code: GXX) (formula: C₁₆H₂₈O₁₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	C	O	0	0
			29	16	13		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	78	Total	O	0	0
			78	78		
7	D	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Xyloside xylosyltransferase 1



• Molecule 2: Coagulation factor IX



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	88.91Å 88.91Å 42.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.08 30.79 – 2.08	Depositor EDS
% Data completeness (in resolution range)	97.7 (40.00-2.08) 97.7 (30.79-2.08)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.191 , 0.246 0.191 , 0.243	Depositor DCC
R_{free} test set	1128 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	29.5	Xtriage
Anisotropy	0.318	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.1	EDS
Estimated twinning fraction	0.022 for -h,-k,l 0.034 for h,-h-k,-l 0.017 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 22151 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2849	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, MN, GXX, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	1/2516 (0.0%)	0.64	0/3414
2	D	0.61	1/269 (0.4%)	0.61	0/362
All	All	0.59	2/2785 (0.1%)	0.64	0/3776

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	265	TRP	CD2-CE2	5.46	1.47	1.41
2	D	72	TRP	CD2-CE2	5.20	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2440	0	2375	43	0
2	D	263	0	217	1	0
3	A	1	0	0	0	0
4	A	25	0	11	2	0
5	A	5	0	0	0	0
6	D	29	0	26	1	0
7	A	78	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	D	8	0	0	0	0
All	All	2849	0	2629	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:LEU:HD23	1:A:299:MET:HE1	1.51	0.91
1:A:99:HIS:HD2	1:A:139:HIS:HE1	1.21	0.86
1:A:296:LEU:HD23	1:A:299:MET:CE	2.11	0.81
1:A:139:HIS:HD2	1:A:171:HIS:NE2	1.87	0.73
1:A:294:LEU:HB3	1:A:299:MET:HE1	1.71	0.72
1:A:288:ASN:HD22	1:A:352:ASN:HD21	1.38	0.71
1:A:384:ASN:H	1:A:384:ASN:HD22	1.39	0.68
1:A:99:HIS:CD2	1:A:139:HIS:HE1	2.08	0.67
1:A:296:LEU:HA	1:A:299:MET:HE3	1.77	0.66
1:A:134:GLU:OE2	1:A:235:ARG:NH1	2.28	0.66
1:A:256:MET:H	1:A:354:GLN:NE2	1.97	0.62
1:A:256:MET:H	1:A:354:GLN:HE22	1.46	0.62
1:A:99:HIS:HD2	1:A:139:HIS:CE1	2.13	0.57
1:A:294:LEU:HB3	1:A:299:MET:CE	2.36	0.56
1:A:118:ARG:HH22	1:A:156:GLU:CD	2.10	0.54
1:A:384:ASN:HD22	1:A:384:ASN:N	2.01	0.54
4:A:402:UDP:O2B	6:D:101:GXX:H21	2.08	0.53
1:A:317:LEU:CD2	1:A:339:GLU:HG3	2.39	0.52
1:A:192:SER:O	1:A:194:GLY:HA2	2.11	0.50
1:A:236:GLU:O	1:A:239:GLU:HG2	2.12	0.49
1:A:217:GLU:O	1:A:219:PRO:HD3	2.11	0.49
1:A:369:GLN:HB2	7:A:510:HOH:O	2.12	0.49
1:A:288:ASN:ND2	1:A:352:ASN:HD21	2.07	0.49
1:A:288:ASN:ND2	1:A:290:GLY:H	2.11	0.48
1:A:295:ASN:C	1:A:299:MET:HE3	2.33	0.48
1:A:254:ARG:HG2	7:A:540:HOH:O	2.13	0.48
1:A:296:LEU:CA	1:A:299:MET:HE3	2.43	0.48
1:A:288:ASN:HD21	1:A:382:HIS:CE1	2.32	0.47
1:A:211:HIS:HB2	1:A:300:ARG:HG3	1.97	0.47
1:A:317:LEU:HD22	1:A:339:GLU:HG3	1.97	0.46
1:A:373:ARG:NH1	1:A:375:GLU:OE2	2.48	0.46
1:A:133:HIS:CD2	1:A:133:HIS:H	2.33	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:LEU:HD13	1:A:154:LEU:CD1	2.45	0.46
1:A:317:LEU:HD22	1:A:339:GLU:CG	2.46	0.46
2:D:52:GLU:HG2	2:D:63:LYS:HD3	1.97	0.46
1:A:257:GLN:NE2	1:A:359:TRP:H	2.15	0.45
1:A:295:ASN:N	1:A:299:MET:HE2	2.32	0.44
1:A:97:ASP:OD1	1:A:135:VAL:HG13	2.18	0.44
4:A:402:UDP:H3'	4:A:402:UDP:PA	2.57	0.43
1:A:136:LEU:HD21	1:A:138:LEU:HD21	2.01	0.43
1:A:183:VAL:HG21	1:A:212:GLN:NE2	2.35	0.42
1:A:184:VAL:HG12	1:A:188:GLN:HE21	1.84	0.41
1:A:94:VAL:CG2	1:A:94:VAL:O	2.69	0.41
1:A:134:GLU:OE1	1:A:235:ARG:NH1	2.54	0.41
1:A:256:MET:CE	1:A:349:CYS:HB2	2.51	0.41
1:A:134:GLU:CD	1:A:235:ARG:NH1	2.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/306 (97%)	287 (97%)	10 (3%)	0	100	100
2	D	33/50 (66%)	33 (100%)	0	0	100	100
All	All	330/356 (93%)	320 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/268 (98%)	249 (95%)	13 (5%)	30	26
2	D	30/45 (67%)	28 (93%)	2 (7%)	20	16
All	All	292/313 (93%)	277 (95%)	15 (5%)	29	25

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	135	VAL
1	A	154	LEU
1	A	184	VAL
1	A	189	LYS
1	A	208	VAL
1	A	282	GLU
1	A	309	LEU
1	A	324	ARG
1	A	353	ARG
1	A	369	GLN
1	A	375	GLU
1	A	384	ASN
1	A	391	GLU
2	D	63	LYS
2	D	78	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	HIS
1	A	139	HIS
1	A	188	GLN
1	A	212	GLN
1	A	257	GLN
1	A	288	ASN
1	A	354	GLN
1	A	384	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	UDP	A	402	3	20,26,26	0.81	0	24,40,40	1.56	2 (8%)
5	SO4	A	403	-	4,4,4	0.51	0	6,6,6	0.24	0
6	GXX	D	101	2	31,31,32	0.75	0	41,45,47	1.35	4 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UDP	A	402	3	-	0/12/32/32	0/2/2/2
5	SO4	A	403	-	-	0/0/0/0	0/0/0/0
6	GXX	D	101	2	-	0/10/61/64	0/3/3/3

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	101	GXX	O22-C22-C12	-2.45	104.57	110.01
4	A	402	UDP	O4'-C1'-N1	2.02	111.94	108.10
6	D	101	GXX	C1-C2-C3	2.13	114.20	109.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	101	GXX	C5-C4-C3	3.07	113.28	109.55
6	D	101	GXX	C52-C42-C32	3.11	113.06	109.23
4	A	402	UDP	C4-N3-C2	5.80	120.32	114.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	402	UDP	2	0
6	D	101	GXX	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/306 (97%)	-0.02	6 (2%) 68 73	18, 29, 47, 74	1 (0%)
2	D	35/50 (70%)	1.06	10 (28%) 1 1	26, 39, 75, 92	0
All	All	334/356 (93%)	0.09	16 (4%) 34 42	18, 29, 53, 92	1 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	84	LEU	9.2
1	A	93	VAL	7.3
1	A	94	VAL	6.8
2	D	69	TYR	4.0
2	D	81	ASN	3.7
2	D	83	GLU	3.7
2	D	75	PHE	3.6
2	D	80	LYS	3.6
2	D	79	GLY	3.3
1	A	161	ALA	3.2
2	D	77	PHE	2.6
2	D	78	GLU	2.5
1	A	291	VAL	2.4
2	D	76	GLY	2.4
1	A	95	PRO	2.3
1	A	109	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	GXX	D	101	29/30	0.94	0.13	-0.16	25,28,45,46	0
5	SO4	A	403	5/5	0.99	0.11	-0.66	36,38,41,42	0
4	UDP	A	402	25/25	0.94	0.10	-0.72	30,36,40,44	0
3	MN	A	401	1/1	0.99	0.10	-	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.