



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:14 PM GMT

PDB ID : 1WNV  
Title : D136A mutant of Heme Oxygenase from *Corynebacterium diphtheriae* (HmuO)  
Authors : Matsui, T.; Unno, M.; Ikeda-Saito, M.  
Deposited on : 2004-08-10  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	<div> <div>5%</div> <div>85%</div> <div>11%</div> <div>.</div> </div>
1	B	215	<div> <div>%</div> <div>87%</div> <div>9%</div> <div>..</div> </div>
1	C	215	<div> <div>%</div> <div>87%</div> <div>8%</div> <div>..</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5426 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heme oxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	0	0
			1649	1041	293	312	3			
1	B	210	Total	C	N	O	S	0	0	0
			1666	1052	296	315	3			
1	C	207	Total	C	N	O	S	0	0	0
			1649	1041	293	312	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	LYS	GLU	SEE REMARK 999	UNP P71119
A	60	VAL	ALA	SEE REMARK 999	UNP P71119
A	92	GLY	ASP	SEE REMARK 999	UNP P71119
A	93	SER	GLY	SEE REMARK 999	UNP P71119
A	136	ALA	ASP	ENGINEERED	UNP P71119
A	192	HIS	ASN	SEE REMARK 999	UNP P71119
B	334	LYS	GLU	SEE REMARK 999	UNP P71119
B	360	VAL	ALA	SEE REMARK 999	UNP P71119
B	392	GLY	ASP	SEE REMARK 999	UNP P71119
B	393	SER	GLY	SEE REMARK 999	UNP P71119
B	436	ALA	ASP	ENGINEERED	UNP P71119
B	492	HIS	ASN	SEE REMARK 999	UNP P71119
C	634	LYS	GLU	SEE REMARK 999	UNP P71119
C	660	VAL	ALA	SEE REMARK 999	UNP P71119
C	692	GLY	ASP	SEE REMARK 999	UNP P71119
C	693	SER	GLY	SEE REMARK 999	UNP P71119
C	736	ALA	ASP	ENGINEERED	UNP P71119
C	792	HIS	ASN	SEE REMARK 999	UNP P71119

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		
4	B	119	Total	O	0	0
			119	119		
4	C	124	Total	O	0	0
			124	124		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.13 Å   62.78 Å   107.99 Å 90.00°   101.00°   90.00°	Depositor
Resolution (Å)	20.00 – 1.85 40.50 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.1 (20.00-1.85) 98.1 (40.50-1.85)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.15 (at 1.86 Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.164   ,   0.204 0.177   ,   0.215	Depositor DCC
$R_{free}$ test set	6049 reflections (11.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.6	Xtriage
Anisotropy	0.295	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 46.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 59721 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5426	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.91	1/1681 (0.1%)	1.00	10/2270 (0.4%)
1	B	1.10	3/1698 (0.2%)	0.95	4/2293 (0.2%)
1	C	1.10	4/1681 (0.2%)	1.02	6/2270 (0.3%)
All	All	1.04	8/5060 (0.2%)	0.99	20/6833 (0.3%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	381	GLU	CD-OE2	8.64	1.35	1.25
1	B	409	TYR	CD1-CE1	7.80	1.51	1.39
1	C	681	GLU	CG-CD	6.29	1.61	1.51
1	C	709	TYR	CE1-CZ	-6.04	1.30	1.38
1	C	681	GLU	CD-OE1	5.97	1.32	1.25
1	C	681	GLU	CD-OE2	5.29	1.31	1.25
1	B	381	GLU	CG-CD	5.21	1.59	1.51
1	A	7	GLY	N-CA	5.02	1.53	1.46

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	154	ASP	CB-CG-OD2	11.69	128.82	118.30
1	C	699	ARG	NE-CZ-NH2	-11.05	114.78	120.30
1	C	699	ARG	NE-CZ-NH1	8.26	124.43	120.30
1	B	454	ASP	CB-CG-OD2	7.87	125.38	118.30
1	B	408	ASP	CB-CG-OD2	7.40	124.96	118.30
1	A	31	ASP	CB-CG-OD2	7.15	124.74	118.30
1	C	631	ASP	CB-CG-OD2	6.87	124.48	118.30
1	A	210	ASP	CB-CG-OD2	6.70	124.33	118.30
1	A	112	ARG	NE-CZ-NH2	6.37	123.48	120.30
1	C	718	ASP	CB-CG-OD1	6.07	123.77	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	112	ARG	NE-CZ-NH1	-5.90	117.35	120.30
1	A	29	MET	CG-SD-CE	-5.86	90.83	100.20
1	C	708	ASP	CB-CG-OD1	-5.54	113.31	118.30
1	A	44	ARG	NE-CZ-NH1	5.45	123.03	120.30
1	A	174	ASP	CB-CG-OD2	5.38	123.14	118.30
1	A	187	ASP	CB-CG-OD2	5.36	123.13	118.30
1	B	399	ARG	NE-CZ-NH1	5.23	122.91	120.30
1	C	679	ARG	NE-CZ-NH1	5.21	122.91	120.30
1	B	386	ASP	CB-CG-OD2	5.18	122.96	118.30
1	A	86	ASP	CB-CG-OD2	5.17	122.95	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1649	0	1612	6	0
1	B	1666	0	1631	8	0
1	C	1649	0	1612	13	0
2	A	5	0	0	0	0
2	B	10	0	0	1	0
2	C	10	0	0	0	0
3	A	43	0	30	1	0
3	B	43	0	30	1	0
3	C	43	0	30	2	0
4	A	65	0	0	4	0
4	B	119	0	0	0	0
4	C	124	0	0	2	0
All	All	5426	0	4945	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:634:LYS:HG3	4:C:44:HOH:O	1.68	0.92
1:C:642:PHE:HB2	1:C:747:MET:HE1	1.81	0.63
1:A:204:ASN:ND2	1:A:208:PHE:CE2	2.68	0.62
1:B:473:LYS:NZ	2:B:2011:SO4:O2	2.33	0.60
1:A:208:PHE:CD2	4:A:2056:HOH:O	2.53	0.58
1:A:82:VAL:HG23	4:A:2022:HOH:O	2.02	0.58
1:C:646:GLN:HE21	1:C:646:GLN:HA	1.72	0.55
1:C:642:PHE:CG	1:C:747:MET:HE2	2.43	0.53
3:C:903:HEM:HHC	3:C:903:HEM:HBB2	1.89	0.53
1:C:646:GLN:NE2	1:C:646:GLN:HA	2.24	0.52
3:A:901:HEM:HBD1	4:A:2072:HOH:O	2.10	0.51
1:C:741:GLN:HG3	4:C:77:HOH:O	2.11	0.51
1:C:704:PRO:HD2	1:C:810:ASP:OD2	2.11	0.50
1:B:442:VAL:HG13	1:B:446:MET:CE	2.42	0.49
1:A:137:LEU:HD22	1:A:169:LEU:HD11	1.94	0.49
1:C:738:SER:HB3	3:C:903:HEM:HBD1	1.95	0.48
1:B:432:ARG:NH1	1:B:504:ASN:OD1	2.46	0.48
3:B:902:HEM:HHD	3:B:902:HEM:HBC2	1.96	0.47
1:B:510:ASP:HA	1:B:513:LYS:HE3	1.97	0.47
1:B:454:ASP:OD2	1:B:455:PRO:HD2	2.15	0.46
1:A:13:LYS:NZ	4:A:2072:HOH:O	2.46	0.46
1:B:322:LYS:HG2	1:B:505:HIS:CD2	2.52	0.45
1:C:747:MET:HE3	1:C:751:TYR:HD1	1.81	0.45
1:B:442:VAL:HG13	1:B:446:MET:HE1	1.98	0.45
1:C:642:PHE:HB2	1:C:747:MET:CE	2.46	0.45
1:B:474:ASP:O	1:B:478:GLU:HG3	2.17	0.43
1:C:689:LYS:HB3	1:C:689:LYS:HE2	1.68	0.42
1:C:695:GLU:OE2	1:C:699:ARG:HD3	2.21	0.41
1:C:700:ILE:HD13	1:C:700:ILE:HG21	1.75	0.41
1:A:203:PHE:O	1:A:207:VAL:HG23	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/215 (95%)	200 (98%)	5 (2%)	0	100	100
1	B	208/215 (97%)	205 (99%)	3 (1%)	0	100	100
1	C	205/215 (95%)	203 (99%)	2 (1%)	0	100	100
All	All	618/645 (96%)	608 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/174 (97%)	162 (96%)	7 (4%)	37	17
1	B	170/174 (98%)	164 (96%)	6 (4%)	43	23
1	C	169/174 (97%)	164 (97%)	5 (3%)	48	29
All	All	508/522 (97%)	490 (96%)	18 (4%)	43	23

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	141	GLN
1	A	162	HIS
1	A	168	LYS
1	A	170	LYS
1	A	204	ASN
1	A	213	LYS
1	B	321	GLU
1	B	346	GLN
1	B	432	ARG
1	B	441	GLN
1	B	442	VAL
1	B	445	ARG
1	C	614	GLN
1	C	646	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	689	LYS
1	C	708	ASP
1	C	783	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	A	182	ASN
1	B	346	GLN
1	B	378	ASN
1	B	448	GLN
1	B	505	HIS
1	B	506	GLN
1	C	646	GLN
1	C	678	ASN
1	C	741	GLN
1	C	748	GLN
1	C	762	HIS
1	C	806	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	2013	-	4,4,4	0.45	0	6,6,6	0.30	0
3	HEM	A	901	1,4	30,50,50	2.11	5 (16%)	24,82,82	2.80	11 (45%)
2	SO4	B	2011	-	4,4,4	0.51	0	6,6,6	0.60	0
2	SO4	B	2012	-	4,4,4	0.81	0	6,6,6	0.47	0
3	HEM	B	902	1,4	30,50,50	2.33	8 (26%)	24,82,82	2.58	11 (45%)
2	SO4	C	2014	-	4,4,4	0.55	0	6,6,6	0.65	0
2	SO4	C	2015	-	4,4,4	0.35	0	6,6,6	0.59	0
3	HEM	C	903	1,4	30,50,50	2.53	6 (20%)	24,82,82	2.33	10 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	2013	-	-	0/0/0/0	0/0/0/0
3	HEM	A	901	1,4	-	0/10/54/54	0/0/8/8
2	SO4	B	2011	-	-	0/0/0/0	0/0/0/0
2	SO4	B	2012	-	-	0/0/0/0	0/0/0/0
3	HEM	B	902	1,4	-	0/10/54/54	0/0/8/8
2	SO4	C	2014	-	-	0/0/0/0	0/0/0/0
2	SO4	C	2015	-	-	0/0/0/0	0/0/0/0
3	HEM	C	903	1,4	-	0/10/54/54	0/0/8/8

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	903	HEM	C3B-C4B	-9.63	1.43	1.51
3	B	902	HEM	C3B-C4B	-8.99	1.43	1.51
3	A	901	HEM	C3B-C4B	-8.22	1.44	1.51
3	C	903	HEM	C3D-C4D	-4.37	1.46	1.51
3	A	901	HEM	C3D-C4D	-4.15	1.46	1.51
3	B	902	HEM	C3D-C4D	-4.07	1.46	1.51
3	A	901	HEM	C2C-C1C	-3.82	1.45	1.52
3	B	902	HEM	C2C-C1C	-3.30	1.46	1.52
3	B	902	HEM	C2D-C1D	-2.70	1.43	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	902	HEM	C2B-C1B	-2.61	1.43	1.51
3	C	903	HEM	C2B-C1B	-2.25	1.44	1.51
3	A	901	HEM	C2B-C1B	-2.21	1.44	1.51
3	A	901	HEM	C2D-C1D	-2.20	1.44	1.51
3	C	903	HEM	C2D-C1D	-2.19	1.44	1.51
3	B	902	HEM	C4C-NC	2.11	1.38	1.36
3	B	902	HEM	C3C-CAC	2.24	1.55	1.51
3	B	902	HEM	FE-NC	3.34	2.09	1.95
3	C	903	HEM	C1C-NC	3.86	1.40	1.36
3	C	903	HEM	C4C-NC	4.89	1.42	1.36

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	HEM	C3B-CAB-CBB	-5.69	115.73	124.46
3	B	902	HEM	CAA-CBA-CGA	-3.49	106.36	112.75
3	A	901	HEM	C3C-CAC-CBC	-3.48	119.12	124.46
3	A	901	HEM	CAA-C2A-C1A	-3.45	123.26	127.01
3	C	903	HEM	CAA-CBA-CGA	-3.38	106.56	112.75
3	B	902	HEM	C3B-CAB-CBB	-3.22	119.52	124.46
3	B	902	HEM	C3C-CAC-CBC	-3.09	119.72	124.46
3	A	901	HEM	CAA-CBA-CGA	-2.98	107.29	112.75
3	C	903	HEM	C3C-CAC-CBC	-2.60	120.47	124.46
3	B	902	HEM	C3B-C4B-NB	-2.42	107.00	111.63
3	C	903	HEM	C3B-C4B-NB	-2.39	107.06	111.63
3	A	901	HEM	CMA-C3A-C4A	-2.37	124.44	128.36
3	A	901	HEM	CBA-CAA-C2A	-2.36	108.30	112.53
3	C	903	HEM	CMA-C3A-C4A	-2.26	124.63	128.36
3	B	902	HEM	CBD-CAD-C3D	-2.06	107.55	113.55
3	B	902	HEM	C2C-C1C-CHC	2.05	126.81	123.68
3	C	903	HEM	CMD-C2D-C3D	2.19	124.01	114.35
3	A	901	HEM	CMD-C2D-C3D	2.61	125.88	114.35
3	C	903	HEM	CMB-C2B-C3B	2.88	123.72	116.53
3	C	903	HEM	C3B-C4B-CHC	3.17	127.63	123.16
3	B	902	HEM	CMD-C2D-C3D	3.19	128.48	114.35
3	A	901	HEM	CAD-C3D-C2D	3.71	123.89	113.22
3	A	901	HEM	CMB-C2B-C3B	3.77	125.94	116.53
3	B	902	HEM	CMB-C2B-C3B	3.80	126.02	116.53
3	B	902	HEM	CAD-C3D-C4D	4.07	126.81	112.47
3	C	903	HEM	CMC-C2C-C3C	4.35	127.40	116.53
3	C	903	HEM	CAD-C3D-C2D	4.77	126.94	113.22
3	C	903	HEM	CAD-C3D-C4D	5.00	130.09	112.47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	902	HEM	CMC-C2C-C3C	5.23	129.59	116.53
3	B	902	HEM	CAD-C3D-C2D	5.35	128.61	113.22
3	A	901	HEM	CMC-C2C-C3C	5.43	130.08	116.53
3	A	901	HEM	CAD-C3D-C4D	5.49	131.82	112.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	HEM	1	0
2	B	2011	SO4	1	0
3	B	902	HEM	1	0
3	C	903	HEM	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/215 (96%)	0.14	10 (4%) 34 32	25, 37, 55, 60	0
1	B	210/215 (97%)	-0.06	3 (1%) 78 78	20, 29, 48, 58	0
1	C	207/215 (96%)	-0.20	3 (1%) 78 78	20, 30, 46, 58	0
All	All	624/645 (96%)	-0.04	16 (2%) 59 57	20, 32, 52, 60	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	93	SER	4.0
1	B	306	ALA	3.6
1	A	213	LYS	2.8
1	A	187	ASP	2.8
1	A	95	GLU	2.7
1	A	98	SER	2.6
1	B	325	HIS	2.6
1	A	101	THR	2.4
1	C	765	GLY	2.3
1	C	618	GLN	2.2
1	C	636	ARG	2.2
1	A	27	THR	2.0
1	A	92	GLY	2.0
1	B	505	HIS	2.0
1	A	156	GLU	2.0
1	A	137	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	2013	5/5	0.91	0.22	1.59	44,44,45,46	5
3	HEM	A	901	43/43	0.96	0.12	0.68	30,43,86,88	0
3	HEM	B	902	43/43	0.96	0.12	0.24	27,36,57,69	0
3	HEM	C	903	43/43	0.96	0.10	-0.08	23,34,49,52	0
2	SO4	B	2011	5/5	0.97	0.08	-	32,32,36,37	5
2	SO4	C	2014	5/5	0.93	0.12	-	38,39,41,42	5
2	SO4	B	2012	5/5	0.98	0.10	-	25,26,30,32	5
2	SO4	C	2015	5/5	0.97	0.10	-	28,28,34,34	5

### 6.5 Other polymers [i](#)

There are no such residues in this entry.