



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:49 PM GMT

PDB ID : 4WP2
Title : Chaetomium Mex67 UBA domain
Authors : Aibara, S.; Stewart, M.
Deposited on : 2014-10-17
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	61	<div><div>7%</div><div>98%</div><div>.</div></div>
1	B	61	<div><div>28%</div><div>79%</div><div>16%</div><div>5%</div></div>
1	C	61	<div><div>11%</div><div>93%</div><div>7%</div></div>
1	D	61	<div><div>31%</div><div>82%</div><div>13%</div><div>5%</div></div>
1	E	61	<div><div>11%</div><div>97%</div><div>.</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	61	<div><div></div><div>28%</div><div></div><div>85%</div><div>7%</div><div>8%</div></div>
1	G	61	<div><div></div><div>10%</div><div></div><div>93%</div><div>7%</div><div></div></div>
1	H	61	<div><div></div><div>21%</div><div></div><div>89%</div><div>7%</div><div>5%</div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7657 atoms, of which 3623 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative mRNA export protein.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	B	58	Total	As	C	H	N	O	S	0	0	0
			899	1	282	447	78	88	3			
1	C	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	D	58	Total	As	C	H	N	O	S	0	0	0
			899	1	282	447	78	88	3			
1	E	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	F	56	Total	As	C	H	N	O	S	0	0	0
			865	1	272	430	75	84	3			
1	G	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	H	58	Total	As	C	H	N	O	S	0	0	0
			899	1	282	447	78	88	3			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	597	SER	-	expression tag	UNP G0SET4
A	598	ASN	-	expression tag	UNP G0SET4
A	599	ALA	-	expression tag	UNP G0SET4
B	597	SER	-	expression tag	UNP G0SET4
B	598	ASN	-	expression tag	UNP G0SET4
B	599	ALA	-	expression tag	UNP G0SET4
C	597	SER	-	expression tag	UNP G0SET4
C	598	ASN	-	expression tag	UNP G0SET4
C	599	ALA	-	expression tag	UNP G0SET4
D	597	SER	-	expression tag	UNP G0SET4
D	598	ASN	-	expression tag	UNP G0SET4
D	599	ALA	-	expression tag	UNP G0SET4
E	597	SER	-	expression tag	UNP G0SET4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	598	ASN	-	expression tag	UNP G0SET4
E	599	ALA	-	expression tag	UNP G0SET4
F	597	SER	-	expression tag	UNP G0SET4
F	598	ASN	-	expression tag	UNP G0SET4
F	599	ALA	-	expression tag	UNP G0SET4
G	597	SER	-	expression tag	UNP G0SET4
G	598	ASN	-	expression tag	UNP G0SET4
G	599	ALA	-	expression tag	UNP G0SET4
H	597	SER	-	expression tag	UNP G0SET4
H	598	ASN	-	expression tag	UNP G0SET4
H	599	ALA	-	expression tag	UNP G0SET4

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	46	Total O 46 46	0	0
2	B	41	Total O 41 41	0	0
2	C	38	Total O 38 38	0	0
2	D	56	Total O 56 56	0	0
2	E	49	Total O 49 49	0	0
2	F	47	Total O 47 47	0	0
2	G	40	Total O 40 40	0	0
2	H	42	Total O 42 42	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative mRNA export protein





- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein



- Molecule 1: Putative mRNA export protein



- Molecule 1: Putative mRNA export protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	95.90 Å 95.90 Å 74.97 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.53 – 1.70 47.95 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.53-1.70) 100.0 (47.95-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 1.70 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.198 , 0.240 0.245 , 0.277	Depositor DCC
R_{free} test set	1000 reflections (2.27%)	DCC
Wilson B-factor (Å ²)	36.2	Xtriage
Anisotropy	0.435	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.7	EDS
Estimated twinning fraction	0.047 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	25 of 44111 reflections (0.057%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7657	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 80.68 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.2251e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.29	0/468	0.46	0/632
1	B	0.31	0/449	0.49	0/606
1	C	0.29	0/468	0.47	0/632
1	D	0.32	0/449	0.50	0/606
1	E	0.32	0/468	0.47	0/632
1	F	0.29	0/432	0.47	0/584
1	G	0.30	0/468	0.47	0/632
1	H	0.32	0/449	0.50	0/606
All	All	0.31	0/3651	0.48	0/4930

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	471	463	457	1	0
1	B	452	447	441	8	0
1	C	471	463	457	3	1
1	D	452	447	441	7	0
1	E	471	463	457	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	435	430	424	3	0
1	G	471	463	457	2	1
1	H	452	447	441	2	0
2	A	46	0	0	1	1
2	B	41	0	0	2	0
2	C	38	0	0	1	0
2	D	56	0	0	1	0
2	E	49	0	0	0	0
2	F	47	0	0	0	0
2	G	40	0	0	0	1
2	H	42	0	0	1	0
All	All	4034	3623	3575	23	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:608:GLU:OE2	2:A:701:HOH:O	2.11	0.69
1:E:621:GLN:OE1	1:F:604:GLN:NE2	2.29	0.66
1:B:615:MET:CE	1:B:619:TYR:HB3	2.29	0.63
1:B:615:MET:HE3	1:B:619:TYR:HB3	1.81	0.62
1:H:600:ALA:N	2:H:702:HOH:O	2.33	0.62
1:D:623:CAS:CE2	1:D:638:SER:HB3	2.36	0.56
1:D:623:CAS:CE2	1:D:638:SER:CB	2.91	0.49
1:C:652:ILE:HD12	1:D:611:ARG:HD2	1.95	0.47
1:D:627:ALA:HB3	1:E:627:ALA:HB3	1.98	0.46
1:B:627:ALA:HB3	1:C:627:ALA:HB3	1.97	0.45
1:G:608:GLU:OE2	1:G:612:ARG:NH1	2.50	0.45
1:D:600:ALA:N	2:D:702:HOH:O	2.49	0.45
1:C:612:ARG:NH1	2:C:736:HOH:O	2.49	0.44
1:B:654:GLN:HG3	1:B:655:PRO:HD2	2.00	0.44
1:B:623:CAS:CE2	1:B:638:SER:HB3	2.48	0.43
1:F:654:GLN:HG3	1:F:655:PRO:HD2	2.01	0.43
1:B:633:GLU:HG3	2:B:715:HOH:O	2.19	0.43
1:F:627:ALA:HB3	1:G:627:ALA:HB3	2.00	0.43
1:D:657:VAL:HG23	1:D:657:VAL:O	2.18	0.43
1:D:654:GLN:HG3	1:D:655:PRO:HD2	2.01	0.42
1:B:615:MET:SD	1:B:639:PHE:CD1	3.12	0.42
1:H:623:CAS:CE2	1:H:638:SER:HB3	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:656:GLN:NE2	2:B:703:HOH:O	2.54	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:731:HOH:O	2:G:716:HOH:O[5_665]	2.09	0.11
1:C:643:LYS:HZ3	1:G:597:SER:O[4_455]	1.57	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	58/61 (95%)	57 (98%)	1 (2%)	0	100	100
1	B	55/61 (90%)	54 (98%)	1 (2%)	0	100	100
1	C	58/61 (95%)	57 (98%)	1 (2%)	0	100	100
1	D	55/61 (90%)	55 (100%)	0	0	100	100
1	E	58/61 (95%)	58 (100%)	0	0	100	100
1	F	53/61 (87%)	53 (100%)	0	0	100	100
1	G	58/61 (95%)	57 (98%)	1 (2%)	0	100	100
1	H	55/61 (90%)	55 (100%)	0	0	100	100
All	All	450/488 (92%)	446 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	49/49 (100%)	49 (100%)	0	100	100
1	B	47/49 (96%)	47 (100%)	0	100	100
1	C	49/49 (100%)	49 (100%)	0	100	100
1	D	47/49 (96%)	47 (100%)	0	100	100
1	E	49/49 (100%)	49 (100%)	0	100	100
1	F	45/49 (92%)	45 (100%)	0	100	100
1	G	49/49 (100%)	49 (100%)	0	100	100
1	H	47/49 (96%)	46 (98%)	1 (2%)	61	42
All	All	382/392 (97%)	381 (100%)	1 (0%)	94	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	637	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	604	GLN
1	B	645	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	623	1	4,8,9	1.41	0	2,9,11	1.49	1 (50%)
1	CAS	B	623	1	4,8,9	1.52	0	2,9,11	1.70	1 (50%)
1	CAS	C	623	1	4,8,9	1.24	0	2,9,11	1.63	1 (50%)
1	CAS	D	623	1	4,8,9	1.65	2 (50%)	2,9,11	1.38	0
1	CAS	E	623	1	4,8,9	1.26	0	2,9,11	1.67	1 (50%)
1	CAS	F	623	1	4,8,9	1.49	0	2,9,11	1.50	0
1	CAS	G	623	1	4,8,9	1.26	0	2,9,11	1.51	1 (50%)
1	CAS	H	623	1	4,8,9	1.51	0	2,9,11	1.56	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	623	1	-	0/0/7/9	0/0/0/0
1	CAS	B	623	1	-	0/0/7/9	0/0/0/0
1	CAS	C	623	1	-	0/0/7/9	0/0/0/0
1	CAS	D	623	1	-	0/0/7/9	0/0/0/0
1	CAS	E	623	1	-	0/0/7/9	0/0/0/0
1	CAS	F	623	1	-	0/0/7/9	0/0/0/0
1	CAS	G	623	1	-	0/0/7/9	0/0/0/0
1	CAS	H	623	1	-	0/0/7/9	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	623	CAS	AS-CE2	-2.29	1.89	1.96
1	D	623	CAS	O-C	2.00	1.29	1.19

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	623	CAS	O-C-CA	-2.36	119.38	125.72
1	C	623	CAS	O-C-CA	-2.29	119.58	125.72
1	H	623	CAS	O-C-CA	-2.18	119.87	125.72
1	B	623	CAS	O-C-CA	-2.14	119.99	125.72
1	A	623	CAS	O-C-CA	-2.05	120.21	125.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	G	623	CAS	O-C-CA	-2.04	120.26	125.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	623	CAS	1	0
1	D	623	CAS	2	0
1	H	623	CAS	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	60/61 (98%)	0.67	4 (6%) 21 23	20, 29, 54, 65	0
1	B	57/61 (93%)	1.77	17 (29%) 1 0	21, 39, 87, 100	0
1	C	60/61 (98%)	0.86	7 (11%) 6 7	19, 33, 71, 89	0
1	D	57/61 (93%)	2.39	19 (33%) 0 0	17, 30, 76, 92	0
1	E	60/61 (98%)	1.03	7 (11%) 6 7	18, 27, 47, 99	0
1	F	55/61 (90%)	1.83	17 (30%) 1 0	18, 35, 82, 108	0
1	G	60/61 (98%)	0.87	6 (10%) 9 9	18, 32, 68, 80	0
1	H	57/61 (93%)	1.27	13 (22%) 1 1	17, 31, 80, 104	0
All	All	466/488 (95%)	1.32	90 (19%) 2 1	17, 31, 74, 108	0

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	657	VAL	14.8
1	D	648	PRO	9.4
1	B	600	ALA	9.3
1	B	657	VAL	9.1
1	F	600	ALA	8.9
1	D	655	PRO	8.1
1	D	650	ALA	7.9
1	B	654	GLN	7.9
1	D	654	GLN	7.7
1	D	652	ILE	7.5
1	E	657	VAL	7.3
1	H	649	GLU	7.1
1	F	652	ILE	6.7
1	D	653	SER	6.6
1	F	649	GLU	6.4
1	E	644	ALA	6.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	657	VAL	6.1
1	H	648	PRO	6.0
1	F	654	GLN	6.0
1	H	656	GLN	5.9
1	D	644	ALA	5.6
1	F	617	VAL	5.4
1	D	645	ASN	5.3
1	D	643	LYS	5.1
1	H	652	ILE	5.0
1	H	654	GLN	5.0
1	D	656	GLN	4.7
1	B	651	PHE	4.7
1	D	619	TYR	4.7
1	G	657	VAL	4.6
1	H	655	PRO	4.5
1	B	652	ILE	4.4
1	C	657	VAL	4.3
1	B	649	GLU	4.3
1	F	636	LEU	4.2
1	F	650	ALA	4.2
1	B	650	ALA	4.2
1	B	645	ASN	4.1
1	G	597	SER	4.0
1	B	656	GLN	3.9
1	C	597	SER	3.8
1	B	646	VAL	3.8
1	B	648	PRO	3.7
1	C	645	ASN	3.7
1	E	656	GLN	3.6
1	F	648	PRO	3.5
1	D	640	GLU	3.4
1	F	647	PRO	3.3
1	F	655	PRO	3.3
1	A	657	VAL	3.2
1	B	655	PRO	3.0
1	A	644	ALA	3.0
1	F	646	VAL	3.0
1	B	647	PRO	3.0
1	C	643	LYS	3.0
1	A	645	ASN	3.0
1	C	644	ALA	3.0
1	D	647	PRO	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	649	GLU	2.9
1	B	639	PHE	2.8
1	F	645	ASN	2.8
1	D	646	VAL	2.8
1	H	650	ALA	2.8
1	F	639	PHE	2.7
1	G	633	GLU	2.7
1	H	653	SER	2.7
1	D	613	THR	2.7
1	H	643	LYS	2.6
1	F	610	SER	2.5
1	D	651	PHE	2.5
1	B	619	TYR	2.5
1	B	644	ALA	2.5
1	H	651	PHE	2.5
1	A	597	SER	2.4
1	H	645	ASN	2.4
1	G	644	ALA	2.3
1	C	646	VAL	2.2
1	F	653	SER	2.2
1	E	650	ALA	2.2
1	E	648	PRO	2.2
1	E	646	VAL	2.2
1	F	643	LYS	2.2
1	G	619	TYR	2.2
1	B	618	GLU	2.1
1	G	645	ASN	2.1
1	D	600	ALA	2.1
1	F	618	GLU	2.1
1	H	647	PRO	2.1
1	E	652	ILE	2.0
1	C	600	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CAS	H	623	9/10	0.94	0.11	-	19,28,67,67	0
1	CAS	A	623	9/10	0.96	0.11	-	19,23,47,47	0
1	CAS	C	623	9/10	0.97	0.10	-	20,27,43,43	0
1	CAS	B	623	9/10	0.93	0.13	-	21,32,52,52	0
1	CAS	E	623	9/10	0.86	0.14	-	18,23,51,51	0
1	CAS	D	623	9/10	0.85	0.20	-	17,24,83,83	0
1	CAS	G	623	9/10	0.95	0.12	-	18,27,69,69	0
1	CAS	F	623	9/10	0.89	0.15	-	20,30,68,68	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.