



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:16 PM GMT

PDB ID : 3WQ7
Title : New crystal form of the hyperthermophilic family 12 endo-cellulase from Pyrococcus furiosus
Authors : Kataoka, M.; Ishikawa, K.
Deposited on : 2014-01-23
Resolution : 1.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4802 (1.70-1.66)
Clashscore	102246	5317 (1.70-1.66)
Ramachandran outliers	100387	5225 (1.70-1.66)
Sidechain outliers	100360	5224 (1.70-1.66)
RSRZ outliers	91569	4813 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	<div> <div>2%</div> <div>72%</div> <div>12%</div> <div>•</div> <div>15%</div> </div>
1	B	319	<div> <div>10%</div> <div>56%</div> <div>26%</div> <div>•</div> <div>15%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	504	-	-	-	X
3	GOL	A	505	-	-	-	X
3	GOL	A	506	-	-	-	X
3	GOL	A	507	-	-	-	X
3	GOL	A	508	-	-	-	X
3	GOL	A	511	-	-	-	X
3	GOL	A	512	-	-	-	X
3	GOL	A	514[A]	-	-	-	X
3	GOL	A	514[B]	-	-	-	X
3	GOL	A	515	-	-	-	X
3	GOL	A	516	-	-	-	X
3	GOL	B	503	-	-	-	X
3	GOL	B	504	-	-	X	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4967 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Endoglucanase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	3	0
			2195	1427	351	415	2			
1	B	270	Total	C	N	O	S	0	3	0
			2195	1427	351	415	2			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 12 6 6	0	1
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	289	Total O 291 291	0	2
4	B	179	Total O 180 180	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endoglucanase A



• Molecule 1: Endoglucanase A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	134.73 Å 62.57 Å 86.28 Å 90.00° 95.08° 90.00°	Depositor
Resolution (Å)	43.01 – 1.68 42.97 – 1.68	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.01-1.68) 99.9 (42.97-1.68)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.68 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.181 , 0.217 0.180 , 0.216	Depositor DCC
R_{free} test set	4096 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	18.3	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 81565 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4967	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.42	12/2263 (0.5%)	1.32	8/3105 (0.3%)
1	B	1.26	5/2263 (0.2%)	1.18	3/3105 (0.1%)
All	All	1.34	17/4526 (0.4%)	1.25	11/6210 (0.2%)

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	178	GLU	CD-OE1	-7.37	1.17	1.25
1	A	211	GLY	N-CA	6.89	1.56	1.46
1	A	121	TRP	CE3-CZ3	6.62	1.49	1.38
1	A	121	TRP	CB-CG	-6.40	1.38	1.50
1	B	288	GLY	N-CA	6.28	1.55	1.46

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	284	ASP	CB-CG-OD2	-7.27	111.76	118.30
1	A	139	TYR	CB-CG-CD1	-6.95	116.83	121.00
1	B	314	ASP	CB-CG-OD2	-6.81	112.17	118.30
1	A	284	ASP	CB-CG-OD1	6.00	123.70	118.30
1	A	158	TYR	CD1-CE1-CZ	-5.80	114.58	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2195	0	2128	23	1
1	B	2195	0	2128	73	2
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	90	0	118	13	0
3	B	12	0	14	6	0
4	A	291	0	0	11	1
4	B	180	0	0	20	0
All	All	4967	0	4388	106	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:512:GOL:H31	4:A:889:HOH:O	1.24	1.27
1:B:149:SER:HB2	1:B:154:LEU:HD21	1.31	1.13
1:B:149:SER:CB	1:B:154:LEU:HD21	1.93	0.98
1:A:238[B]:ASN:OD1	4:A:748:HOH:O	1.91	0.88
1:A:183[A]:ARG:HD3	4:A:873:HOH:O	1.75	0.84

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88[B]:ASN:ND2	4:A:720:HOH:O[4_555]	2.05	0.15
1:A:61[A]:GLU:OE1	1:B:168:LYS:NZ[4_555]	2.09	0.11

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/319 (85%)	262 (97%)	9 (3%)	0	100	100
1	B	271/319 (85%)	250 (92%)	20 (7%)	1 (0%)	39	19
All	All	542/638 (85%)	512 (94%)	29 (5%)	1 (0%)	52	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	89	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/284 (84%)	238 (100%)	1 (0%)	93	90
1	B	239/284 (84%)	226 (95%)	13 (5%)	27	8
All	All	478/568 (84%)	464 (97%)	14 (3%)	48	25

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	208	GLN
1	B	243	TYR
1	B	274	LEU
1	B	155	THR
1	B	270	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 21 ligands modelled in this entry, 4 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GOL	A	503	-	5,5,5	0.36	0	5,5,5	0.77	0
3	GOL	A	504	-	5,5,5	0.61	0	5,5,5	0.90	0
3	GOL	A	505	-	5,5,5	0.50	0	5,5,5	1.75	1 (20%)
3	GOL	A	506	-	5,5,5	0.53	0	5,5,5	0.52	0
3	GOL	A	507	-	5,5,5	0.49	0	5,5,5	2.15	2 (40%)
3	GOL	A	508	-	5,5,5	0.95	0	5,5,5	1.31	1 (20%)
3	GOL	A	509	-	5,5,5	0.51	0	5,5,5	1.14	1 (20%)
3	GOL	A	510	-	5,5,5	1.28	1 (20%)	5,5,5	1.10	0
3	GOL	A	511	-	5,5,5	0.63	0	5,5,5	0.78	0
3	GOL	A	512	-	5,5,5	1.00	0	5,5,5	3.05	3 (60%)
3	GOL	A	513	-	5,5,5	0.98	0	5,5,5	1.16	0
3	GOL	A	514[A]	-	5,5,5	0.42	0	5,5,5	1.59	1 (20%)
3	GOL	A	514[B]	-	5,5,5	0.52	0	5,5,5	0.49	0
3	GOL	A	515	-	5,5,5	0.45	0	5,5,5	1.88	2 (40%)
3	GOL	A	516	-	5,5,5	2.12	2 (40%)	5,5,5	2.68	2 (40%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	503	-	5,5,5	0.38	0	5,5,5	0.75	0
3	GOL	B	504	-	5,5,5	1.29	1 (20%)	5,5,5	1.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	503	-	-	0/4/4/4	0/0/0/0
3	GOL	A	504	-	-	0/4/4/4	0/0/0/0
3	GOL	A	505	-	-	0/4/4/4	0/0/0/0
3	GOL	A	506	-	-	0/4/4/4	0/0/0/0
3	GOL	A	507	-	-	0/4/4/4	0/0/0/0
3	GOL	A	508	-	-	0/4/4/4	0/0/0/0
3	GOL	A	509	-	-	0/4/4/4	0/0/0/0
3	GOL	A	510	-	-	0/4/4/4	0/0/0/0
3	GOL	A	511	-	-	0/4/4/4	0/0/0/0
3	GOL	A	512	-	-	0/4/4/4	0/0/0/0
3	GOL	A	513	-	-	0/4/4/4	0/0/0/0
3	GOL	A	514[A]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	514[B]	-	-	0/4/4/4	0/0/0/0
3	GOL	A	515	-	-	0/4/4/4	0/0/0/0
3	GOL	A	516	-	-	0/4/4/4	0/0/0/0
3	GOL	B	503	-	-	0/4/4/4	0/0/0/0
3	GOL	B	504	-	-	0/4/4/4	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	516	GOL	C1-C2	-3.66	1.38	1.52
3	A	516	GOL	O3-C3	-2.44	1.31	1.42
3	B	504	GOL	O2-C2	-2.34	1.36	1.43
3	A	510	GOL	O2-C2	2.60	1.51	1.43

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	516	GOL	O1-C1-C2	-5.56	83.20	110.18
3	A	512	GOL	O2-C2-C1	-4.94	86.01	108.65
3	A	507	GOL	O3-C3-C2	-3.60	92.75	110.18
3	A	505	GOL	O2-C2-C3	-3.39	93.12	108.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	512	GOL	O2-C2-C3	-3.05	94.65	108.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	504	GOL	3	0
3	A	507	GOL	1	0
3	A	509	GOL	1	0
3	A	512	GOL	2	0
3	A	513	GOL	2	0
3	A	514[A]	GOL	1	0
3	A	516	GOL	3	0
3	B	503	GOL	2	0
3	B	504	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/319 (84%)	-0.41	5 (1%) 70 73	9, 16, 32, 68	0
1	B	270/319 (84%)	0.58	31 (11%) 6 6	11, 33, 54, 76	0
All	All	540/638 (84%)	0.08	36 (6%) 21 21	9, 22, 50, 76	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	GLY	4.6
1	B	274	LEU	4.3
1	B	71	GLY	4.0
1	B	277	TYR	3.8
1	B	316	PRO	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	A	515	6/6	0.85	0.25	15.46	30,32,32,37	6
3	GOL	B	503	6/6	0.87	0.17	8.15	30,34,36,38	6
3	GOL	B	504	6/6	0.89	0.17	6.16	24,27,28,31	6
3	GOL	A	506	6/6	0.88	0.14	5.92	32,37,46,47	6
3	GOL	A	514[B]	6/6	0.96	0.12	5.67	19,20,24,29	6
3	GOL	A	514[A]	6/6	0.96	0.12	5.53	15,18,20,22	6
3	GOL	A	516	6/6	0.96	0.13	4.51	15,23,29,29	0
3	GOL	A	511	6/6	0.84	0.14	4.47	33,43,44,46	6
3	GOL	A	504	6/6	0.84	0.14	4.32	38,44,46,51	0
3	GOL	A	508	6/6	0.70	0.24	3.10	25,34,36,37	6
3	GOL	A	505	6/6	0.88	0.13	2.40	17,31,38,38	6
3	GOL	A	512	6/6	0.95	0.09	2.40	14,25,29,30	0
3	GOL	A	507	6/6	0.91	0.14	2.27	24,33,36,42	6
3	GOL	A	513	6/6	0.97	0.10	1.58	12,29,33,34	0
3	GOL	A	509	6/6	0.95	0.07	0.61	22,42,44,52	0
2	CA	A	501	1/1	0.98	0.11	-0.62	35,35,35,35	1
2	CA	A	502	1/1	0.99	0.04	-1.19	23,23,23,23	1
2	CA	B	501	1/1	0.78	0.10	-1.39	63,63,63,63	0
2	CA	B	502	1/1	0.99	0.04	-3.09	26,26,26,26	1
3	GOL	A	510	6/6	0.91	0.16	-	22,34,39,43	6
3	GOL	A	503	6/6	0.81	0.13	-	52,55,58,60	0

6.5 Other polymers

There are no such residues in this entry.