



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 07:57 PM GMT

PDB ID : 4WU8  
Title : Structure of trPtNAP-NCP145  
Authors : Chua, E.Y.D.; Davey, C.A.  
Deposited on : 2014-10-31  
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4776 (2.50-2.42)
Clashscore	102246	1030 (2.48-2.44)
Ramachandran outliers	100387	1024 (2.48-2.44)
Sidechain outliers	100360	1024 (2.48-2.44)
RSRZ outliers	91569	4787 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	145	<div> <div>2%</div> <div>52% 38% 10%</div> </div>
2	J	145	<div> <div>52% 36% 12%</div> </div>
3	A	135	<div> <div>62% 8% 29%</div> </div>
3	E	135	<div> <div>2%</div> <div>61% 7% 29%</div> </div>
4	B	102	<div> <div>2%</div> <div>70% 11% 20%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	F	102	
5	C	129	
5	G	129	
6	D	125	
6	H	125	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	CX3	I	100	-	-	-	X
7	CX3	J	101[A]	-	-	-	X
7	CX3	J	101[B]	-	-	-	X
8	SO4	D	201	-	-	-	X
8	SO4	H	201	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 12128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	I	145	Total	C	N	O	P	0	0	0
			2970	1421	538	867	144			

- Molecule 2 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	J	145	Total	C	N	O	P	0	0	0
			2969	1421	535	869	144			

- Molecule 3 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	96	Total	C	N	O	S	0	0	0
			791	500	151	137	3			
3	E	96	Total	C	N	O	S	0	0	0
			791	500	151	137	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	engineered mutation	UNP P84233
E	102	ALA	GLY	engineered mutation	UNP P84233

- Molecule 4 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	82	Total	C	N	O	S	0	0	0
			653	412	127	113	1			
4	F	87	Total	C	N	O	S	0	0	0
			703	442	142	118	1			

- Molecule 5 is a protein called Histone H2A type 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	C	106	Total 818	C 516	N 160	O 142	0	0	0
5	G	106	Total 818	C 516	N 160	O 142	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	99	ARG	GLY	engineered mutation	UNP P06897
C	123	SER	ALA	engineered mutation	UNP P06897
G	99	ARG	GLY	engineered mutation	UNP P06897
G	123	SER	ALA	engineered mutation	UNP P06897


- Molecule 6 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	D	95	Total 745	C 469	N 134	O 140	S 2	0	0	0
6	H	95	Total 745	C 469	N 134	O 140	S 2	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	THR	SER	engineered mutation	UNP P02281
H	29	THR	SER	engineered mutation	UNP P02281

- Molecule 7 is [2-(3-{bis[2-(amino-kappaN)ethyl]amino-kappaN}propyl)-1H-benzo[de]isoquinoline-1,3(2H)-dionato(2-)]platinum(1+) (three-letter code: CX3) (formula: C<sub>19</sub>H<sub>22</sub>N<sub>4</sub>O<sub>2</sub>Pt).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	I	1	Total	C	N	O	Pt	0	0
			26	19	4	2	1		
7	J	1	Total	C	N	O	Pt	0	1
			52	38	8	4	2		

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	D	1	Total	O	S	0	0
			5	4	1		
8	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	E	1	Total	Mg	0	0
			1	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	J	1	Total	O	0	0
			1	1		
10	A	3	Total	O	0	0
			3	3		
10	B	6	Total	O	0	0
			6	6		
10	C	4	Total	O	0	0
			4	4		
10	D	3	Total	O	0	0
			3	3		
10	E	8	Total	O	0	0
			8	8		
10	F	8	Total	O	0	0
			8	8		
10	G	1	Total	O	0	0
			1	1		
10	H	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: DNA (145-MER)




#### • Molecule 2: DNA (145-MER)



#### • Molecule 3: Histone H3.2



#### • Molecule 3: Histone H3.2





SER
GLY
ARG
GLY
LVS
GLY
GLY
LVS
GLY
LEU
GLY
LVS
GLY
GLY
ALA
ALA
LYS
ARG
HIS
ARG
LVS
V21
L22
R23
D24
R23
K31
P32
R35
R39
S47
L62
I66
M84
Y88
G102

SER	GLY	ARG	GLY	LYS	GLY	LYS	GLY	LEU	GLY	LYS	GLY	GLY	ALA
K16	R17	H18	R19	T20	D21	N22	I23	T24	K25	P26	S27	L28	I29
T30	K31	P32	S33	L34	I35	R36	T37	G38	T39	G40	E41	A42	V43

SER	GLY	ARG	GLY	LYS	GLN	GLY	GLY	LYS	THR	ARG	ALA	LYS	A14	K15	T16	R17	H31	R32	R35	G46	K75	I79	P80	R81	H82	L83	Q84	R88	K118	K119	THR	GLU	SER	SER	LYS	SER	ALA	LYS	SER	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.41Å 109.63Å 183.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	94.05 – 2.45 47.09 – 2.45	Depositor EDS
% Data completeness (in resolution range)	93.1 (94.05-2.45) 93.1 (47.09-2.45)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.214 , 0.263 0.213 , 0.261	Depositor DCC
$R_{free}$ test set	1471 reflections (2.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	54.6	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 37.5	EDS
Estimated twinning fraction	0.027 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	1 of 73892 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12128	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CX3, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	I	0.49	6/3332 (0.2%)	1.16	40/5141 (0.8%)
2	J	0.57	7/3330 (0.2%)	1.19	47/5138 (0.9%)
3	A	0.49	0/803	0.69	0/1078
3	E	0.64	0/803	0.85	0/1078
4	B	0.54	0/660	0.71	0/883
4	F	0.72	0/711	0.87	2/948 (0.2%)
5	C	0.63	0/828	0.80	1/1117 (0.1%)
5	G	0.51	0/828	0.72	2/1117 (0.2%)
6	D	0.62	0/756	0.82	3/1015 (0.3%)
6	H	0.58	0/756	0.69	0/1015
All	All	0.56	13/12807 (0.1%)	1.02	95/18530 (0.5%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	-14	DG	O3'-P	-16.53	1.41	1.61
1	I	-14	DG	C5-C4	12.42	1.47	1.38
2	J	-14	DG	C5-C4	11.68	1.46	1.38
2	J	-14	DG	N7-C5	-10.17	1.33	1.39
1	I	-14	DG	N7-C5	-8.54	1.34	1.39

The worst 5 of 95 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	-14	DG	N3-C4-C5	-23.41	116.89	128.60
1	I	-14	DG	N3-C4-C5	-21.59	117.81	128.60
2	J	-14	DG	C2-N3-C4	18.01	120.90	111.90
2	J	-14	DG	N3-C4-N9	17.90	136.74	126.00
1	I	-14	DG	C2-N3-C4	17.85	120.83	111.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	2970	0	1640	41	0
2	J	2969	0	1641	43	0
3	A	791	0	828	7	0
3	E	791	0	828	12	0
4	B	653	0	696	5	0
4	F	703	0	755	5	0
5	C	818	0	877	23	0
5	G	818	0	877	10	0
6	D	745	0	773	14	0
6	H	745	0	773	9	0
7	I	26	0	20	3	0
7	J	52	0	40	6	0
8	D	5	0	0	1	0
8	H	5	0	0	1	0
9	E	1	0	0	0	0
10	A	3	0	0	0	0
10	B	6	0	0	0	0
10	C	4	0	0	0	0
10	D	3	0	0	0	0
10	E	8	0	0	0	0
10	F	8	0	0	1	0
10	G	1	0	0	0	0
10	H	2	0	0	0	0
10	J	1	0	0	0	0
All	All	12128	0	9748	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:49:ARG:HG3	3:E:49:ARG:HH11	1.35	0.92

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:G:17:ARG:HH12	5:G:31:HIS:HD2	1.02	0.92
5:G:17:ARG:HH12	5:G:31:HIS:CD2	1.91	0.87
1:I:62:DT:H2''	1:I:63:DG:O5'	1.81	0.81
2:J:-51:DC:H2''	2:J:-50:DT:H5'	1.63	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	94/135 (70%)	92 (98%)	2 (2%)	0	100	100
3	E	94/135 (70%)	93 (99%)	1 (1%)	0	100	100
4	B	80/102 (78%)	79 (99%)	1 (1%)	0	100	100
4	F	85/102 (83%)	81 (95%)	2 (2%)	2 (2%)	7	4
5	C	104/129 (81%)	100 (96%)	4 (4%)	0	100	100
5	G	104/129 (81%)	100 (96%)	4 (4%)	0	100	100
6	D	93/125 (74%)	89 (96%)	3 (3%)	1 (1%)	17	19
6	H	93/125 (74%)	89 (96%)	3 (3%)	1 (1%)	17	19
All	All	747/982 (76%)	723 (97%)	20 (3%)	4 (0%)	34	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	F	18	HIS
6	D	101	GLY
6	H	101	GLY
4	F	26	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	84/110 (76%)	81 (96%)	3 (4%)	42	58
3	E	84/110 (76%)	81 (96%)	3 (4%)	42	58
4	B	67/78 (86%)	65 (97%)	2 (3%)	48	65
4	F	72/78 (92%)	68 (94%)	4 (6%)	26	36
5	C	84/101 (83%)	73 (87%)	11 (13%)	5	4
5	G	84/101 (83%)	81 (96%)	3 (4%)	42	58
6	D	81/105 (77%)	76 (94%)	5 (6%)	23	31
6	H	81/105 (77%)	72 (89%)	9 (11%)	8	8
All	All	637/788 (81%)	597 (94%)	40 (6%)	22	30

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	D	30	ARG
3	E	59	GLU
6	H	84	SER
6	D	120	SER
3	E	129	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
3	E	68	GLN
6	H	92	GLN
5	G	31	HIS
6	D	92	GLN
4	F	18	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
8	SO4	D	201	-	4,4,4	0.71	0	6,6,6	0.58	0
8	SO4	H	201	-	4,4,4	0.74	0	6,6,6	0.71	0
7	CX3	I	100	1	22,30,30	2.07	4 (18%)	20,45,45	2.47	3 (15%)
7	CX3	J	101[A]	2	22,30,30	1.90	4 (18%)	20,45,45	1.83	3 (15%)
7	CX3	J	101[B]	-	22,30,30	1.84	4 (18%)	20,45,45	1.76	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	SO4	D	201	-	-	0/0/0/0	0/0/0/0
8	SO4	H	201	-	-	0/0/0/0	0/0/0/0
7	CX3	I	100	1	-	1/4/27/27	0/3/5/5
7	CX3	J	101[A]	2	-	0/4/27/27	0/3/5/5
7	CX3	J	101[B]	-	-	0/4/27/27	0/3/5/5

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	J	101[B]	CX3	C12-N4	2.25	1.41	1.38
7	J	101[A]	CX3	C12-N4	2.37	1.41	1.38
7	J	101[A]	CX3	C8-N4	2.52	1.42	1.38
7	J	101[B]	CX3	C8-N4	2.54	1.42	1.38
7	I	100	CX3	C12-N4	3.12	1.42	1.38

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	J	101[A]	CX3	C7-N4-C8	2.04	121.10	119.10
7	J	101[B]	CX3	C11-C12-N4	4.39	119.72	116.09
7	J	101[B]	CX3	C9-C8-N4	4.74	120.00	116.09
7	J	101[A]	CX3	C9-C8-N4	4.77	120.03	116.09
7	J	101[A]	CX3	C11-C12-N4	5.25	120.43	116.09

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	I	100	CX3	C6-C7-N4-C12

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	D	201	SO4	1	0
8	H	201	SO4	1	0
7	I	100	CX3	3	0
7	J	101[A]	CX3	1	0
7	J	101[B]	CX3	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	I	145/145 (100%)	-0.27	1 (0%) 89 90	41, 94, 136, 165	0
2	J	145/145 (100%)	-0.29	0 100 100	49, 95, 140, 168	0
3	A	96/135 (71%)	-0.17	0 100 100	30, 47, 70, 121	0
3	E	96/135 (71%)	-0.22	1 (1%) 84 86	24, 35, 55, 107	0
4	B	82/102 (80%)	-0.13	2 (2%) 62 65	31, 43, 68, 124	0
4	F	87/102 (85%)	0.07	2 (2%) 64 66	24, 33, 57, 164	0
5	C	106/129 (82%)	-0.16	1 (0%) 85 87	24, 40, 67, 119	0
5	G	106/129 (82%)	-0.17	1 (0%) 85 87	33, 48, 83, 107	0
6	D	95/125 (76%)	-0.06	1 (1%) 82 84	30, 42, 75, 118	0
6	H	95/125 (76%)	0.02	1 (1%) 82 84	31, 47, 92, 132	0
All	All	1053/1272 (82%)	-0.15	10 (0%) 85 87	24, 48, 122, 168	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	16	LYS	9.5
5	C	119	LYS	6.0
6	H	122	LYS	5.4
6	D	122	LYS	4.0
4	B	21	VAL	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	CX3	J	101[A]	26/26	0.86	0.40	24.50	96,100,107,108	26
8	SO4	D	201	5/5	0.96	0.24	6.46	59,60,66,72	0
7	CX3	J	101[B]	26/26	0.86	0.40	5.82	112,125,157,159	26
8	SO4	H	201	5/5	0.97	0.19	2.47	57,58,63,76	0
7	CX3	I	100	26/26	0.88	0.27	2.39	134,148,186,194	0
9	MG	E	201	1/1	0.98	0.12	-0.57	30,30,30,30	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.