



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:43 PM GMT

PDB ID : 3WVJ  
Title : The crystal structure of native glycosidic hydrolase  
Authors : Chen, C.C.; Huang, J.W.; Zhao, P.; Ko, T.P.; Huang, C.H.; Chan, H.C.;  
Huang, Z.; Liu, W.; Cheng, Y.S.; Liu, J.R.; Guo, R.T.  
Deposited on : 2014-05-22  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	<div> <div>7%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>..</div> </div> </div>
1	B	220	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>18%</div> <div>..</div> </div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1792	1164	289	334	5			
1	B	220	Total	C	N	O	S	0	0	0
			1802	1169	291	337	5			

- Molecule 2 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYL AMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: B3P) (formula:  $C_{11}H_{26}N_2O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	11	2	6		
2	A	1	Total	C	N	O	0	0
			19	11	2	6		
2	B	1	Total	C	N	O	0	0
			19	11	2	6		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			19	11	2	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	164	Total	O	0	0
			164	164		
3	B	178	Total	O	0	0
			178	178		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Beta-glucanase



#### • Molecule 1: Beta-glucanase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.45Å 74.45Å 182.92Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 1.95 24.37 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (25.00-1.95) 99.9 (24.37-1.95)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.14 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.190 , 0.232 0.192 , 0.234	Depositor DCC
$R_{free}$ test set	2194 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.6	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 37.0	EDS
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 43664 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4012	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.09	3/1855 (0.2%)	1.13	9/2523 (0.4%)
1	B	1.22	7/1865 (0.4%)	1.30	14/2535 (0.6%)
All	All	1.16	10/3720 (0.3%)	1.22	23/5058 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	153	TRP	CD2-CE2	7.33	1.50	1.41
1	B	182	TRP	CD2-CE2	7.17	1.50	1.41
1	B	96	ARG	CD-NE	-6.86	1.34	1.46
1	A	182	TRP	CD2-CE2	6.10	1.48	1.41
1	B	223	TRP	CD2-CE2	6.06	1.48	1.41
1	B	109	ARG	CD-NE	-5.42	1.37	1.46
1	A	54	TRP	CD2-CE2	5.36	1.47	1.41
1	B	215	TRP	CD2-CE2	5.21	1.47	1.41
1	B	54	TRP	CD2-CE2	5.11	1.47	1.41
1	B	182	TRP	CG-CD1	5.07	1.43	1.36

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	96	ARG	NE-CZ-NH2	-20.62	109.99	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	96	ARG	NE-CZ-NH2	-17.52	111.54	120.30
1	A	96	ARG	NE-CZ-NH1	16.20	128.40	120.30
1	B	109	ARG	NE-CZ-NH2	-15.97	112.31	120.30
1	B	96	ARG	NE-CZ-NH1	14.36	127.48	120.30
1	B	109	ARG	NE-CZ-NH1	10.80	125.70	120.30
1	B	109	ARG	CG-CD-NE	-8.14	94.71	111.80
1	B	228	ASP	N-CA-CB	-7.51	97.08	110.60
1	A	96	ARG	CD-NE-CZ	6.85	133.19	123.60
1	B	96	ARG	CD-NE-CZ	6.05	132.07	123.60
1	B	205	VAL	CG1-CB-CG2	5.97	120.46	110.90
1	A	145	ASP	CB-CG-OD2	5.91	123.62	118.30
1	B	96	ARG	CG-CD-NE	-5.89	99.42	111.80
1	B	218	ILE	CB-CA-C	-5.55	100.50	111.60
1	B	201	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	A	81	ASP	CB-CG-OD1	5.52	123.27	118.30
1	B	98	LYS	CD-CE-NZ	-5.47	99.12	111.70
1	B	183	ARG	C-N-CD	5.43	139.80	128.40
1	A	132	ASN	C-N-CD	5.27	139.46	128.40
1	A	96	ARG	CG-CD-NE	-5.22	100.84	111.80
1	A	109	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	230	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	B	109	ARG	CD-NE-CZ	5.06	130.69	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	227	TYR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1792	0	1665	24	0
1	B	1802	0	1673	33	1
2	A	38	0	52	0	0
2	B	38	0	52	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	164	0	0	4	0
3	B	178	0	0	7	0
All	All	4012	0	3442	53	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:148:LYS:HE3	1:B:166:ASN:HD21	1.39	0.87
1:B:34:ASN:HB3	3:B:435:HOH:O	1.75	0.87
1:B:147:THR:HG23	3:B:498:HOH:O	1.78	0.81
1:A:131:ASN:HD21	1:B:51:LYS:NZ	1.82	0.75
1:B:165:HIS:HD2	3:B:507:HOH:O	1.69	0.75
1:A:230:ARG:HD2	3:A:409:HOH:O	1.87	0.73
1:B:155:LYS:HD2	1:B:203:ILE:O	1.90	0.72
1:B:109:ARG:HD3	1:B:238:GLU:OE2	1.89	0.71
1:B:148:LYS:CE	1:B:166:ASN:HD21	2.04	0.69
1:B:165:HIS:HE1	3:B:429:HOH:O	1.75	0.68
1:A:75:LYS:HD3	1:A:175:PHE:CZ	2.32	0.65
1:A:82:ARG:NH2	1:A:229:GLY:O	2.31	0.64
1:B:155:LYS:HE3	3:B:557:HOH:O	1.99	0.63
1:B:176:HIS:HD2	1:B:192:ASP:OD2	1.82	0.62
1:A:131:ASN:HD21	1:B:51:LYS:HZ1	1.47	0.60
1:B:61:ASN:HB2	1:B:218:ILE:HG13	1.84	0.59
1:A:176:HIS:HE1	3:A:480:HOH:O	1.86	0.58
1:B:145:ASP:OD1	1:B:147:THR:HB	2.04	0.57
1:A:170:ASP:HB3	1:A:173:GLN:HG2	1.87	0.57
1:A:115:ASN:HD21	1:A:228:ASP:HB2	1.70	0.56
1:A:176:HIS:HD2	1:A:192:ASP:OD2	1.88	0.56
1:A:222:GLU:H	1:A:222:GLU:CD	2.09	0.56
1:A:30:ALA:N	3:A:459:HOH:O	2.38	0.56
1:A:115:ASN:OD1	1:A:230:ARG:HG2	2.06	0.56
1:A:46:SER:HB2	3:A:551:HOH:O	2.08	0.53
1:A:128:PRO:HB2	1:B:66:PRO:HG2	1.90	0.52
1:B:195:LYS:O	1:B:195:LYS:HG3	2.09	0.50
1:B:61:ASN:HB2	1:B:218:ILE:CG1	2.41	0.50
1:B:148:LYS:HE3	1:B:166:ASN:ND2	2.18	0.49
1:B:164:LEU:HB2	3:B:474:HOH:O	2.13	0.48
1:A:51:LYS:HE3	1:B:131:ASN:HD21	1.79	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:228:ASP:OD2	1:B:230:ARG:CB	2.62	0.48
1:B:147:THR:HG22	1:B:147:THR:O	2.14	0.47
1:B:176:HIS:CD2	1:B:192:ASP:OD2	2.67	0.47
1:B:42:SER:O	1:B:43:ASN:HB2	2.15	0.47
1:B:125:TYR:CD1	2:B:302:B3P:H62	2.50	0.47
1:A:136:GLU:OE2	1:A:138:ASP:OD1	2.33	0.46
1:B:109:ARG:HE	1:B:177:THR:HG22	1.79	0.46
1:A:96:ARG:HB3	1:A:211:MET:HG2	1.98	0.46
1:B:228:ASP:OD2	1:B:230:ARG:HB3	2.15	0.46
1:B:103:TYR:CZ	1:B:184:PRO:HG3	2.51	0.46
1:A:131:ASN:HD21	1:B:51:LYS:HZ2	1.62	0.45
1:A:58:SER:OG	1:A:59:VAL:N	2.50	0.44
1:B:134:TRP:CD1	1:B:134:TRP:C	2.90	0.44
1:B:147:THR:O	1:B:147:THR:CG2	2.64	0.44
1:A:245:ASN:CG	1:A:246:GLY:N	2.71	0.43
1:A:230:ARG:HB2	1:A:230:ARG:HE	1.48	0.43
1:A:180:PHE:HA	1:A:188:ASP:O	2.19	0.42
1:A:183:ARG:HB3	1:A:184:PRO:CD	2.50	0.41
1:B:109:ARG:HH21	1:B:177:THR:CG2	2.34	0.41
1:A:109:ARG:HD3	1:A:238:GLU:OE1	2.21	0.41
2:B:301:B3P:H22	2:B:301:B3P:H112	1.54	0.41
1:B:165:HIS:CD2	3:B:507:HOH:O	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ASN:OD1	1:B:202:ASN:ND2[4_565]	2.05	0.15

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/220 (99%)	204 (94%)	13 (6%)	0	100	100
1	B	218/220 (99%)	207 (95%)	10 (5%)	1 (0%)	34	21
All	All	435/440 (99%)	411 (94%)	23 (5%)	1 (0%)	52	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	43	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/189 (100%)	182 (97%)	6 (3%)	46	33
1	B	189/189 (100%)	172 (91%)	17 (9%)	12	3
All	All	377/378 (100%)	354 (94%)	23 (6%)	23	9

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	SER
1	A	73	ASN
1	A	96	ARG
1	A	148	LYS
1	A	222	GLU
1	A	230	ARG
1	B	31	THR
1	B	32	VAL
1	B	73	ASN
1	B	75	LYS
1	B	96	ARG
1	B	109	ARG
1	B	129	SER
1	B	156	ASN
1	B	174	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	177	THR
1	B	194	LYS
1	B	195	LYS
1	B	205	VAL
1	B	218	ILE
1	B	226	ARG
1	B	230	ARG
1	B	249	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	131	ASN
1	A	166	ASN
1	A	176	HIS
1	A	202	ASN
1	B	131	ASN
1	B	156	ASN
1	B	165	HIS
1	B	166	ASN
1	B	173	GLN
1	B	176	HIS
1	B	202	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	B3P	A	301	-	18,18,18	1.68	3 (16%)	17,23,23	2.67	6 (35%)
2	B3P	A	302	-	18,18,18	3.53	7 (38%)	17,23,23	2.33	5 (29%)
2	B3P	B	301	-	18,18,18	1.14	2 (11%)	17,23,23	2.94	8 (47%)
2	B3P	B	302	-	18,18,18	2.29	4 (22%)	17,23,23	2.24	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B3P	A	301	-	-	0/28/28/28	0/0/0/0
2	B3P	A	302	-	-	0/28/28/28	0/0/0/0
2	B3P	B	301	-	-	0/28/28/28	0/0/0/0
2	B3P	B	302	-	-	0/28/28/28	0/0/0/0

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	B3P	C5-C4	-5.11	1.48	1.53
2	A	302	B3P	C9-C8	-4.61	1.49	1.53
2	B	302	B3P	C10-C8	-4.12	1.49	1.53
2	A	301	B3P	C11-C8	-4.06	1.49	1.53
2	A	301	B3P	C6-C4	-3.38	1.50	1.53
2	A	301	B3P	C7-C4	-2.76	1.51	1.53
2	A	302	B3P	O6-C7	2.08	1.48	1.42
2	B	301	B3P	C7-C4	2.24	1.55	1.53
2	A	302	B3P	C1-C2	2.83	1.63	1.51
2	A	302	B3P	C6-C4	2.93	1.56	1.53
2	B	301	B3P	C10-C8	3.70	1.56	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	302	B3P	C3-N1	3.86	1.53	1.46
2	B	302	B3P	C7-C4	4.14	1.57	1.53
2	B	302	B3P	C6-C4	4.52	1.57	1.53
2	A	302	B3P	C5-C4	5.94	1.58	1.53
2	A	302	B3P	C7-C4	11.05	1.63	1.53

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	B3P	C2-N2-C8	-6.62	106.01	116.10
2	B	302	B3P	C11-C8-C10	-6.49	98.25	110.16
2	A	301	B3P	C7-C4-C6	-5.93	99.28	110.16
2	A	302	B3P	O6-C7-C4	-5.62	99.43	111.47
2	A	301	B3P	O2-C10-C8	-4.94	100.89	111.47
2	A	301	B3P	C2-N2-C8	-4.52	109.21	116.10
2	B	301	B3P	O2-C10-C8	-4.30	102.27	111.47
2	A	302	B3P	C11-C8-C9	-3.96	102.89	110.16
2	B	302	B3P	C3-N1-C4	-3.94	110.10	116.10
2	B	301	B3P	C7-C4-C6	-3.81	103.17	110.16
2	B	301	B3P	C11-C8-C9	-3.69	103.40	110.16
2	A	301	B3P	C10-C8-C9	-3.36	103.99	110.16
2	A	301	B3P	O4-C5-C4	-3.25	104.51	111.47
2	A	302	B3P	O3-C11-C8	-3.25	104.51	111.47
2	A	302	B3P	C2-N2-C8	-2.99	111.54	116.10
2	B	301	B3P	O4-C5-C4	-2.88	105.30	111.47
2	B	302	B3P	O5-C6-C4	-2.65	105.79	111.47
2	B	301	B3P	C1-C3-N1	-2.27	103.82	112.08
2	A	302	B3P	O5-C6-C4	-2.21	106.72	111.47
2	A	301	B3P	C6-C4-C5	2.10	114.02	110.16
2	B	301	B3P	C3-N1-C4	3.41	121.30	116.10
2	B	301	B3P	C10-C8-C9	4.02	117.54	110.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	B3P	1	0
2	B	302	B3P	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	219/220 (99%)	0.24	16 (7%) 18 28	28, 42, 67, 88	0
1	B	220/220 (100%)	0.01	9 (4%) 41 52	28, 38, 53, 96	0
All	All	439/440 (99%)	0.13	25 (5%) 27 37	28, 39, 62, 96	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	30	ALA	7.7
1	A	30	ALA	5.7
1	B	31	THR	5.1
1	A	84	TYR	4.8
1	B	249	GLN	4.4
1	A	139	ILE	4.2
1	A	122	PHE	3.8
1	A	248	PRO	3.4
1	A	245	ASN	3.4
1	A	247	VAL	3.1
1	B	222	GLU	3.1
1	A	31	THR	3.0
1	B	122	PHE	2.9
1	A	108	VAL	2.8
1	A	173	GLN	2.8
1	A	123	PHE	2.7
1	A	180	PHE	2.7
1	B	34	ASN	2.7
1	B	137	ILE	2.6
1	A	34	ASN	2.6
1	A	174	ASP	2.5
1	B	139	ILE	2.3
1	B	32	VAL	2.2
1	A	210	ILE	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	189	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	B3P	B	301	19/19	0.87	0.13	1.06	40,51,64,66	0
2	B3P	A	302	19/19	0.93	0.10	-0.11	22,27,44,48	0
2	B3P	A	301	19/19	0.93	0.09	-0.16	27,38,52,52	0
2	B3P	B	302	19/19	0.93	0.11	-0.21	26,35,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.