



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:24 AM GMT

PDB ID : 2WYV  
Title : High resolution structure of Thermus thermophilus enoyl-acyl carrier protein reductase NAD-form  
Authors : Otero, J.M.; Noel, A.J.; Guardado-Calvo, P.; Llamas-Saiz, A.L.; Van Raaij, M.J.  
Deposited on : 2009-11-20  
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	<div> <div>90%</div> <div>8%</div> <div>.</div> </div>
1	B	261	<div> <div>3%</div> <div>85%</div> <div>10%</div> <div>5%</div> </div>
1	C	261	<div> <div>7%</div> <div>89%</div> <div>7%</div> <div>.</div> </div>
1	D	261	<div> <div>4%</div> <div>90%</div> <div>8%</div> <div>.</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENOYL-[ACYL CARRIER PROTEIN] REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	1	0
			1956	1250	338	361	7			
1	B	247	Total	C	N	O	S	0	0	0
			1877	1197	324	350	6			
1	C	253	Total	C	N	O	S	0	1	0
			1930	1232	334	357	7			
1	D	257	Total	C	N	O	S	0	1	0
			1953	1246	338	362	7			

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		
2	A	1	Total	Na	0	0
			1	1		
2	C	1	Total	Na	0	0
			1	1		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	161	Total	O	0	0
			161	161		
4	B	135	Total	O	0	0
			135	135		
4	C	117	Total	O	0	0
			117	117		
4	D	132	Total	O	0	0
			132	132		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: ENOYL-[ACYL CARRIER PROTEIN] REDUCTASE




- Molecule 1: ENOYL-[ACYL CARRIER PROTEIN] REDUCTASE



- Molecule 1: ENOYL-[ACYL CARRIER PROTEIN] REDUCTASE



- Molecule 1: ENOYL-[ACYL CARRIER PROTEIN] REDUCTASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.27Å 107.80Å 85.98Å 90.00° 95.89° 90.00°	Depositor
Resolution (Å)	39.75 – 1.86 39.75 – 1.86	Depositor EDS
% Data completeness (in resolution range)	95.9 (39.75-1.86) 94.0 (39.75-1.86)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.75 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, $R_{free}$	0.182 , 0.239 0.187 , 0.243	Depositor DCC
$R_{free}$ test set	1098 reflections (1.43%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtriage
Anisotropy	0.661	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 77755 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8352	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	0/1992	0.72	0/2697
1	B	0.75	1/1905 (0.1%)	0.73	0/2578
1	C	0.75	0/1963	0.73	1/2655 (0.0%)
1	D	0.76	0/1988	0.75	1/2691 (0.0%)
All	All	0.75	1/7848 (0.0%)	0.73	2/10621 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	151	GLU	CG-CD	5.38	1.60	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	145	LEU	CA-CB-CG	5.41	127.74	115.30
1	C	76	LEU	CA-CB-CG	5.07	126.95	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1956	0	2007	34	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1877	0	1922	24	0
1	C	1930	0	1981	14	0
1	D	1953	0	2004	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	B	44	0	26	7	0
3	D	44	0	26	0	0
4	A	161	0	0	3	0
4	B	135	0	0	0	0
4	C	117	0	0	1	0
4	D	132	0	0	1	0
All	All	8352	0	7966	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:LEU:HD13	1:A:33:ALA:HB1	1.33	1.10
1:B:26:ILE:O	1:B:30:LEU:HD23	1.59	1.03
1:D:132:ALA:O	1:D:136:LEU:HD23	1.62	0.99
1:A:4:VAL:HG13	1:B:2:LEU:HB3	1.57	0.84
1:C:3:THR:HG22	1:D:3:THR:HG22	1.60	0.81
1:B:67:VAL:HG11	1:B:94:ILE:HD13	1.63	0.80
1:D:132:ALA:HB1	1:D:136:LEU:HD21	1.63	0.80
1:A:3:THR:HG22	1:B:3:THR:HG22	1.66	0.78
1:B:45:ARG:HG3	1:B:46:LEU:HD13	1.65	0.76
1:A:4:VAL:CG1	1:B:2:LEU:HB3	2.18	0.73
1:B:17:THR:HG23	3:B:1260:NAD:O3B	1.87	0.73
1:A:6:LEU:HD13	1:A:33:ALA:CB	2.16	0.73
1:A:4:VAL:HG22	1:A:6:LEU:HD11	1.72	0.71
1:A:67:VAL:HA	1:A:73:LEU:HD11	1.73	0.71
1:A:207:MET:CE	4:A:2114:HOH:O	2.38	0.71
1:A:6:LEU:HD12	1:A:6:LEU:N	2.09	0.67
1:A:229:LEU:HD11	1:A:233:LEU:HD21	1.78	0.66
1:A:6:LEU:CD1	1:A:33:ALA:HB1	2.21	0.66
1:D:133:GLU:OE1	1:D:181:LYS:HE3	1.96	0.65
1:A:229:LEU:CD1	1:A:233:LEU:HD21	2.28	0.64
1:A:2:LEU:HB3	1:B:4:VAL:CG2	2.28	0.63

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:LEU:HD12	1:A:233:LEU:CD2	2.27	0.63
1:D:132:ALA:HB1	1:D:136:LEU:CD2	2.29	0.62
1:B:17:THR:CG2	3:B:1260:NAD:O2B	2.48	0.61
1:A:2:LEU:HB3	1:B:4:VAL:HG22	1.82	0.60
1:C:26:ILE:O	1:C:30:LEU:HD23	2.01	0.60
1:B:20:ARG:HG3	1:B:195:THR:HG23	1.82	0.60
1:A:229:LEU:O	1:A:233:LEU:HD23	2.01	0.60
1:A:26:ILE:O	1:A:30:LEU:HD23	2.02	0.59
1:A:229:LEU:CD1	1:A:233:LEU:CD2	2.80	0.59
1:A:4:VAL:HG11	1:B:2:LEU:HD23	1.84	0.58
1:A:2:LEU:HD22	4:A:2001:HOH:O	2.05	0.56
1:A:207:MET:HE2	4:A:2114:HOH:O	2.05	0.55
1:A:69:GLN:O	1:A:73:LEU:HD13	2.08	0.54
1:D:132:ALA:C	1:D:136:LEU:HD23	2.28	0.53
1:A:4:VAL:CG1	1:B:2:LEU:HD23	2.39	0.53
1:A:4:VAL:HG22	1:A:6:LEU:CD1	2.38	0.53
1:B:16:VAL:HG12	1:B:53:LEU:HD13	1.90	0.52
1:A:26:ILE:O	1:A:30:LEU:CD2	2.58	0.52
1:C:63:PHE:HB3	1:C:76:LEU:HD22	1.92	0.52
1:A:49:GLU:HA	1:A:52:LYS:HE3	1.92	0.51
1:D:198:ALA:HB1	1:D:204:PHE:CE1	2.46	0.51
1:C:221:ILE:HD13	1:C:248:VAL:O	2.11	0.51
1:A:36:GLU:OE1	1:A:61:LEU:HD11	2.11	0.51
1:A:147[A]:TYR:OH	1:A:207:MET:HG2	2.11	0.50
1:A:61:LEU:H	1:A:61:LEU:HD12	1.77	0.50
1:D:248:VAL:HG13	1:D:248:VAL:O	2.12	0.49
1:B:67:VAL:HG11	1:B:94:ILE:CD1	2.37	0.49
1:C:18:ASN:ND2	1:C:196:VAL:HB	2.27	0.49
1:A:207:MET:HB2	1:C:257:MET:HE1	1.95	0.48
1:D:149:ALA:HA	1:D:152:LYS:O	2.15	0.47
1:B:95:ALA:HB2	3:B:1260:NAD:O3D	2.14	0.47
1:A:232:PHE:HD2	1:A:233:LEU:HD22	1.79	0.46
1:B:155:PRO:O	1:B:156:LYS:HB2	2.15	0.46
1:B:17:THR:HG23	3:B:1260:NAD:O2B	2.15	0.46
1:C:188:ILE:HG23	1:C:229:LEU:HD23	1.98	0.46
1:D:145:LEU:HD13	1:D:188:ILE:HB	1.97	0.46
1:A:147[A]:TYR:HD1	1:A:190:ALA:O	1.99	0.46
1:B:95:ALA:HB3	3:B:1260:NAD:H3D	1.98	0.46
1:D:198:ALA:HA	1:D:201:ILE:HD12	1.98	0.46
1:C:218:ARG:NH1	4:C:2097:HOH:O	2.44	0.45
1:D:256:GLY:O	1:D:257:MET:HB2	2.16	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:MET:HB3	1:D:207:MET:HE1	1.99	0.45
1:B:17:THR:CG2	3:B:1260:NAD:O3B	2.61	0.44
1:C:63:PHE:CB	1:C:76:LEU:HD22	2.48	0.44
1:D:188:ILE:HG23	1:D:229:LEU:HD23	1.99	0.44
1:C:20:ARG:NH1	1:C:194:ARG:O	2.50	0.44
1:D:46:LEU:HD22	1:D:46:LEU:N	2.31	0.44
1:A:6:LEU:CD1	1:A:6:LEU:N	2.80	0.43
1:B:47:ARG:HB3	1:B:48:PRO:HD3	2.00	0.43
1:C:61:LEU:N	1:C:61:LEU:CD1	2.81	0.43
1:C:76:LEU:HD13	1:C:76:LEU:C	2.39	0.43
1:D:13:VAL:HG21	1:D:27:ALA:HB2	2.01	0.43
1:D:46:LEU:CD2	1:D:46:LEU:N	2.81	0.42
1:B:17:THR:HG21	3:B:1260:NAD:O2B	2.17	0.42
1:C:18:ASN:HD21	1:C:196:VAL:HB	1.84	0.42
1:B:99:ARG:O	1:B:103:GLU:HG3	2.20	0.41
1:C:49:GLU:HA	1:C:52:LYS:HE3	2.02	0.41
1:D:45:ARG:HG2	1:D:46:LEU:HD22	2.03	0.40
1:A:130:ARG:HD3	4:D:2062:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/261 (98%)	249 (97%)	7 (3%)	0	100	100
1	B	243/261 (93%)	236 (97%)	5 (2%)	2 (1%)	24	9
1	C	250/261 (96%)	244 (98%)	6 (2%)	0	100	100
1	D	256/261 (98%)	247 (96%)	9 (4%)	0	100	100
All	All	1005/1044 (96%)	976 (97%)	27 (3%)	2 (0%)	52	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	194	ARG
1	B	148	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/197 (99%)	195 (100%)	0	100	100
1	B	187/197 (95%)	185 (99%)	2 (1%)	80	72
1	C	192/197 (98%)	189 (98%)	3 (2%)	70	57
1	D	195/197 (99%)	192 (98%)	3 (2%)	72	60
All	All	769/788 (98%)	761 (99%)	8 (1%)	82	76

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	46	LEU
1	B	105	ARG
1	C	6	LEU
1	C	76	LEU
1	C	208	TYR
1	D	39	LEU
1	D	135	LEU
1	D	145	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAD	B	1260	-	38,48,48	1.76	3 (7%)	47,73,73	2.17	5 (10%)
3	NAD	D	1258	-	38,48,48	1.58	3 (7%)	47,73,73	2.00	8 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	B	1260	-	-	0/22/62/62	0/5/5/5
3	NAD	D	1258	-	-	0/22/62/62	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	1258	NAD	C2A-N1A	2.45	1.38	1.33
3	B	1260	NAD	C2A-N1A	2.86	1.39	1.33
3	B	1260	NAD	C2A-N3A	3.93	1.39	1.32
3	D	1258	NAD	C2A-N3A	4.10	1.39	1.32
3	D	1258	NAD	O7N-C7N	7.31	1.39	1.24
3	B	1260	NAD	O7N-C7N	8.68	1.42	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1260	NAD	N3A-C2A-N1A	-10.79	120.64	128.89
3	D	1258	NAD	N3A-C2A-N1A	-9.89	121.32	128.89
3	D	1258	NAD	C4A-C5A-N7A	-3.62	106.15	109.48
3	B	1260	NAD	C2B-C1B-N9A	-3.32	109.21	114.29
3	D	1258	NAD	O7N-C7N-N7N	-2.79	118.67	122.59
3	D	1258	NAD	C4B-O4B-C1B	-2.45	107.02	109.72
3	D	1258	NAD	PN-O3-PA	-2.00	127.11	132.73
3	B	1260	NAD	O3-PA-O5B	2.74	110.19	102.94
3	D	1258	NAD	C3N-C7N-N7N	3.03	121.13	117.82
3	B	1260	NAD	O4B-C1B-N9A	3.21	114.81	108.10
3	D	1258	NAD	O4B-C1B-N9A	3.43	115.28	108.10
3	D	1258	NAD	O4D-C1D-N1N	3.79	112.30	108.13
3	B	1260	NAD	O4D-C1D-N1N	6.53	115.31	108.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1260	NAD	7	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	257/261 (98%)	-0.04	2 (0%) 87 87	15, 25, 41, 53	0
1	B	247/261 (94%)	-0.05	7 (2%) 56 54	16, 26, 42, 59	0
1	C	253/261 (96%)	0.19	17 (6%) 21 19	15, 26, 45, 74	0
1	D	257/261 (98%)	-0.06	10 (3%) 43 40	14, 25, 44, 62	0
All	All	1014/1044 (97%)	0.01	36 (3%) 46 44	14, 26, 43, 74	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	198	ALA	8.5
1	C	197	ALA	7.2
1	C	196	VAL	7.2
1	C	208	TYR	5.8
1	B	208	TYR	5.7
1	C	207	MET	5.3
1	C	206	LYS	5.1
1	C	199	ARG	5.1
1	B	195	THR	5.1
1	C	195	THR	4.5
1	D	200	SER	4.2
1	C	194	ARG	4.2
1	D	198	ALA	3.9
1	D	202	PRO	3.4
1	C	20	ARG	3.1
1	D	257	MET	3.0
1	D	196	VAL	3.0
1	D	201	ILE	3.0
1	B	20	ARG	3.0
1	B	210	ARG	2.8
1	C	209	ASP	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	210	ARG	2.5
1	A	257	MET	2.5
1	C	4	VAL	2.5
1	D	45	ARG	2.4
1	B	193	VAL	2.3
1	C	99	ARG	2.3
1	D	203	GLY	2.2
1	D	1	MET	2.2
1	A	199	ARG	2.2
1	B	194	ARG	2.2
1	B	100	GLU	2.1
1	D	199	ARG	2.1
1	C	1	MET	2.1
1	C	213	GLN	2.1
1	C	156	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAD	B	1260	44/44	0.66	0.25	1.88	60,72,77,77	0
2	NA	C	1260	1/1	0.98	0.24	1.37	34,34,34,34	0
2	NA	A	1258	1/1	0.96	0.15	0.80	35,35,35,35	0
2	NA	B	1261	1/1	0.99	0.17	0.36	25,25,25,25	0
3	NAD	D	1258	44/44	0.96	0.08	-0.43	21,27,30,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.