



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 06:28 PM BST

PDB ID : 1X0F  
Title : Complex structure of the C-terminal RNA-binding domain of hnRNP D(AUF1) with telomeric DNA  
Authors : Enokizono, Y.; Konishi, Y.; Nagata, K.; Ouhashi, K.; Uesugi, S.; Ishikawa, F.; Katahira, M.  
Deposited on : 2005-03-22

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	4	100%
2	A	79	6% 46% 34% 13%

## 2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:181-A:213, A:224-A:259 (69)	0.14	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 2 single-model clusters were found.

Cluster number	Models
1	5, 7, 8, 9, 12, 14
2	1, 2, 10, 19
3	3, 4, 6
4	17, 18, 20
5	11, 15
Single-model clusters	13; 16

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1417 atoms, of which 696 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(P\*TP\*AP\*GP\*G)-3'.

Mol	Chain	Residues	Atoms						Trace
1	B	4	Total	C	H	N	O	P	0
			130	40	45	17	24	4	

- Molecule 2 is a protein called Heterogeneous nuclear ribonucleoprotein D0.

Mol	Chain	Residues	Atoms						Trace
2	A	79	Total	C	H	N	O	S	0
			1287	409	651	105	117	5	

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-D(P\*TP\*AP\*GP\*G)-3'

Chain B:  100%

T2  
A3  
G4  
G5

- Molecule 2: Heterogeneous nuclear ribonucleoprotein D0

Chain A:  6% 46% 34% 13%

V181 K182 K183 K184 F185 V186 G187 G188 L189 S190 P191 D192 T193 P194 E195 E196 K197 I198 R199 E200 Y201 F202 G203 G204 F205 G206 E207 V208 E209 S210 T211 E212 L213 F214 M215 D216 D217 N218 T219 M220 K221 R222 R223 G224 F225 C226 F227 T228 T229 T230 K231 E232 E233 E234 P235 V236 K237 R238 T239 M240

E241 K242 K243 Y244 H245 N246 V247 G248 L249 S250 K251 C252 E253 I254 K255 V256 A257 M258 S259

### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 8. Colouring as in section 4.1 above.

- Molecule 1: 5'-D(P\*TP\*AP\*GP\*G)-3'

Chain B:  100%

T2  
A3  
G4  
G5

- Molecule 2: Heterogeneous nuclear ribonucleoprotein D0

Chain A:  9% 42% 34% 13%

V181 K182 K183 K184 F185 V186 G187 G188 L189 S190 P191 D192 T193 P194 E195 E196 K197 I198 R199 E200 Y201 F202 G203 G204 F205 G206 E207 V208 E209 S210 T211 E212 L213 F214 M215 D216 D217 N218 T219 M220 K221 R222 R223 G224 F225 C226 F227 T228 T229 T230 K231 E232 E233 E234 P235 V236 K237 R238 T239 M240 E241

K242
K243
Y244
H245
K246
Y247
G248
L249
S250
K251
C252
E253
I254
K255
Y256
A257
K258
S259

## 5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing*.

Of the 228 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
X-PLOR	refinement	3.851

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	85	45	46	27±6
2	A	550	560	559	172±9
All	All	12700	12100	12100	3653

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 147.

5 of 591 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:A:227:PHE:CE2	2:A:257:ALA:HB1	1.07	1.84	19	19
2:A:202:PHE:CE2	2:A:228:ILE:HD12	1.04	1.86	4	20
2:A:227:PHE:CE1	2:A:257:ALA:HB1	1.02	1.90	10	1
2:A:227:PHE:CD2	2:A:257:ALA:HB1	0.95	1.96	14	19
1:B:4:DG:H2''	1:B:5:DG:O5'	0.95	1.58	14	12

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR



entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	67/79 (85%)	55±1 (82±2%)	6±2 (9±2%)	6±1 (8±2%)	2	14
All	All	1340/1580 (85%)	1103 (82%)	124 (9%)	113 (8%)	2	14

5 of 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	190	SER	20
2	A	192	ASP	20
2	A	248	GLY	20
2	A	256	VAL	20
2	A	224	GLY	17

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	61/71 (86%)	34±2 (55±3%)	27±2 (45±3%)	0	2
All	All	1220/1420 (86%)	675 (55%)	545 (45%)	0	2

5 of 41 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	211	ILE	20
2	A	205	PHE	20
2	A	183	LYS	20
2	A	255	LYS	20
2	A	190	SER	20

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided