



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 8, 2016 – 12:37 PM EST

PDB ID : 2X1N
Title : Truncation and Optimisation of Peptide Inhibitors of CDK2, Cyclin A Through Structure Guided Design
Authors : Kontopidis, G.; Andrews, M.J.; McInnes, C.; Plater, A.; Innes, L.; Renachowski, S.; Cowan, A.; Fischer, P.M.; McIntyre, N.A.; Griffiths, G.; Barnett, A.L.; Slawin, A.M.Z.; Jackson, W.; Thomas, M.; Zheleva, D.I.; Wang, S.; Blake, D.G.; Westwood, N.J.
Deposited on : 2009-12-31
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div>4%</div> <div> <div></div> <div>79%</div> <div>16%</div> <div>..</div> </div> </div>
1	C	298	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>13%</div> <div>..</div> </div> </div>
2	B	261	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>..</div> </div> </div>
2	D	261	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>..</div> </div> </div>
3	H	5	<div> <div>20%</div> <div> <div></div> <div>60%</div> <div>40%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9118 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELL DIVISION PROTEIN KINASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	1	0
			2384	1551	403	422	8			
1	C	297	Total	C	N	O	S	0	2	1
			2393	1557	408	420	8			


- Molecule 2 is a protein called CYCLIN-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	258	Total	C	N	O	S	0	1	0
			2091	1355	342	383	11			
2	D	258	Total	C	N	O	S	0	0	0
			2084	1350	339	384	11			

- Molecule 3 is a protein called ACE-LEU-ASN-PFF-NH2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	5	Total	C	F	N	O	0	0	1
			32	21	1	5	5			

- Molecule 4 is 2-METHYL-N-[(1Z)-3-NITROCYCLOHEXA-2,4-DIEN-1-YLIDENE]-4,5-DIHYDRO[1,3]THIAZOLO[4,5-H]QUINAZOLIN-8-AMINE (three-letter code: X1N) (formula: C₁₆H₁₃N₅O₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			24	16	5	2	1		
4	C	1	Total	C	N	O	S	0	0
			24	16	5	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	22	Total	O	0	0
			22	22		
5	B	24	Total	O	0	0
			24	24		
5	C	17	Total	O	0	0
			17	17		
5	D	21	Total	O	0	0
			21	21		
5	H	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CELL DIVISION PROTEIN KINASE 2




• Molecule 1: CELL DIVISION PROTEIN KINASE 2



• Molecule 2: CYCLIN-A2



• Molecule 2: CYCLIN-A2



• Molecule 3: ACE-LEU-ASN-PFF-NH2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.56Å 114.26Å 157.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.75 14.96 – 2.75	Depositor EDS
% Data completeness (in resolution range)	98.0 (30.00-2.75) 98.6 (14.96-2.75)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.200 , 0.255 0.204 , 0.254	Depositor DCC
R_{free} test set	1097 reflections (3.24%)	DCC
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.384	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9118	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NH2, X1N, ACE, PFF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2449	0.83	2/3325 (0.1%)
1	C	0.62	1/2461 (0.0%)	0.85	5/3340 (0.1%)
2	B	0.70	1/2144 (0.0%)	0.78	2/2910 (0.1%)
2	D	0.65	1/2134 (0.0%)	0.82	1/2897 (0.0%)
3	H	1.77	1/16 (6.2%)	1.58	0/21
All	All	0.65	4/9204 (0.0%)	0.82	10/12493 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	175	VAL	N-CA	11.37	1.69	1.46
3	H	3	ASN	CB-CG	-6.67	1.35	1.51
2	D	175	VAL	N-CA	6.51	1.59	1.46
1	C	177	CYS	CB-SG	-6.09	1.72	1.82

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	38	ASP	CB-CG-OD1	6.13	123.82	118.30
1	A	256	ASP	CB-CG-OD2	5.72	123.45	118.30
1	A	92	ASP	CB-CG-OD1	5.59	123.33	118.30
1	C	64	VAL	CB-CA-C	-5.46	101.02	111.40
2	B	216	ASP	CB-CG-OD1	5.38	123.15	118.30
2	B	424	LEU	CB-CG-CD2	-5.30	101.98	111.00
1	C	1	MET	CG-SD-CE	5.21	108.54	100.20
2	D	175	VAL	CB-CA-C	5.16	121.20	111.40
1	C	196	MET	CG-SD-CE	5.08	108.32	100.20
1	C	177	CYS	CA-CB-SG	-5.04	104.93	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2384	0	2432	58	0
1	C	2393	0	2452	47	0
2	B	2091	0	2120	24	0
2	D	2084	0	2107	20	0
3	H	32	0	27	1	0
4	A	24	0	13	1	0
4	C	24	0	13	0	0
5	A	22	0	0	0	0
5	B	24	0	0	0	0
5	C	17	0	0	0	0
5	D	21	0	0	0	0
5	H	2	0	0	0	0
All	All	9118	0	9164	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (140) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:175:VAL:CA	2:B:175:VAL:N	1.69	1.51
1:A:41:THR:HG22	1:A:42:GLU:H	1.08	1.11
1:C:164:VAL:HG12	1:C:165:THR:H	1.18	1.02
2:B:282:THR:O	2:B:285:THR:HG23	1.63	0.99
1:C:39:THR:O	1:C:40:GLU:HB2	1.63	0.96
1:A:36:ARG:HG2	1:A:36:ARG:HH11	1.24	0.96
1:A:39:THR:OG1	2:B:289:LYS:HD3	1.66	0.94
1:A:162:GLU:OE1	1:A:180:TYR:OH	1.89	0.91
2:D:176:PRO:HB3	2:D:179:HIS:ND1	1.88	0.89
1:A:41:THR:HG22	1:A:42:GLU:N	1.86	0.88
1:C:166:LEU:O	1:C:168:TYR:N	2.09	0.86
1:A:41:THR:CG2	1:A:42:GLU:H	1.91	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:ARG:HG2	1:A:36:ARG:NH1	1.83	0.82
1:A:36:ARG:O	1:A:37:LEU:HD13	1.81	0.81
1:C:164:VAL:HG12	1:C:165:THR:N	1.97	0.79
1:A:36:ARG:CG	1:A:36:ARG:HH11	1.97	0.77
1:A:60:HIS:HD2	1:A:62:ASN:H	1.34	0.76
1:C:162:GLU:HG2	1:C:163:VAL:N	2.02	0.75
1:A:40:GLU:O	1:A:41:THR:HB	1.87	0.74
2:B:205:ASP:OD2	2:B:250:ARG:HD2	1.88	0.72
1:C:15:TYR:OH	1:C:51:GLU:OE1	2.04	0.71
1:C:51:GLU:O	1:C:55:LEU:HB2	1.91	0.71
1:A:164:VAL:O	1:A:164:VAL:CG1	2.40	0.69
1:A:164:VAL:HG13	1:A:164:VAL:O	1.92	0.69
2:B:282:THR:O	2:B:285:THR:CG2	2.40	0.69
1:A:39:THR:O	1:A:40:GLU:HB2	1.94	0.68
1:A:2:GLU:HG2	1:C:73:GLU:HG3	1.76	0.68
1:A:163:VAL:HG22	1:A:163:VAL:O	1.92	0.68
2:B:176:PRO:HA	2:B:179:HIS:ND1	2.09	0.67
2:D:404:HIS:HD2	2:D:406:GLN:H	1.43	0.67
2:B:175:VAL:C	2:B:175:VAL:N	2.48	0.66
1:C:162:GLU:HG3	1:C:169:ARG:HH22	1.60	0.66
2:D:346:PRO:O	2:D:349:LYS:HG2	1.96	0.66
1:A:40:GLU:O	1:A:41:THR:CB	2.44	0.66
1:A:39:THR:O	1:A:40:GLU:CB	2.43	0.65
1:A:162:GLU:HG3	1:A:164:VAL:HB	1.79	0.64
1:C:95:ALA:O	1:C:96:LEU:HD23	1.98	0.64
2:D:343:ASP:HB3	2:D:345:ASP:OD2	1.97	0.63
1:A:231:THR:HG22	1:A:236:TYR:CZ	2.35	0.62
1:A:60:HIS:CD2	1:A:62:ASN:H	2.18	0.62
2:D:207:THR:HG23	2:D:210:MET:H	1.65	0.62
1:C:164:VAL:CG1	1:C:165:THR:H	2.03	0.61
2:B:207:THR:HG23	2:B:210:MET:H	1.64	0.61
1:C:165:THR:O	1:C:166:LEU:HB2	2.00	0.60
1:C:253:PRO:HB2	1:C:254:PRO:HD3	1.84	0.60
1:C:160:THR:C	1:C:162:GLU:H	2.06	0.59
1:C:60:HIS:HD2	1:C:62:ASN:H	1.50	0.59
1:A:49:ILE:HG23	2:B:306:LEU:HD12	1.83	0.59
1:C:60:HIS:CD2	1:C:62:ASN:H	2.20	0.59
1:A:14:THR:O	1:A:15:TYR:HB2	2.03	0.59
1:A:95:ALA:O	1:A:96:LEU:HD23	2.03	0.59
1:C:161:HIS:O	1:C:162:GLU:O	2.19	0.59
1:A:39:THR:OG1	2:B:289:LYS:CD	2.45	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:332:LEU:HD13	2:B:421:VAL:HB	1.86	0.58
1:A:39:THR:HG22	2:B:292:LEU:HB3	1.84	0.58
1:A:2:GLU:HG2	1:C:73:GLU:CG	2.33	0.58
1:C:162:GLU:CG	1:C:163:VAL:N	2.66	0.58
4:A:1297:X1N:N3	4:A:1297:X1N:H2B	2.18	0.58
2:D:207:THR:HG22	2:D:210:MET:HG3	1.86	0.58
1:A:154:VAL:HG12	1:A:155:PRO:HD2	1.86	0.57
1:A:115:LEU:HD11	1:A:185:ASP:HB3	1.86	0.56
1:C:165:THR:O	1:C:166:LEU:CB	2.52	0.56
1:C:162:GLU:HG2	1:C:163:VAL:H	1.71	0.56
1:C:72:THR:HG22	1:C:73:GLU:N	2.22	0.54
2:B:176:PRO:HB3	2:B:179:HIS:ND1	2.22	0.54
1:A:41:THR:CG2	1:A:42:GLU:N	2.58	0.54
1:A:51:GLU:O	1:A:55:LEU:HB2	2.08	0.54
1:C:1:MET:HE3	1:C:1:MET:HA	1.90	0.54
2:B:336:LEU:HD13	2:B:362:LEU:HD23	1.91	0.53
1:C:5:GLN:HB2	1:C:24:LYS:HE2	1.91	0.53
1:C:64:VAL:CG2	1:C:143:LEU:O	2.57	0.53
2:D:176:PRO:CB	2:D:179:HIS:ND1	2.69	0.52
1:A:6:LYS:NZ	1:A:34:LYS:NZ	2.57	0.52
1:A:36:ARG:C	1:A:37:LEU:HD13	2.30	0.52
1:C:115:LEU:HD11	1:C:185:ASP:HB3	1.90	0.52
1:A:33:LYS:HB3	1:A:78:LEU:HB2	1.92	0.51
2:B:265:SER:HB3	2:B:272:PRO:HB3	1.90	0.51
2:D:219:VAL:HG22	2:D:232:LEU:HD11	1.91	0.51
1:A:84:HIS:O	1:A:85:GLN:HB3	2.10	0.51
1:A:227:TRP:CD2	1:A:230:VAL:HG13	2.46	0.51
2:D:229:ASN:HD22	2:D:334:MET:CE	2.24	0.51
2:D:404:HIS:CD2	2:D:406:GLN:H	2.26	0.50
2:D:329:VAL:HG11	2:D:364:LEU:HD13	1.94	0.49
1:A:227:TRP:O	1:A:230:VAL:HG22	2.11	0.49
1:A:91:MET:CE	1:A:195:GLU:HG2	2.43	0.49
1:C:64:VAL:HG22	1:C:143:LEU:O	2.13	0.49
1:A:71:HIS:CE1	2:B:296:HIS:CE1	3.00	0.49
2:D:178:TYR:HD2	2:D:178:TYR:C	2.16	0.49
1:A:73:GLU:CD	1:C:2:GLU:HG2	2.33	0.48
2:D:289:LYS:HZ2	2:D:293:ARG:HD2	1.78	0.48
1:A:212:LEU:HD22	1:A:216:PHE:CZ	2.48	0.48
1:A:36:ARG:HD3	1:A:36:ARG:O	2.13	0.48
1:C:127:ASP:OD1	1:C:165:THR:HG23	2.13	0.48
2:B:327:CYS:HB3	2:B:419:HIS:NE2	2.29	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:176:PRO:CA	2:B:179:HIS:ND1	2.77	0.47
1:C:164:VAL:CG1	1:C:165:THR:N	2.70	0.46
1:C:162:GLU:HG2	1:C:164:VAL:H	1.80	0.46
2:D:178:TYR:CD2	2:D:178:TYR:C	2.88	0.46
1:A:121:HIS:O	1:A:122:ARG:HG3	2.15	0.46
2:D:345:ASP:OD2	2:D:346:PRO:HD3	2.15	0.46
1:A:6:LYS:HZ1	1:A:34:LYS:NZ	2.14	0.46
1:C:86:ASP:OD2	1:C:89:LYS:HD3	2.15	0.46
1:C:15:TYR:HD2	1:C:35:ILE:HG12	1.82	0.45
2:B:289:LYS:O	2:B:293:ARG:HG2	2.16	0.45
1:C:74:ASN:N	1:C:74:ASN:OD1	2.50	0.45
1:C:39:THR:O	1:C:40:GLU:CB	2.45	0.45
1:A:37:LEU:HA	1:A:37:LEU:HD12	1.72	0.45
2:D:229:ASN:HD22	2:D:334:MET:HE2	1.82	0.44
1:A:101:LEU:N	1:A:102:PRO:CD	2.79	0.44
1:A:230:VAL:HG23	1:A:231:THR:HG23	1.99	0.44
1:A:39:THR:O	1:A:40:GLU:CG	2.66	0.44
1:C:115:LEU:HD23	1:C:115:LEU:HA	1.82	0.44
2:D:207:THR:CG2	2:D:210:MET:HG3	2.47	0.44
1:A:159:TYR:N	1:A:159:TYR:CD1	2.86	0.43
1:C:161:HIS:O	1:C:162:GLU:C	2.57	0.43
1:C:64:VAL:CG2	1:C:144:ALA:HA	2.48	0.43
2:D:367:VAL:HG12	2:D:368:THR:HG23	1.99	0.43
1:A:163:VAL:CG2	1:A:163:VAL:O	2.63	0.43
2:B:430:LEU:HD23	2:B:430:LEU:HA	1.83	0.43
1:C:160:THR:C	1:C:162:GLU:N	2.71	0.42
2:B:323:GLN:HA	2:B:324:PRO:HA	1.85	0.42
1:C:72:THR:HG22	1:C:73:GLU:H	1.83	0.42
1:C:253:PRO:CB	1:C:254:PRO:HD3	2.49	0.42
1:C:189:LEU:HD12	1:C:189:LEU:HA	1.62	0.42
1:C:124:LEU:HD21	1:C:182:THR:HA	2.02	0.42
1:A:121:HIS:C	1:A:122:ARG:HG3	2.40	0.42
1:C:64:VAL:HG23	1:C:143:LEU:O	2.20	0.41
1:A:159:TYR:N	1:A:159:TYR:HD1	2.17	0.41
2:B:279:VAL:O	2:B:282:THR:OG1	2.33	0.41
1:C:71:HIS:CE1	2:D:296:HIS:NE2	2.89	0.41
1:A:202:LEU:HD13	1:A:203:PHE:CE2	2.56	0.41
1:A:35:ILE:C	1:A:36:ARG:HG3	2.41	0.41
1:A:121:HIS:HD2	2:B:185:TYR:CE1	2.39	0.41
1:A:227:TRP:CE3	1:A:230:VAL:CG1	3.04	0.41
1:C:288:ASP:N	1:C:288:ASP:OD1	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:271:TYR:HD1	2:B:271:TYR:O	2.03	0.41
2:D:210:MET:HG2	3:H:4:PFF:HB2	2.03	0.41
1:A:193:PHE:CZ	1:A:255:LEU:HD21	2.56	0.40
1:C:17:VAL:HG13	1:C:19:TYR:CE1	2.57	0.40
1:A:227:TRP:CG	1:A:230:VAL:HG13	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/298 (99%)	283 (96%)	8 (3%)	4 (1%)	14	38
1	C	297/298 (100%)	280 (94%)	8 (3%)	9 (3%)	5	16
2	B	257/261 (98%)	250 (97%)	7 (3%)	0	100	100
2	D	256/261 (98%)	252 (98%)	2 (1%)	2 (1%)	24	55
3	H	2/5 (40%)	2 (100%)	0	0	100	100
All	All	1107/1123 (99%)	1067 (96%)	25 (2%)	15 (1%)	14	38

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	TYR
1	A	36	ARG
1	A	41	THR
1	C	162	GLU
1	C	163	VAL
1	C	167	TRP
1	C	40	GLU
1	C	41	THR
1	C	166	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	177	ASP
2	D	346	PRO
1	C	15	TYR
1	C	160	THR
1	A	40	GLU
1	C	38	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/263 (100%)	245 (94%)	17 (6%)	21	48
1	C	263/263 (100%)	241 (92%)	22 (8%)	14	34
2	B	233/235 (99%)	225 (97%)	8 (3%)	44	76
2	D	232/235 (99%)	225 (97%)	7 (3%)	48	80
3	H	2/2 (100%)	2 (100%)	0	100	100
All	All	992/998 (99%)	938 (95%)	54 (5%)	27	58

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ARG
1	A	37	LEU
1	A	55	LEU
1	A	56	LYS
1	A	71	HIS
1	A	74	ASN
1	A	75	LYS
1	A	96	LEU
1	A	115	LEU
1	A	122	ARG
1	A	126	ARG
1	A	165	THR
1	A	200	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	202	LEU
1	A	230	VAL
1	A	248	PHE
1	A	287	GLN
2	B	199	TYR
2	B	206	ILE
2	B	250	ARG
2	B	271	TYR
2	B	292	LEU
2	B	323	GLN
2	B	374	GLU
2	B	375	SER
1	C	1	MET
1	C	2	GLU
1	C	38	ASP
1	C	39	THR
1	C	41	THR
1	C	64	VAL
1	C	71	HIS
1	C	74	ASN
1	C	75	LYS
1	C	87	LEU
1	C	96	LEU
1	C	115	LEU
1	C	122	ARG
1	C	158	THR
1	C	161	HIS
1	C	162	GLU
1	C	163	VAL
1	C	217	ARG
1	C	232	SER
1	C	247	ASP
1	C	248	PHE
1	C	252	VAL
2	D	178	TYR
2	D	188	GLU
2	D	196	LYS
2	D	292	LEU
2	D	328	LYS
2	D	345	ASP
2	D	364	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	71	HIS
1	A	85	GLN
2	B	208	ASN
2	B	296	HIS
2	B	323	GLN
2	B	404	HIS
1	C	60	HIS
1	C	71	HIS
1	C	85	GLN
1	C	246	GLN
2	D	208	ASN
2	D	404	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PFF	H	4	3	10,12,13	0.72	0	13,15,17	2.10	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PFF	H	4	3	-	0/4/6/8	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	4	PFF	O-C-CA	-3.36	116.72	125.72
3	H	4	PFF	F-CZ-CE1	-2.69	114.24	118.53
3	H	4	PFF	CG-CB-CA	3.91	123.16	114.12
3	H	4	PFF	F-CZ-CE2	4.13	125.11	118.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	4	PFF	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	X1N	A	1297	-	21,27,27	3.36	7 (33%)	17,39,39	3.70	8 (47%)
4	X1N	C	1298	-	21,27,27	3.48	7 (33%)	17,39,39	3.39	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	X1N	A	1297	-	-	0/3/27/27	0/4/4/4
4	X1N	C	1298	-	-	0/3/27/27	0/4/4/4

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1298	X1N	C20-C3A	-5.85	1.39	1.49
4	A	1297	X1N	C20-C3A	-5.37	1.40	1.49
4	A	1297	X1N	C6B-C5B	-4.49	1.39	1.49
4	C	1298	X1N	C4-C5A	-4.45	1.38	1.47
4	C	1298	X1N	C6B-C5B	-4.28	1.39	1.49
4	A	1297	X1N	C4-C5A	-3.89	1.39	1.47
4	C	1298	X1N	C2B-C1B	-3.01	1.38	1.44
4	A	1297	X1N	C2B-C1B	-2.33	1.39	1.44
4	C	1298	X1N	C4-C5	-2.11	1.37	1.41
4	A	1297	X1N	C1B-N7	2.39	1.35	1.29
4	C	1298	X1N	C4B-C5B	3.00	1.39	1.32
4	A	1297	X1N	C4B-C5B	3.05	1.39	1.32
4	A	1297	X1N	O8B-N7B	11.73	1.46	1.22
4	C	1298	X1N	O8B-N7B	12.07	1.46	1.22

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1297	X1N	C6B-C5B-C4B	-11.59	117.38	123.51
4	C	1298	X1N	C6B-C5B-C4B	-9.70	118.38	123.51
4	A	1297	X1N	N1-C2-N3	-4.09	119.33	125.56
4	C	1298	X1N	N1-C2-N3	-4.09	119.33	125.56
4	A	1297	X1N	C5-C6-N1	-4.07	116.75	123.86
4	C	1298	X1N	C5-C6-N1	-3.71	117.38	123.86
4	A	1297	X1N	C5A-C4-C5	-3.16	117.50	120.55
4	C	1298	X1N	C5A-C4-C5	-2.53	118.11	120.55
4	A	1297	X1N	C60-C6A-C1A	2.12	111.42	109.84
4	C	1298	X1N	C5A-C4-N3	2.13	120.81	117.98
4	A	1297	X1N	C5A-C4-N3	2.58	121.40	117.98
4	C	1298	X1N	C60-C6A-C1A	3.07	112.13	109.84
4	A	1297	X1N	C20-C3A-N2A	3.10	132.03	123.88
4	C	1298	X1N	C20-C3A-N2A	4.20	134.93	123.88
4	C	1298	X1N	C60-C5-C4	4.22	123.80	119.08
4	A	1297	X1N	C60-C5-C4	4.93	124.60	119.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1297	X1N	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/298 (99%)	-0.12	11 (3%) 45 38	21, 37, 73, 98	1 (0%)
1	C	297/298 (99%)	-0.08	15 (5%) 32 24	19, 37, 74, 99	0
2	B	258/261 (98%)	-0.18	7 (2%) 58 51	21, 41, 61, 99	0
2	D	258/261 (98%)	-0.18	4 (1%) 74 70	21, 40, 60, 99	0
3	H	2/5 (40%)	1.32	1 (50%) 0 0	52, 52, 52, 77	0
All	All	1111/1123 (98%)	-0.13	38 (3%) 49 42	19, 38, 70, 99	1 (0%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	39	THR	11.1
2	D	175	VAL	7.0
1	C	13	GLY	5.2
1	C	14	THR	5.1
1	C	40	GLU	5.1
2	B	175	VAL	5.0
1	A	15	TYR	4.8
1	C	295	HIS	4.2
1	C	96	LEU	3.8
1	C	36	ARG	3.5
1	A	73	GLU	3.3
3	H	3	ASN	3.3
2	B	271	TYR	3.3
2	B	323	GLN	3.3
1	A	74	ASN	3.2
1	A	36	ARG	3.2
1	A	38	ASP	3.2
1	C	73	GLU	3.1
1	A	295	HIS	3.0
1	A	14	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	161	HIS	2.9
1	C	38	ASP	2.7
1	A	244	ALA	2.7
1	A	247	ASP	2.6
2	B	201	LYS	2.5
1	C	15	TYR	2.5
2	D	176	PRO	2.4
1	C	37	LEU	2.4
2	D	432	LEU	2.4
1	C	200	ARG	2.3
1	C	297	ARG	2.3
1	A	95	ALA	2.3
2	B	176	PRO	2.2
1	A	200	ARG	2.2
1	C	74	ASN	2.2
2	B	378[A]	ARG	2.1
2	B	180	GLU	2.1
2	D	226	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PFF	H	4	12/13	0.79	0.24	-	40,49,68,69	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	X1N	C	1298	24/24	0.90	0.22	0.22	41,52,86,95	0
4	X1N	A	1297	24/24	0.93	0.16	-0.29	32,47,72,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.