wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 06:20 PM BST

PDB ID	:	1X5Q
Title	:	Solution structure of the first PDZ domain of scribble homolog protein (hScrib)
Authors	:	Qin, X.R.; Nagashima, T.; Hayashi, F.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2005-05-16

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 91%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	110	42% 35% • 20%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 15 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:13-A:23, A:28-A:104 (88)	0.27	15

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20
2	2, 8

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1561 atoms, of which 773 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called LAP4 protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	110	1561	478	773	146	163	1	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q14160
A	2	SER	-	CLONING ARTIFACT	UNP Q14160
A	3	SER	-	CLONING ARTIFACT	UNP Q14160
A	4	GLY	-	CLONING ARTIFACT	UNP Q14160
A	5	SER	-	CLONING ARTIFACT	UNP Q14160
A	6	SER	-	CLONING ARTIFACT	UNP Q14160
A	7	GLY	-	CLONING ARTIFACT	UNP Q14160
A	105	SER	-	CLONING ARTIFACT	UNP Q14160
A	106	GLY	-	CLONING ARTIFACT	UNP Q14160
A	107	PRO	-	CLONING ARTIFACT	UNP Q14160
A	108	SER	-	CLONING ARTIFACT	UNP Q14160
A	109	SER	-	CLONING ARTIFACT	UNP Q14160
A	110	GLY	-	CLONING ARTIFACT	UNP Q14160

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: LAP4 protein

Chain A: 42% 35% • 20%

G1 S2 S3 G4 S5 S6 G7 E8 P9 A10 R11 I12 E13 E14 E15 E16 L17 T18 L19 T20 I21 L22 R23 Q24 T25 G26 G27 L28 G29 I30 S38 T39 P40 F49 I50 S51 R52 V53 A59 A62 G63 V64 D68 K69 L70 L71 E72 V73 N74 G75 V76 Q79 E82

H83 H84 E85 E88 A89 A93 G94 T95 A96 V97 Q98 M99 R100 V101 W102 R103 E104 S105 G106 P107 S108 S109 G110

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 15. Colouring as in section 4.1 above.

- Molecule 1: LAP4 protein

Chain A: 40% 35% • • 20%

G1 S2 S3 G4 S5 S6 G7 E8 P9 A10 R11 I12 E13 E14 E15 E16 L17 R23 Q24 T25 G26 G27 L28 G29 I30 Y41 K42 E46 F49 I50 S51 R52 V53 S54 P58 A59 A60 R61 A62 V66 G67 D68 K69 L70 L71 E72 V73 N74 G75 V76 A77 L78 Q79

E82 H83 H84 E85 E88 A89 A93 G94 T95 A96 V97 Q98 M99 R100 V101 W102 R103 E104 S105 G106 P107 S108 S109 G110

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *Torsion angle dynamics.*

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function, structures with the lowest energy, structures with the least restraint violations.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.7
CYANA	refinement	1.0.7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 11207
Number of chemical shift lists	1
Total number of shifts	1222
Number of shifts mapped to atoms	1222
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	91%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	655	653	653	34±4
All	All	13100	13060	13060	682

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

5 of 188 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:73:VAL:O	1:A:76:VAL:HG12	0.94	1.63	11	20
1:A:70:LEU:HD21	1:A:73:VAL:HG23	0.93	1.38	12	2
1:A:93:ALA:HB1	1:A:97:VAL:HG22	0.92	1.42	5	17
1:A:22:LEU:HD12	1:A:23:ARG:N	0.90	1.80	13	1
1:A:70:LEU:HD22	1:A:101:VAL:HG12	0.86	1.47	20	8

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	88/110 (80%)	72±2 (82±3%)	13±3 (15±3%)	3±1 (3±1%)	9 42
All	All	1760/2200 (80%)	1443 (82%)	265 (15%)	52 (3%)	9 42

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	95	THR	16
1	A	14	GLU	13
1	A	28	LEU	7
1	A	35	GLY	4
1	A	43	GLY	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	65/79 (82%)	52±3 (80±4%)	13±3 (20±4%)	4 35
All	All	1300/1580 (82%)	1038 (80%)	262 (20%)	4 35

5 of 34 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	88	GLU	19
1	A	74	ASN	19
1	A	70	LEU	14
1	A	17	LEU	14
1	A	68	ASP	13

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 91% for the well-defined parts and 88% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 11207

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1222
Number of shifts mapped to atoms	1222
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	106	-0.02 ± 0.10	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	88	-0.27 ± 0.17	None needed (< 0.5 ppm)
$^{13}C'$	105	-0.04 ± 0.09	None needed (< 0.5 ppm)
^{15}N	98	-0.66 ± 0.27	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 91%, i.e. 925 atoms were assigned a chemical shift out of a possible 1021. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	434/436 (100%)	173/174 (99%)	176/176 (100%)	85/86 (99%)
Sidechain	454/542 (84%)	276/313 (88%)	173/202 (86%)	5/27 (19%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	37/43 (86%)	19/23 (83%)	17/17 (100%)	1/3 (33%)
Overall	925/1021 (91%)	468/510 (92%)	366/395 (93%)	91/116 (78%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: