



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:12 PM GMT

PDB ID : 4X5O
Title : Human histidine tRNA synthetase
Authors : Kim, Y.K.; Jeon, Y.H.
Deposited on : 2014-12-05
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	509	<div> <div>2%</div> <div>65%</div> <div>20%</div> <div>•</div> <div>15%</div> </div>
1	B	509	<div> <div>4%</div> <div>53%</div> <div>15%</div> <div>31%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6256 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histidine-tRNA ligase, cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	434	Total	C	N	O	S	0	0	0
			3429	2185	586	641	17			
1	B	350	Total	C	N	O	S	0	0	0
			2816	1804	478	522	12			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	O	0	0
			4	4		
2	B	7	Total	O	0	0
			7	7		



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	97.67Å 97.67Å 254.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.02 – 2.80 29.02 – 2.79	Depositor EDS
% Data completeness (in resolution range)	97.4 (29.02-2.80) 96.9 (29.02-2.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.80Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.248 , 0.314 0.266 , 0.282	Depositor DCC
R_{free} test set	1540 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	66.0	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 30688 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6256	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/3478	0.47	0/4679
1	B	0.25	0/2854	0.49	1/3834 (0.0%)
All	All	0.25	0/6332	0.48	1/8513 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	366	LEU	CA-CB-CG	6.49	130.22	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	329	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3429	0	3498	70	0
1	B	2816	0	2902	51	0
2	A	4	0	0	0	0
2	B	7	0	0	0	0
All	All	6256	0	6400	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:MET:HB3	1:A:257:LYS:HE3	1.47	0.97
1:A:164:THR:H	1:A:165:ARG:HA	1.42	0.84
1:A:257:LYS:HE2	1:A:259:LEU:HD22	1.64	0.78
1:A:227:SER:HB2	1:A:230:LYS:HG2	1.71	0.71
1:A:232:ARG:NH2	1:A:329:ASP:OD1	2.25	0.69
1:B:330:TYR:HD2	1:B:366:LEU:HD22	1.56	0.69
1:B:92:ILE:HG12	1:B:149:ARG:HE	1.60	0.67
1:B:119:ASP:O	1:B:121:GLY:N	2.26	0.67
1:B:331:TYR:HD2	1:B:335:ILE:HG12	1.59	0.67
1:A:340:LEU:HG	1:A:357:VAL:HG21	1.77	0.67
1:A:257:LYS:HD3	1:A:259:LEU:HB2	1.76	0.66
1:A:257:LYS:NZ	1:A:264:ALA:HB2	2.11	0.66
1:A:260:ALA:HB3	1:A:263:VAL:HG23	1.79	0.65
1:A:418:LYS:O	1:A:445:ASN:ND2	2.29	0.65
1:B:363:TYR:CD1	1:B:366:LEU:HB3	2.32	0.65
1:A:428:VAL:HG21	1:A:440:LEU:HD23	1.78	0.63
1:A:56:LEU:HD11	1:B:370:PHE:HB3	1.82	0.62
1:B:306:PHE:HA	1:B:309:LEU:HD12	1.81	0.62
1:B:363:TYR:HD1	1:B:366:LEU:HB3	1.65	0.61
1:A:425:LEU:HD23	1:A:440:LEU:HD21	1.82	0.61
1:A:130:ASP:OD2	1:A:132:THR:OG1	2.17	0.61
1:A:253:MET:HB3	1:A:257:LYS:CE	2.29	0.60
1:A:257:LYS:HZ3	1:A:264:ALA:HB2	1.67	0.60
1:A:244:VAL:HG12	1:A:245:SER:H	1.67	0.59
1:A:257:LYS:HB2	1:A:259:LEU:H	1.67	0.58
1:A:310:THR:HA	1:A:315:ASP:HB2	1.86	0.57
1:B:148:LYS:HG3	1:B:178:ILE:HG12	1.87	0.57
1:A:253:MET:CB	1:A:257:LYS:HE3	2.29	0.57
1:B:72:VAL:HG21	1:B:394:GLU:HB2	1.86	0.56
1:A:213:ASP:HB3	1:A:216:ILE:HG13	1.85	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:376:LYS:HG3	1:A:378:PRO:HD3	1.86	0.56
1:B:413:VAL:HG21	1:B:428:VAL:HG22	1.88	0.56
1:B:159:ASP:H	1:B:169:ARG:HG2	1.70	0.56
1:A:164:THR:N	1:A:165:ARG:HA	2.14	0.55
1:B:336:TYR:OH	1:B:381:GLY:O	2.19	0.54
1:A:498:ILE:O	1:A:502:THR:OG1	2.25	0.54
1:B:315:ASP:N	1:B:315:ASP:OD1	2.40	0.53
1:B:316:ASP:N	1:B:316:ASP:OD1	2.42	0.52
1:A:97:PHE:HB3	1:A:127:LEU:HD23	1.92	0.52
1:A:202:LEU:HD13	1:A:389:ILE:HG21	1.93	0.51
1:B:475:VAL:HG13	1:B:488:ASP:HB2	1.92	0.51
1:A:139:LEU:HD12	1:A:367:VAL:HG11	1.93	0.50
1:A:479:ARG:HB2	1:A:486:GLU:HG2	1.93	0.50
1:B:411:VAL:HG21	1:B:498:ILE:HD13	1.94	0.49
1:B:80:ILE:HD13	1:B:198:ILE:HG21	1.93	0.49
1:A:54:PHE:HE1	1:B:99:LEU:HD12	1.77	0.49
1:A:159:ASP:O	1:A:169:ARG:NE	2.28	0.49
1:A:418:LYS:N	1:A:418:LYS:HD3	2.27	0.49
1:A:257:LYS:HD3	1:A:259:LEU:H	1.77	0.49
1:B:214:ARG:NH1	1:B:218:ASP:OD2	2.46	0.49
1:B:325:ALA:O	1:B:328:LEU:HB2	2.13	0.48
1:A:115:TYR:OH	1:A:169:ARG:HD3	2.14	0.48
1:B:441:LEU:HD12	1:B:451:GLN:HG2	1.96	0.48
1:A:230:LYS:O	1:A:234:ILE:HG12	2.14	0.48
1:B:417:GLN:OE1	1:B:468:GLU:HG3	2.14	0.47
1:A:424:ARG:O	1:A:428:VAL:HG23	2.15	0.47
1:A:421:LEU:HD22	1:A:444:LYS:HG2	1.96	0.47
1:B:111:SER:O	1:B:111:SER:OG	2.31	0.47
1:A:413:VAL:HG21	1:A:428:VAL:HG22	1.96	0.47
1:B:472:LYS:HA	1:B:472:LYS:HD3	1.63	0.47
1:B:394:GLU:HA	1:B:397:LEU:HB2	1.97	0.46
1:A:284:LEU:O	1:A:290:SER:OG	2.24	0.46
1:A:254:VAL:HA	1:A:257:LYS:HD2	1.98	0.46
1:A:311:LEU:HD11	1:B:422:GLU:HG3	1.97	0.45
1:B:99:LEU:HB2	1:B:102:THR:OG1	2.17	0.45
1:A:192:LEU:HD12	1:A:309:LEU:HD11	1.98	0.45
1:A:65:TYR:HB2	1:B:93:ASP:HB3	1.98	0.45
1:A:232:ARG:CZ	1:A:328:LEU:HB2	2.47	0.45
1:B:424:ARG:O	1:B:428:VAL:HG23	2.16	0.45
1:A:117:LEU:HA	1:B:117:LEU:HA	1.97	0.45
1:A:90:GLU:HB3	1:B:67:PRO:HB3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:LYS:NZ	1:A:116:ASP:OD1	2.49	0.45
1:A:60:LYS:HE3	1:B:123:GLU:HG3	1.98	0.45
1:A:221:PHE:HB3	1:A:226:VAL:HG22	1.99	0.45
1:B:117:LEU:HD13	1:B:125:LEU:HB2	1.99	0.44
1:B:367:VAL:HB	1:B:377:VAL:HB	1.99	0.44
1:B:115:TYR:OH	1:B:169:ARG:HD3	2.17	0.44
1:A:410:GLN:HG3	1:A:502:THR:HG21	1.99	0.44
1:A:288:LYS:HA	1:A:288:LYS:HD2	1.85	0.44
1:B:212:ASN:OD1	1:B:213:ASP:N	2.41	0.44
1:A:327:GLY:O	1:A:330:TYR:HB3	2.17	0.44
1:A:60:LYS:O	1:A:169:ARG:HD2	2.18	0.43
1:A:177:ASP:OD2	1:A:363:TYR:OH	2.30	0.43
1:B:330:TYR:CD2	1:B:366:LEU:HD22	2.44	0.43
1:A:198:ILE:O	1:A:202:LEU:HG	2.18	0.43
1:A:95:PRO:O	1:A:128:ARG:NH1	2.52	0.43
1:A:475:VAL:HG12	1:A:490:ARG:HA	2.00	0.43
1:B:193:LYS:HD3	1:B:312:PHE:HB3	2.01	0.43
1:B:328:LEU:HD12	1:B:328:LEU:HA	1.64	0.43
1:B:363:TYR:CD1	1:B:363:TYR:O	2.72	0.43
1:B:139:LEU:HD12	1:B:367:VAL:HG11	2.00	0.43
1:A:271:VAL:HG22	1:A:323:SER:HA	2.01	0.43
1:B:364:ASP:O	1:B:376:LYS:HG3	2.18	0.42
1:A:356:SER:O	1:A:388:ARG:NH1	2.52	0.42
1:A:286:ASP:O	1:A:290:SER:OG	2.37	0.42
1:B:80:ILE:HG21	1:B:174:CYS:SG	2.59	0.42
1:A:160:ASN:HA	1:A:161:PRO:HD3	1.79	0.42
1:A:412:LEU:HD12	1:A:439:GLU:O	2.20	0.42
1:A:441:LEU:HD23	1:A:441:LEU:HA	1.91	0.42
1:B:373:LYS:HG2	1:B:375:ARG:HB2	2.01	0.42
1:A:235:CYS:HA	1:A:238:VAL:HG22	2.02	0.41
1:B:92:ILE:HG12	1:B:149:ARG:NE	2.32	0.41
1:A:148:LYS:HG3	1:A:178:ILE:HG12	2.02	0.41
1:B:420:LEU:O	1:B:424:ARG:HG3	2.20	0.41
1:B:331:TYR:CD2	1:B:335:ILE:HG12	2.46	0.41
1:A:96:VAL:O	1:A:128:ARG:HG2	2.21	0.41
1:A:480:SER:HB2	1:A:501:ARG:HH22	1.86	0.41
1:B:112:LYS:HD3	1:B:112:LYS:HA	1.76	0.41
1:B:54:PHE:CG	1:B:55:VAL:N	2.88	0.41
1:A:139:LEU:HD22	1:A:144:LEU:HB2	2.03	0.40
1:A:302:LEU:HA	1:A:302:LEU:HD23	1.84	0.40
1:A:250:LYS:O	1:A:254:VAL:HG23	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:417:GLN:C	1:A:418:LYS:HD3	2.41	0.40
1:B:443:LYS:HD2	1:B:446:PRO:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	428/509 (84%)	395 (92%)	30 (7%)	3 (1%)	26	62
1	B	340/509 (67%)	313 (92%)	22 (6%)	5 (2%)	13	40
All	All	768/1018 (75%)	708 (92%)	52 (7%)	8 (1%)	19	52

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	184	PRO
1	A	253	MET
1	B	184	PRO
1	B	118	LYS
1	B	120	GLN
1	B	369	MET
1	B	385	GLY
1	A	372	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/438 (85%)	361 (97%)	10 (3%)	52	85
1	B	307/438 (70%)	302 (98%)	5 (2%)	70	93
All	All	678/876 (77%)	663 (98%)	15 (2%)	60	89

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	ARG
1	A	149	ARG
1	A	165	ARG
1	A	211	VAL
1	A	226	VAL
1	A	257	LYS
1	A	262	GLU
1	A	334	VAL
1	A	369	MET
1	A	411	VAL
1	B	58	THR
1	B	96	VAL
1	B	149	ARG
1	B	326	ARG
1	B	489	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	445	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	434/509 (85%)	0.00	11 (2%) 61 48	36, 65, 100, 125	0
1	B	350/509 (68%)	0.23	22 (6%) 23 14	39, 72, 111, 128	0
All	All	784/1018 (77%)	0.10	33 (4%) 40 28	36, 67, 105, 128	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	322	LEU	4.8
1	B	297	GLU	4.5
1	B	375	ARG	3.9
1	B	401	GLU	3.3
1	B	215	ARG	3.2
1	B	110	ASP	3.2
1	B	189	ALA	3.1
1	B	372	PRO	3.1
1	B	217	LEU	3.1
1	A	161	PRO	3.0
1	B	376	LYS	3.0
1	B	298	GLY	2.9
1	B	366	LEU	2.9
1	A	503	GLY	2.8
1	A	164	THR	2.8
1	B	373	LYS	2.8
1	A	256	GLU	2.7
1	A	372	PRO	2.7
1	A	252	GLU	2.5
1	B	361	GLY	2.5
1	B	340	LEU	2.5
1	A	399	ALA	2.5
1	B	400	LEU	2.5
1	B	402	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	373	LYS	2.4
1	B	496	GLU	2.4
1	B	191	CYS	2.4
1	A	100	LYS	2.3
1	B	323	SER	2.2
1	A	472	LYS	2.1
1	A	229	SER	2.1
1	B	341	LEU	2.1
1	B	330	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.