



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:30 PM GMT

PDB ID : 1X7I
Title : Crystal structure of the native copper homeostasis protein (cutCm) with calcium binding from Shigella flexneri 2a str. 301
Authors : Zhu, D.Y.; Zhu, Y.Q.; Huang, R.H.; Xiang, Y.; Wang, D.C.
Deposited on : 2004-08-14
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	<div> <div>4%</div> <div>71%</div> <div>20%</div> <div>8%</div> </div>
1	B	256	<div> <div>6%</div> <div>73%</div> <div>20%</div> <div>7%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4129 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper homeostasis protein cutC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	S	0	0	0
			1782	1117	319	330	16			
1	B	238	Total	C	N	O	S	0	0	0
			1799	1128	322	333	16			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	249	LEU	-	EXPRESSION TAG	UNP P67825
A	250	GLU	-	EXPRESSION TAG	UNP P67825
A	251	HIS	-	EXPRESSION TAG	UNP P67825
A	252	HIS	-	EXPRESSION TAG	UNP P67825
A	253	HIS	-	EXPRESSION TAG	UNP P67825
A	254	HIS	-	EXPRESSION TAG	UNP P67825
A	255	HIS	-	EXPRESSION TAG	UNP P67825
A	256	HIS	-	EXPRESSION TAG	UNP P67825
B	249	LEU	-	EXPRESSION TAG	UNP P67825
B	250	GLU	-	EXPRESSION TAG	UNP P67825
B	251	HIS	-	EXPRESSION TAG	UNP P67825
B	252	HIS	-	EXPRESSION TAG	UNP P67825
B	253	HIS	-	EXPRESSION TAG	UNP P67825
B	254	HIS	-	EXPRESSION TAG	UNP P67825
B	255	HIS	-	EXPRESSION TAG	UNP P67825
B	256	HIS	-	EXPRESSION TAG	UNP P67825


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	267	Total 267	O 267	0	0
3	B	280	Total 280	O 280	0	0

- Molecule 1: Copper homeostasis protein cutC



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	75.33Å 97.67Å 132.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.83 – 1.70 45.83 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.5 (45.83-1.70) 97.7 (45.83-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.73 (at 1.70Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.198 , 0.226 0.196 , 0.224	Depositor DCC
R_{free} test set	2665 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	11.0	Xtriage
Anisotropy	0.204	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 52833 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4129	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/1812	0.59	1/2450 (0.0%)
1	B	0.27	0/1829	0.58	0/2473
All	All	0.27	0/3641	0.59	1/4923 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	LEU	CA-CB-CG	5.63	128.25	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1782	0	1789	51	1
1	B	1799	0	1808	44	0
2	B	1	0	0	0	0
3	A	267	0	0	4	1
3	B	280	0	0	3	0
All	All	4129	0	3597	93	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 13.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:GLN:HE22	1:A:51:THR:H	1.08	0.96
1:A:179:VAL:HG11	1:A:195:VAL:HG21	1.50	0.94
1:B:18:GLN:HE22	1:B:51:THR:H	1.18	0.91
1:B:92:THR:HG22	1:B:93:GLY:H	1.38	0.88
1:A:179:VAL:HG11	1:A:195:VAL:CG2	2.07	0.84
1:A:40:LEU:HD11	1:A:44:LYS:HE3	1.62	0.82
1:A:3:LEU:HD13	1:A:23:ASP:HB2	1.65	0.79
1:A:207:MET:SD	1:B:66:CYS:SG	2.82	0.77
1:A:1:MET:HB3	1:A:246:GLN:HG2	1.66	0.76
1:B:203:GLN:HE21	1:B:227:ARG:HH11	1.40	0.69
1:B:80:ARG:O	1:B:84:GLU:HG3	1.93	0.69
1:A:3:LEU:CD1	1:A:23:ASP:HB2	2.22	0.68
1:A:135:ASN:HD21	1:A:167:HIS:CE1	2.11	0.68
1:B:103:MET:O	1:B:107:GLU:HG3	1.93	0.68
1:A:18:GLN:NE2	1:A:51:THR:H	1.88	0.67
1:B:58:ILE:O	1:B:92:THR:HG23	1.93	0.66
1:B:135:ASN:OD1	1:B:167:HIS:HE1	1.78	0.66
1:B:92:THR:HG22	1:B:93:GLY:N	2.11	0.64
1:A:40:LEU:HD11	1:A:44:LYS:CE	2.28	0.63
1:B:203:GLN:NE2	1:B:227:ARG:HH11	1.95	0.63
1:A:3:LEU:HD13	1:A:23:ASP:CB	2.28	0.62
1:A:15:LEU:O	1:A:19:GLN:HG3	2.00	0.62
1:A:80:ARG:HG3	3:A:475:HOH:O	2.01	0.60
1:B:239:LYS:O	1:B:243:GLU:HG3	2.00	0.60
1:B:18:GLN:NE2	1:B:51:THR:H	1.95	0.60
1:B:5:GLU:OE1	1:B:196:HIS:HD2	1.84	0.60
1:A:5:GLU:OE1	1:A:196:HIS:HD2	1.85	0.59
1:A:135:ASN:HD21	1:A:167:HIS:HE1	1.51	0.58
1:B:246:GLN:HE21	1:B:246:GLN:HA	1.69	0.57
1:B:246:GLN:NE2	1:B:249:LEU:HD12	2.19	0.57
1:A:18:GLN:HE21	1:A:52:ILE:H	1.53	0.57
1:A:176:GLY:O	1:A:177:ALA:HB3	2.05	0.57
1:A:122:ARG:HD2	1:A:125:ASP:OD2	2.04	0.57
1:A:40:LEU:HD13	1:A:40:LEU:O	2.05	0.56
1:B:205:SER:HB2	1:B:225:TYR:HB3	1.90	0.54
1:A:121:HIS:HD2	1:A:123:ALA:H	1.56	0.53
1:A:176:GLY:O	1:A:177:ALA:CB	2.57	0.53
1:A:205:SER:HB3	1:A:225:TYR:HB3	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:LEU:HD13	1:A:40:LEU:C	2.28	0.53
1:B:3:LEU:HD13	1:B:4:LEU:N	2.23	0.52
1:B:3:LEU:HB3	1:B:194:GLU:HG2	1.92	0.52
1:B:246:GLN:HA	1:B:246:GLN:NE2	2.24	0.52
1:A:76:LEU:HB3	1:A:80:ARG:NH2	2.25	0.51
1:A:155:LEU:C	1:A:155:LEU:HD13	2.30	0.51
1:B:83:ARG:HG3	1:B:116:LEU:HD13	1.93	0.51
1:A:27:LEU:HD11	1:A:46:VAL:HG21	1.94	0.49
1:A:76:LEU:O	1:A:80:ARG:HD3	2.13	0.49
1:B:176:GLY:O	1:B:177:ALA:CB	2.60	0.48
1:A:18:GLN:NE2	1:A:52:ILE:H	2.10	0.48
1:A:18:GLN:HE22	1:A:51:THR:N	1.92	0.48
1:B:203:GLN:HG3	1:B:227:ARG:HB2	1.96	0.48
1:B:176:GLY:O	1:B:177:ALA:HB3	2.13	0.48
1:A:3:LEU:HD12	1:A:4:LEU:N	2.29	0.48
1:A:168:ARG:HH21	1:A:168:ARG:HG3	1.78	0.47
1:B:1:MET:HB3	1:B:246:GLN:HB2	1.96	0.47
1:A:244:ARG:O	1:A:248:LYS:HG3	2.15	0.47
1:A:131:LEU:HD22	1:A:167:HIS:CD2	2.50	0.47
1:B:159:SER:O	1:B:163:GLU:HG3	2.15	0.46
1:A:3:LEU:HB3	1:A:194:GLU:HG2	1.96	0.46
1:A:13:CYS:SG	1:A:230:VAL:HB	2.55	0.46
1:B:15:LEU:O	1:B:19:GLN:HG3	2.16	0.46
1:B:18:GLN:HG3	1:B:52:ILE:HG23	1.98	0.46
1:B:203:GLN:CG	1:B:227:ARG:HB2	2.45	0.46
1:A:131:LEU:HD22	1:A:167:HIS:HD2	1.81	0.46
1:B:13:CYS:SG	1:B:230:VAL:HB	2.55	0.46
1:A:121:HIS:HE1	3:A:264:HOH:O	1.99	0.45
1:B:231:ASP:OD2	1:B:234:ALA:HB2	2.16	0.45
1:B:38:PRO:HA	3:B:1361:HOH:O	2.16	0.45
1:B:199:ALA:HA	3:B:1487:HOH:O	2.17	0.45
1:B:92:THR:CG2	1:B:93:GLY:H	2.20	0.44
1:A:125:ASP:HA	1:A:150:GLN:HG2	2.00	0.44
1:A:155:LEU:O	1:A:155:LEU:HD13	2.18	0.43
1:A:170:ALA:HB2	3:A:298:HOH:O	2.18	0.43
1:A:24:ARG:HD2	1:A:55:HIS:CE1	2.53	0.43
1:A:82:VAL:HG13	1:A:87:PHE:CD1	2.54	0.42
1:A:203:GLN:O	1:A:226:SER:HA	2.19	0.42
1:A:12:GLU:HG3	3:A:267:HOH:O	2.18	0.42
1:B:18:GLN:HE21	1:B:52:ILE:HG12	1.84	0.42
1:B:18:GLN:HE21	1:B:52:ILE:H	1.67	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:GLN:HE21	1:A:52:ILE:HG12	1.85	0.42
1:B:134:LEU:HD11	1:B:164:LEU:HD22	2.02	0.42
1:B:167:HIS:HD2	3:B:1401:HOH:O	2.02	0.42
1:A:210:ARG:HG2	1:A:225:TYR:CE2	2.56	0.41
1:B:205:SER:CB	1:B:225:TYR:HB3	2.51	0.41
1:A:48:GLN:HG3	1:A:49:ARG:HG3	2.03	0.41
1:A:18:GLN:HG3	1:A:52:ILE:HG23	2.02	0.41
1:B:203:GLN:NE2	1:B:227:ARG:NH1	2.67	0.41
1:B:1:MET:HB3	1:B:246:GLN:CB	2.51	0.41
1:A:36:LEU:HD12	1:B:30:ALA:HB2	2.02	0.41
1:B:58:ILE:HD11	1:B:82:VAL:HG21	2.02	0.41
1:B:246:GLN:HE22	1:B:249:LEU:HD12	1.84	0.40
1:B:170:ALA:HB1	1:B:171:PRO:CD	2.51	0.40
1:A:135:ASN:ND2	1:A:167:HIS:CE1	2.85	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:425:HOH:O	3:A:425:HOH:O[3_555]	1.68	0.52
1:A:203:GLN:NE2	1:A:203:GLN:NE2[3_655]	2.00	0.20

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/256 (91%)	224 (97%)	7 (3%)	1 (0%)	39	20
1	B	234/256 (91%)	223 (95%)	8 (3%)	3 (1%)	15	2
All	All	466/512 (91%)	447 (96%)	15 (3%)	4 (1%)	21	5

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	168	ARG
1	A	177	ALA
1	B	177	ALA
1	B	169	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/203 (91%)	179 (97%)	6 (3%)	46	24
1	B	187/203 (92%)	183 (98%)	4 (2%)	61	42
All	All	372/406 (92%)	362 (97%)	10 (3%)	52	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	27	LEU
1	A	84	GLU
1	A	184	LEU
1	A	227	ARG
1	A	246	GLN
1	B	121	HIS
1	B	146	LEU
1	B	184	LEU
1	B	239	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	48	GLN
1	A	121	HIS
1	A	135	ASN
1	A	136	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	150	GLN
1	A	167	HIS
1	A	185	HIS
1	A	196	HIS
1	B	18	GLN
1	B	48	GLN
1	B	55	HIS
1	B	167	HIS
1	B	196	HIS
1	B	203	GLN
1	B	246	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	236/256 (92%)	0.19	11 (4%) 35 39	5, 10, 26, 46	0
1	B	238/256 (92%)	0.30	15 (6%) 23 25	4, 10, 28, 51	0
All	All	474/512 (92%)	0.24	26 (5%) 29 31	4, 10, 27, 51	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	169	ASP	12.3
1	A	248	LYS	6.3
1	B	250	GLU	5.4
1	B	170	ALA	5.0
1	B	211	ASN	4.6
1	A	169	ASP	4.6
1	B	225	TYR	4.0
1	A	1	MET	4.0
1	A	225	TYR	3.8
1	B	202	TRP	3.8
1	B	249	LEU	3.4
1	B	203	GLN	3.2
1	B	228	TYR	3.2
1	A	223	ASP	3.1
1	A	246	GLN	3.0
1	B	1	MET	2.9
1	B	210	ARG	2.9
1	A	167	HIS	2.5
1	B	204	ALA	2.5
1	B	208	ARG	2.4
1	B	246	GLN	2.4
1	A	247	ALA	2.3
1	B	224	GLU	2.3
1	A	168	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	170	ALA	2.2
1	A	228	TYR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	B	1251	1/1	0.99	0.04	-6.42	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.