



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:28 PM GMT

PDB ID : 1X7W
Title : Crystal structure of the human mitochondrial branched-chain alpha-ketoacid dehydrogenase
Authors : Wynn, R.M.; Kato, M.; Machius, M.; Chuang, J.L.; Li, J.; Tomchick, D.R.; Chuang, D.T.
Deposited on : 2004-08-16
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2417 (1.76-1.72)
Clashscore	102246	2570 (1.76-1.72)
Ramachandran outliers	100387	2544 (1.76-1.72)
Sidechain outliers	100360	2544 (1.76-1.72)
RSRZ outliers	91569	2420 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	<div> <div>10%</div> <div>82%</div> <div>9%</div> <div>8%</div> </div>
2	B	342	<div> <div>6%</div> <div>87%</div> <div>9%</div> <div>.</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-oxoisovalerate dehydrogenase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	369	Total	C	N	O	S	0	9	0
			3025	1908	535	562	20			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	292	GLN	SER	ENGINEERED	UNP P12694

- Molecule 2 is a protein called 2-oxoisovalerate dehydrogenase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	329	Total	C	N	O	S	0	6	0
			2613	1679	433	486	15			

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	K	0	0
			1	1		
3	A	1	Total	K	0	0
			1	1		


- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mn	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is THIAMIN DIPHOSPHATE (three-letter code: TDP) (formula: $C_{12}H_{18}N_4O_7P_2S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	350	Total	O	0	0
			350	350		
8	B	267	Total	O	0	0
			267	267		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-oxoisovalerate dehydrogenase alpha subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	145.67Å 145.67Å 69.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.73 35.94 – 1.73	Depositor EDS
% Data completeness (in resolution range)	98.8 (50.00-1.73) 98.8 (35.94-1.73)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.147 , 0.181 0.182 , 0.208	Depositor DCC
R_{free} test set	1469 reflections (1.72%)	DCC
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 59.2	EDS
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 86997 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6291	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TDP, GOL, K, MN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	1/3098 (0.0%)	0.90	2/4186 (0.0%)
2	B	1.00	0/2679	0.90	5/3639 (0.1%)
All	All	0.97	1/5777 (0.0%)	0.90	7/7825 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	176	TYR	CE1-CZ	6.32	1.46	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	61	ARG	NE-CZ-NH2	-7.00	116.80	120.30
2	B	61	ARG	NE-CZ-NH1	6.30	123.45	120.30
1	A	252	ARG	NE-CZ-NH2	-5.95	117.32	120.30
2	B	274	ARG	NE-CZ-NH2	-5.61	117.50	120.30
2	B	181	ASP	CB-CG-OD2	5.48	123.23	118.30
2	B	314	ASP	CB-CG-OD2	5.39	123.15	118.30
1	A	242	ARG	NE-CZ-NH2	-5.11	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3025	0	2916	37	0
2	B	2613	0	2572	17	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
5	A	1	0	0	1	0
6	A	26	0	16	2	0
7	B	6	0	8	0	0
8	A	350	0	0	15	0
8	B	267	0	0	2	0
All	All	6291	0	5512	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:801:TDP:C2	6:A:801:TDP:H2	0.97	1.49
1:A:55:GLU:HG2	8:A:829:HOH:O	1.29	1.23
1:A:186[A]:VAL:CG2	1:A:273:VAL:HG23	1.71	1.21
1:A:186[A]:VAL:CG2	1:A:273:VAL:CG2	2.21	1.19
1:A:186[A]:VAL:HG23	1:A:273:VAL:CG2	1.76	1.16
1:A:185:ARG:HG2	8:A:1024:HOH:O	1.62	0.98
2:B:19:GLN:HG2	2:B:203:ILE:HD11	1.53	0.89
2:B:236:GLU:HG2	8:B:1106:HOH:O	1.77	0.83
1:A:186[A]:VAL:HG22	1:A:273:VAL:CG2	2.11	0.79
1:A:186[A]:VAL:HG21	1:A:273:VAL:HG23	1.65	0.78
2:B:289[C]:SER:OG	2:B:309:ARG:NH1	2.17	0.78
1:A:186[A]:VAL:HG23	1:A:273:VAL:HG21	1.65	0.77
1:A:186[A]:VAL:HG22	1:A:273:VAL:HG23	1.68	0.75
2:B:217:GLN:NE2	8:B:1097:HOH:O	2.17	0.74
1:A:186[A]:VAL:CG2	1:A:273:VAL:HG22	2.18	0.73
1:A:326:GLN:HA	8:A:1058:HOH:O	1.87	0.71
2:B:55[B]:ARG:HD2	2:B:195:ARG:HH22	1.54	0.71
1:A:313:ASP:HA	8:A:1095:HOH:O	1.91	0.69
5:A:704:CL:CL	6:A:801:TDP:H2	2.29	0.69
1:A:7:PRO:N	8:A:990:HOH:O	2.29	0.65
2:B:279:HIS:HE1	2:B:288:ALA:H	1.45	0.64
1:A:287:ARG:HD3	8:A:899:HOH:O	1.98	0.64
1:A:370[A]:GLU:OE2	8:A:857:HOH:O	0.61	0.61
2:B:279:HIS:CE1	2:B:288:ALA:H	2.19	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:ARG:HG2	8:A:1021:HOH:O	2.03	0.57
1:A:74:LEU:HD21	1:A:130[A]:MET:SD	2.46	0.56
1:A:149:HIS:HE1	1:A:162:SER:OG	1.88	0.56
2:B:19:GLN:CG	2:B:203:ILE:HD11	2.32	0.55
1:A:186[A]:VAL:HG22	1:A:273:VAL:HG22	1.83	0.55
2:B:222:VAL:HG11	2:B:336:LEU:HD11	1.93	0.51
1:A:112:GLN:HA	1:A:160:ILE:HD12	1.93	0.50
1:A:326:GLN:CA	8:A:1058:HOH:O	2.53	0.50
1:A:285:THR:HG23	8:A:977:HOH:O	2.10	0.50
1:A:147:PRO:O	1:A:149:HIS:HD2	1.96	0.48
1:A:130[B]:MET:SD	1:A:349:PHE:HA	2.55	0.46
1:A:134:TYR:CE2	1:A:353:GLU:HG2	2.51	0.46
2:B:23:LEU:HD23	2:B:194:TYR:CD1	2.52	0.45
2:B:142:GLY:O	2:B:146[A]:HIS:HB2	2.16	0.45
1:A:285:THR:CG2	8:A:977:HOH:O	2.65	0.44
2:B:209:PRO:HB2	2:B:212:GLN:HG3	1.99	0.44
2:B:164:ILE:HG12	2:B:188:PHE:HB2	1.99	0.44
1:A:358:PRO:HB3	1:A:362:LEU:HD12	1.99	0.44
1:A:275:GLU:OE1	1:A:277:GLN:NE2	2.45	0.43
1:A:376:ARG:NH1	8:A:1144:HOH:O	2.46	0.43
1:A:7:PRO:CA	8:A:990:HOH:O	2.67	0.42
2:B:55[A]:ARG:HG3	2:B:195:ARG:NH2	2.33	0.42
1:A:220:ARG:HB3	8:A:977:HOH:O	2.19	0.42
1:A:186[A]:VAL:HG23	1:A:273:VAL:HG22	1.80	0.41
2:B:55[B]:ARG:HH11	2:B:195:ARG:HH22	1.68	0.41
2:B:20:LYS:HG2	2:B:200:GLU:OE2	2.20	0.41
1:A:147:PRO:O	1:A:148:VAL:HG22	2.21	0.41
1:A:57:VAL:HG12	1:A:323:LEU:HD21	2.02	0.41
1:A:186[A]:VAL:HG12	1:A:187:VAL:N	2.36	0.40
1:A:146[B]:MET:HE2	8:A:819:HOH:O	2.21	0.40
1:A:170:GLN:O	2:B:85:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/400 (94%)	367 (98%)	5 (1%)	2 (0%)	34	14
2	B	334/342 (98%)	324 (97%)	9 (3%)	1 (0%)	46	26
All	All	708/742 (95%)	691 (98%)	14 (2%)	3 (0%)	39	20

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	ILE
2	B	196	ALA
1	A	148	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/336 (94%)	308 (98%)	7 (2%)	60	35
2	B	284/289 (98%)	276 (97%)	8 (3%)	51	25
All	All	599/625 (96%)	584 (98%)	15 (2%)	57	29

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	182	ASN
1	A	185	ARG
1	A	220	ARG
1	A	239	ILE
1	A	287	ARG
1	A	363	LEU
1	A	396	ASP
2	B	20	LYS
2	B	23	LEU
2	B	50	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	117	TYR
2	B	133	ARG
2	B	296	GLN
2	B	308[A]	SER
2	B	308[B]	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	HIS
1	A	276	ASN
1	A	333	GLN
2	B	279	HIS
2	B	296	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	TDP	A	801	4	21,27,27	1.82	6 (28%)	31,40,40	1.79	7 (22%)
7	GOL	B	901	-	5,5,5	0.38	0	5,5,5	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TDP	A	801	4	-	0/16/17/17	0/2/2/2
7	GOL	B	901	-	-	0/4/4/4	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	801	TDP	C4-N3	-2.69	1.37	1.39
6	A	801	TDP	C2A-C2'	2.17	1.56	1.49
6	A	801	TDP	C4A-C4	2.18	1.54	1.49
6	A	801	TDP	C2'-N3'	2.30	1.38	1.34
6	A	801	TDP	P2-O21	2.73	1.58	1.50
6	A	801	TDP	C4'-N3'	5.51	1.43	1.35

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	801	TDP	N1'-C2'-N3'	-3.04	119.97	125.60
6	A	801	TDP	C5'-C35-N3	-2.38	109.35	113.33
6	A	801	TDP	C5'-C6'-N1'	-2.36	119.76	123.86
6	A	801	TDP	C5-C4-N3	2.47	113.12	107.69
6	A	801	TDP	C5A-C5-C4	3.64	130.82	127.56
6	A	801	TDP	C2A-C2'-N1'	3.82	121.61	117.03
6	A	801	TDP	C6'-N1'-C2'	4.89	124.31	115.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	801	TDP	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	369/400 (92%)	0.40	40 (10%) 8 9	12, 18, 31, 46	0
2	B	329/342 (96%)	0.27	20 (6%) 25 28	14, 18, 28, 47	0
All	All	698/742 (94%)	0.34	60 (8%) 13 15	12, 18, 30, 47	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	28	ILE	7.6
1	A	8	GLN	6.7
1	A	387	GLN	5.0
1	A	395	LEU	4.9
1	A	183	ALA	4.4
1	A	397	HIS	4.3
1	A	9	PHE	4.2
1	A	10	PRO	4.2
2	B	301	LEU	4.1
1	A	7	PRO	3.9
1	A	12	ALA	3.6
1	A	13	SER	3.5
1	A	14	ALA	3.2
1	A	396	ASP	3.0
2	B	342	TYR	2.9
2	B	111	VAL	2.9
2	B	15	TYR	2.9
1	A	11	GLY	2.8
1	A	339	LYS	2.8
1	A	27	VAL	2.8
1	A	331	GLU	2.8
1	A	394	PRO	2.8
2	B	157	CYS	2.8
1	A	29	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	39	ARG	2.7
1	A	184	ASN	2.7
2	B	107	PHE	2.7
1	A	286	TYR	2.7
1	A	343	ARG	2.7
1	A	75	TYR	2.6
1	A	391	GLU	2.6
1	A	400	LYS	2.5
2	B	110	ILE	2.5
2	B	341	ASN	2.5
1	A	338	ARG	2.5
1	A	40	GLN	2.5
1	A	15	GLU	2.4
1	A	313	ASP	2.3
2	B	153	PHE	2.3
1	A	201	ALA	2.3
1	A	393	TYR	2.3
2	B	270	ILE	2.3
2	B	235	ARG	2.3
2	B	75	CYS	2.3
2	B	184	PRO	2.2
2	B	114	ALA	2.2
2	B	246	GLY	2.2
1	A	168	ILE	2.2
1	A	217	PHE	2.2
2	B	104	PHE	2.2
1	A	191	PHE	2.1
2	B	160	ILE	2.1
2	B	115	ALA	2.1
1	A	392	HIS	2.1
1	A	26	ASN	2.1
2	B	185	CYS	2.0
2	B	103	ILE	2.0
1	A	79	ARG	2.0
1	A	205	PHE	2.0
1	A	280	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	GOL	B	901	6/6	0.97	0.08	-0.36	17,18,19,19	0
4	MN	A	703	1/1	1.00	0.07	-0.88	14,14,14,14	0
3	K	B	702	1/1	0.99	0.10	-1.37	15,15,15,15	0
6	TDP	A	801	26/26	0.99	0.04	-2.47	7,9,10,12	0
3	K	A	701	1/1	1.00	0.06	-4.66	13,13,13,13	0
5	CL	A	704	1/1	0.97	0.05	-	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.