



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:23 PM GMT

PDB ID : 4XA2
Title : Structure of the Major Type IV pilin of *Acinetobacter baumannii*
Authors : Piepenbrink, K.H.; Sundberg, E.J.
Deposited on : 2014-12-12
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	8664 (2.00-1.96)
Clashscore	102246	9905 (2.00-1.96)
Ramachandran outliers	100387	9792 (2.00-1.96)
Sidechain outliers	100360	9791 (2.00-1.96)
RSRZ outliers	91569	8679 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	<div> <div>5%</div> <div>89%</div> <div>9%</div> <div>•</div> </div>
1	B	498	<div> <div>8%</div> <div>87%</div> <div>11%</div> <div>•</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8222 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Maltose-binding periplasmic protein,MBP-PilA: c.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	0	2	0
			3709	2356	618	725	10			
1	B	488	Total	C	N	O	S	0	3	0
			3697	2348	615	723	11			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P0AEY0
A	82	ALA	ASP	conflict	UNP P0AEY0
A	83	ALA	LYS	conflict	UNP P0AEY0
A	172	ALA	GLU	conflict	UNP P0AEY0
A	173	ALA	ASN	conflict	UNP P0AEY0
A	239	ALA	LYS	conflict	UNP P0AEY0
A	359	ALA	GLU	conflict	UNP P0AEY0
A	362	ALA	LYS	conflict	UNP P0AEY0
A	363	ALA	ASP	conflict	UNP P0AEY0
B	0	MET	-	initiating methionine	UNP P0AEY0
B	82	ALA	ASP	conflict	UNP P0AEY0
B	83	ALA	LYS	conflict	UNP P0AEY0
B	172	ALA	GLU	conflict	UNP P0AEY0
B	173	ALA	ASN	conflict	UNP P0AEY0
B	239	ALA	LYS	conflict	UNP P0AEY0
B	359	ALA	GLU	conflict	UNP P0AEY0
B	362	ALA	LYS	conflict	UNP P0AEY0
B	363	ALA	ASP	conflict	UNP P0AEY0

- Molecule 2 is MALTOTRIOSE (three-letter code: MLR) (formula: C₁₈H₃₂O₁₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			34	18	16		
2	B	1	Total	C	O	0	0
			34	18	16		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	396	Total	O	0	0
			396	396		
4	B	340	Total	O	0	0
			340	340		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein,MBP-PilA: c



- Molecule 1: Maltose-binding periplasmic protein,MBP-PilA: c



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.02Å 128.30Å 92.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.02 – 1.98 46.25 – 1.98	Depositor EDS
% Data completeness (in resolution range)	80.0 (41.02-1.98) 78.1 (46.25-1.98)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 1.98Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.181 , 0.241 0.186 , 0.242	Depositor DCC
R_{free} test set	2016 reflections (4.01%)	DCC
Wilson B-factor (Å ²)	18.3	Xtriage
Anisotropy	0.324	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.1	EDS
Estimated twinning fraction	0.036 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 53304 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8222	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLR, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3791	0.55	1/5152 (0.0%)
1	B	0.40	0/3782	0.53	0/5141
All	All	0.42	0/7573	0.54	1/10293 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	ARG	NE-CZ-NH2	-5.20	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3709	0	3679	34	0
1	B	3697	0	3666	41	0
2	A	34	0	32	1	0
2	B	34	0	32	0	0
3	A	8	0	12	0	0
3	B	4	0	6	0	0
4	A	396	0	0	15	1
4	B	340	0	0	10	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8222	0	7427	75	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:241:ASN:OD1	4:B:1301:HOH:O	1.93	0.84
1:A:18:ASN:OD1	4:A:1675:HOH:O	1.94	0.83
1:B:119:LYS:NZ	4:B:1302:HOH:O	2.15	0.78
1:B:32:GLY:O	4:B:1544:HOH:O	2.02	0.76
1:A:32:GLY:O	4:A:1301:HOH:O	2.06	0.74
1:A:291[A]:GLU:OE1	4:A:1590:HOH:O	2.09	0.70
1:A:29:LYS:N	4:A:1302:HOH:O	2.25	0.69
1:B:189:LYS:NZ	1:B:358:ASP:OD1	2.27	0.65
1:B:222:THR:HG22	1:B:224:MET:H	1.63	0.64
1:A:51:ALA:O	4:A:1553:HOH:O	2.15	0.63
1:A:80:THR:OG1	4:A:1689:HOH:O	2.16	0.63
1:A:1063:GLU:HG2	4:A:1602:HOH:O	2.01	0.61
1:B:25:LYS:HE2	1:B:29:LYS:HE2	1.83	0.61
1:A:28:GLU:C	4:A:1302:HOH:O	2.42	0.57
1:B:348:ILE:HG22	1:B:354[A]:ARG:NH2	2.19	0.57
1:B:348:ILE:HG22	1:B:354[A]:ARG:HH22	1.70	0.56
1:A:157:THR:HG23	4:A:1463:HOH:O	2.06	0.56
1:A:1134:LYS:O	1:A:1135:SER:HB2	2.05	0.56
1:B:1109:ASP:OD2	1:B:1122:ARG:NH1	2.39	0.55
1:B:285:LEU:HD23	1:B:290:LEU:HD21	1.87	0.55
1:A:297:LYS:NZ	4:A:1691:HOH:O	2.31	0.55
1:B:133:PRO:O	1:B:137:LYS:HE2	2.08	0.53
1:A:29:LYS:C	4:A:1302:HOH:O	2.47	0.52
1:A:91:PRO:HD2	4:A:1530:HOH:O	2.10	0.52
1:A:1044:LYS:HD3	1:A:1106:TRP:CD2	2.45	0.51
1:A:287:ASP:OD1	1:A:306:SER:OG	2.27	0.51
1:B:68:GLY:HA3	1:B:332:ASN:O	2.11	0.50
1:B:288:GLU:CD	1:B:288:GLU:H	2.15	0.50
1:B:245:THR:OG1	1:B:246:VAL:N	2.45	0.48
1:A:229:PRO:HA	1:A:232:TRP:CE2	2.48	0.48
1:B:129:TRP:CD1	1:B:248:PRO:HB2	2.48	0.48
1:B:205:ASN:OD1	1:B:206:ALA:N	2.45	0.48
1:A:311:LEU:HB3	1:A:317:ILE:HD13	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:LYS:HE2	4:A:1563:HOH:O	2.13	0.48
1:B:130:GLU:O	1:B:133:PRO:HD2	2.14	0.47
1:B:22:GLU:OE1	4:B:1617:HOH:O	2.20	0.47
1:A:352:SER:HB3	4:A:1497:HOH:O	2.15	0.47
1:A:181:VAL:HG12	1:A:183:VAL:HG22	1.96	0.47
1:A:46:LYS:HB2	1:A:46:LYS:HE3	1.78	0.47
1:A:47:PHE:CG	1:A:60:ILE:HD12	2.50	0.46
1:A:30:ASP:HB2	1:A:283:TYR:OH	2.15	0.46
1:A:79:ILE:HG13	1:A:79:ILE:O	2.15	0.46
1:B:137:LYS:HE3	1:B:137:LYS:HB2	1.77	0.46
1:B:25:LYS:CE	1:B:29:LYS:HE2	2.47	0.45
1:B:337:SER:HB3	4:B:1563:HOH:O	2.16	0.45
1:B:67:PHE:HB3	1:B:104:ILE:HD13	1.99	0.45
1:B:171:TYR:OH	1:B:174:GLY:HA2	2.17	0.45
1:B:209:ASP:HB2	4:B:1433:HOH:O	2.16	0.44
1:B:1056:THR:HB	4:B:1343:HOH:O	2.16	0.44
1:A:344[B]:ARG:HD3	2:A:1201:MLR:O6C	2.17	0.44
1:A:233:SER:HB2	1:A:297:LYS:HD3	1.99	0.44
1:B:31:THR:HB	1:B:33:ILE:HD12	1.99	0.44
1:B:288:GLU:OE1	1:B:288:GLU:N	2.51	0.43
1:B:139:LEU:HD21	1:B:223:ALA:HB2	2.00	0.43
1:A:270:SER:HA	1:A:271:PRO:HD3	1.76	0.43
1:B:295:LYS:HA	1:B:295:LYS:HD3	1.66	0.43
1:A:1:LYS:HG2	1:A:2:ILE:N	2.33	0.42
1:B:165:GLY:O	1:B:185:ASN:ND2	2.51	0.42
1:A:68:GLY:HA3	1:A:332:ASN:O	2.19	0.42
1:A:1132:ARG:NH2	1:A:1138:ALA:HB2	2.34	0.42
1:B:125:PRO:HG2	1:B:127:LYS:NZ	2.35	0.42
1:A:67:PHE:HB3	1:A:104:ILE:HD13	2.02	0.41
1:B:90:TYR:HA	1:B:91:PRO:HD3	1.95	0.41
1:B:1127:VAL:CG1	1:B:1131:LEU:HB2	2.50	0.41
1:A:151:LEU:HA	4:A:1463:HOH:O	2.20	0.41
1:B:246:VAL:HG23	4:B:1635:HOH:O	2.20	0.41
1:B:229:PRO:HA	1:B:232:TRP:CE2	2.56	0.41
1:B:179:LYS:HD3	1:B:179:LYS:HA	1.91	0.41
1:B:25:LYS:HD3	4:B:1617:HOH:O	2.21	0.41
1:B:133:PRO:HB3	1:B:203:HIS:CD2	2.56	0.41
1:A:30:ASP:OD1	1:A:30:ASP:N	2.53	0.41
1:B:254:PRO:HB3	4:B:1578:HOH:O	2.21	0.41
1:B:1099:ILE:HG22	1:B:1127:VAL:HG22	2.03	0.40
1:B:128:THR:HA	1:B:249:THR:HG22	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:PHE:O	1:A:283:TYR:HB2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1360:HOH:O	4:B:1370:HOH:O[1_655]	2.00	0.20
4:A:1334:HOH:O	4:A:1339:HOH:O[2_657]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	490/498 (98%)	482 (98%)	8 (2%)	0	100	100
1	B	489/498 (98%)	479 (98%)	10 (2%)	0	100	100
All	All	979/996 (98%)	961 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/387 (98%)	378 (100%)	1 (0%)	94	96

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	379/387 (98%)	377 (100%)	2 (0%)	92	93
All	All	758/774 (98%)	755 (100%)	3 (0%)	93	95

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	PHE
1	B	127	LYS
1	B	258	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1046	ASN
1	B	39	HIS
1	B	241	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLR	A	1201	-	36,36,36	1.28	3 (8%)	53,53,53	0.96	2 (3%)
3	EDO	A	1202	-	3,3,3	0.51	0	2,2,2	0.52	0
3	EDO	A	1203	-	3,3,3	0.45	0	2,2,2	0.41	0
2	MLR	B	1201	-	36,36,36	1.27	3 (8%)	53,53,53	0.98	2 (3%)
3	EDO	B	1202	-	3,3,3	0.44	0	2,2,2	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLR	A	1201	-	-	0/14/74/74	0/3/3/3
3	EDO	A	1202	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1203	-	-	0/1/1/1	0/0/0/0
2	MLR	B	1201	-	-	0/14/74/74	0/3/3/3
3	EDO	B	1202	-	-	0/1/1/1	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1201	MLR	O5B-C1B	2.39	1.47	1.41
2	B	1201	MLR	O5C-C1C	2.55	1.48	1.41
2	A	1201	MLR	O5C-C1C	2.62	1.48	1.41
2	A	1201	MLR	O5B-C1B	3.18	1.50	1.41
2	A	1201	MLR	O5A-C1A	3.18	1.49	1.43
2	B	1201	MLR	O5A-C1A	3.77	1.50	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1201	MLR	C1B-O4A-C4A	-3.10	109.76	118.00
2	A	1201	MLR	C1B-O4A-C4A	-2.64	110.97	118.00
2	B	1201	MLR	C1C-O4B-C4B	-2.55	111.21	118.00
2	A	1201	MLR	O4B-C1C-O5C	-2.07	105.30	110.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1201	MLR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/498 (98%)	0.58	23 (4%) 35 39	10, 21, 42, 51	0
1	B	488/498 (97%)	0.76	40 (8%) 14 17	9, 27, 47, 57	0
All	All	978/996 (98%)	0.67	63 (6%) 23 26	9, 24, 45, 57	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	315	PRO	6.1
1	B	240	VAL	5.6
1	B	125	PRO	4.7
1	B	241	ASN	4.4
1	A	80	THR	4.2
1	B	245	THR	4.0
1	B	239	ALA	3.9
1	B	193	THR	3.8
1	B	171	TYR	3.7
1	B	137	LYS	3.7
1	B	238	SER	3.6
1	B	350	ALA	3.4
1	A	29	LYS	3.3
1	B	242	TYR	3.2
1	A	1140	GLY	3.1
1	B	173	ALA	3.1
1	A	22	GLU	3.1
1	B	201	ASN	3.1
1	B	318	ALA	3.0
1	B	237	THR	3.0
1	B	124	ASN	3.0
1	B	314	ASP	3.0
1	A	71	ALA	3.0
1	B	117	TYR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	143	GLY	3.0
1	A	79	ILE	2.9
1	A	52	ALA	2.9
1	B	358	ASP	2.9
1	B	235	ILE	2.8
1	A	1141	SER	2.8
1	B	126	PRO	2.8
1	A	87	ASP	2.8
1	A	21	ALA	2.7
1	A	5	GLY	2.6
1	A	291[A]	GLU	2.6
1	B	354[A]	ARG	2.6
1	B	141	ALA	2.6
1	B	307	TYR	2.6
1	A	278	GLU	2.5
1	B	80	THR	2.5
1	B	306	SER	2.5
1	B	222	THR	2.4
1	A	28	GLU	2.4
1	B	134	ALA	2.4
1	B	246	VAL	2.4
1	A	82	ALA	2.4
1	B	324	ALA	2.4
1	B	257	PRO	2.4
1	A	306	SER	2.3
1	B	317	ILE	2.3
1	A	74	GLY	2.2
1	A	31	THR	2.2
1	B	145	SER	2.2
1	B	195	LEU	2.2
1	B	186	ALA	2.2
1	B	319	ALA	2.1
1	A	1135	SER	2.1
1	A	280	LEU	2.1
1	B	139	LEU	2.1
1	B	353	GLY	2.1
1	A	310	GLU	2.1
1	A	1134	LYS	2.1
1	A	35	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	A	1202	4/4	0.91	0.15	0.73	24,25,26,28	0
2	MLR	B	1201	34/34	0.95	0.14	0.27	14,18,24,28	0
2	MLR	A	1201	34/34	0.96	0.12	-0.73	10,14,19,22	0
3	EDO	A	1203	4/4	0.92	0.09	-1.98	37,38,41,49	0
3	EDO	B	1202	4/4	0.93	0.24	-	31,33,36,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.