



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 08:05 PM GMT

PDB ID : 4XBZ
Title : Crystal Structure of EvdO1 from *Micromonospora carbonacea* var. *aurantiaca*
Authors : McCulloch, K.M.; McCranie, E.K.; Sarwar, M.; Mathieu, J.L.; Gitschlag, B.L.;
Du, Y.; Bachmann, B.O.; Iverson, T.M.
Deposited on : 2014-12-17
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	
1	B	334	
1	C	334	
1	D	334	
1	E	334	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	334	<div><div>%</div><div><div></div><div>84%</div><div>8%</div><div>7%</div></div></div>
1	G	334	<div><div></div><div><div>76%</div><div>11%</div><div>13%</div></div></div>
1	H	334	<div><div>%</div><div><div></div><div>83%</div><div>10%</div><div>7%</div></div></div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 20394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EvdO1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	0	0	0
			2467	1560	442	454	11			
1	B	286	Total	C	N	O	S	0	0	0
			2270	1436	406	418	10			
1	C	317	Total	C	N	O	S	8	0	0
			2495	1573	451	460	11			
1	D	290	Total	C	N	O	S	0	0	0
			2320	1470	414	426	10			
1	E	283	Total	C	N	O	S	1	0	0
			2254	1428	405	411	10			
1	F	309	Total	C	N	O	S	2	0	0
			2449	1549	439	450	11			
1	G	292	Total	C	N	O	S	1	0	0
			2313	1462	414	427	10			
1	H	312	Total	C	N	O	S	0	0	0
			2449	1549	438	451	11			

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Ni	0	0
			1	1		
2	D	1	Total	Ni	0	0
			1	1		
2	E	1	Total	Ni	0	0
			1	1		
2	H	1	Total	Ni	0	0
			1	1		
2	B	1	Total	Ni	0	0
			1	1		
2	C	1	Total	Ni	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ni	0	0
			1	1		
2	F	1	Total	Ni	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	G	1	Total	C	O	0	0
			6	3	3		
3	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	163	Total	O	0	0
			163	163		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	134	Total 134	O 134	0	0
4	C	209	Total 209	O 209	0	0
4	D	165	Total 165	O 165	0	0
4	E	168	Total 168	O 168	0	0
4	F	178	Total 178	O 178	0	0
4	G	168	Total 168	O 168	0	0
4	H	148	Total 148	O 148	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: EvdO1

Chain A: 





• Molecule 1: EvdO1

Chain B: 





• Molecule 1: EvdO1

Chain C: 




• Molecule 1: EvdO1

Chain D: 









• Molecule 1: EvdO1





• Molecule 1: EvdO1



• Molecule 1: EvdO1



• Molecule 1: EvdO1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	91.61Å 109.43Å 146.05Å 90.00° 108.27° 90.00°	Depositor
Resolution (Å)	46.32 – 2.30 46.32 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (46.32-2.30) 99.4 (46.32-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.29Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.183 , 0.237 0.182 , 0.236	Depositor DCC
R_{free} test set	6080 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.608	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.8	EDS
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 120948 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	20394	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2534	0.55	0/3461
1	B	0.37	0/2327	0.51	0/3171
1	C	0.43	0/2564	0.55	0/3500
1	D	0.40	0/2379	0.54	0/3244
1	E	0.40	0/2311	0.54	0/3149
1	F	0.41	0/2514	0.53	0/3431
1	G	0.42	0/2370	0.54	0/3232
1	H	0.40	0/2514	0.54	0/3432
All	All	0.41	0/19513	0.54	0/26620

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2467	0	2399	36	0
1	B	2270	0	2193	20	0
1	C	2495	0	2395	30	0
1	D	2320	0	2250	25	0
1	E	2254	0	2189	33	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2449	0	2379	17	0
1	G	2313	0	2242	24	0
1	H	2449	0	2374	21	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
3	C	6	0	7	0	0
3	E	6	0	8	0	0
3	G	6	0	8	0	0
3	H	6	0	8	0	0
4	A	163	0	0	12	0
4	B	134	0	0	6	0
4	C	209	0	0	7	0
4	D	165	0	0	14	1
4	E	168	0	0	12	1
4	F	178	0	0	8	1
4	G	168	0	0	7	1
4	H	148	0	0	5	0
All	All	20394	0	18468	200	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 200 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:GLU:O	4:A:501:HOH:O	1.84	0.96
1:B:169:GLU:OE2	4:B:501:HOH:O	1.85	0.94
1:E:255:GLU:OE1	4:E:501:HOH:O	1.86	0.92
1:F:312:ARG:O	4:F:501:HOH:O	1.87	0.91
1:A:86:ARG:NH1	4:A:503:HOH:O	2.09	0.84

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:667:HOH:O	4:F:673:HOH:O[2_655]	1.92	0.28
4:D:655:HOH:O	4:G:668:HOH:O[1_545]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/334 (92%)	298 (97%)	10 (3%)	0	100	100
1	B	276/334 (83%)	263 (95%)	12 (4%)	1 (0%)	39	48
1	C	313/334 (94%)	301 (96%)	12 (4%)	0	100	100
1	D	280/334 (84%)	270 (96%)	10 (4%)	0	100	100
1	E	273/334 (82%)	264 (97%)	9 (3%)	0	100	100
1	F	301/334 (90%)	290 (96%)	10 (3%)	1 (0%)	46	57
1	G	284/334 (85%)	271 (95%)	11 (4%)	2 (1%)	26	31
1	H	306/334 (92%)	292 (95%)	14 (5%)	0	100	100
All	All	2341/2672 (88%)	2249 (96%)	88 (4%)	4 (0%)	52	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	66	ARG
1	G	198	GLU
1	G	199	TRP
1	F	-10	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/278 (93%)	251 (97%)	7 (3%)	52	70
1	B	236/278 (85%)	234 (99%)	2 (1%)	86	94
1	C	259/278 (93%)	255 (98%)	4 (2%)	72	85
1	D	243/278 (87%)	241 (99%)	2 (1%)	86	94
1	E	234/278 (84%)	231 (99%)	3 (1%)	76	87
1	F	256/278 (92%)	252 (98%)	4 (2%)	70	84
1	G	241/278 (87%)	238 (99%)	3 (1%)	78	89
1	H	255/278 (92%)	251 (98%)	4 (2%)	70	84
All	All	1982/2224 (89%)	1953 (98%)	29 (2%)	72	85

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	144	LEU
1	E	159	ASP
1	H	123	ASP
1	D	205	VAL
1	E	276	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 8 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	402	2	5,5,5	0.24	0	5,5,5	0.59	0
3	GOL	B	402	2	5,5,5	0.33	0	5,5,5	0.20	0
3	GOL	C	402	2	5,5,5	0.19	0	5,5,5	0.48	0
3	GOL	E	402	2	5,5,5	0.22	0	5,5,5	0.38	0
3	GOL	G	402	2	5,5,5	0.27	0	5,5,5	0.45	0
3	GOL	H	402	-	5,5,5	0.29	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	402	2	-	0/4/4/4	0/0/0/0
3	GOL	B	402	2	-	0/4/4/4	0/0/0/0
3	GOL	C	402	2	-	0/4/4/4	0/0/0/0
3	GOL	E	402	2	-	0/4/4/4	0/0/0/0
3	GOL	G	402	2	-	0/4/4/4	0/0/0/0
3	GOL	H	402	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/334 (93%)	-0.12	1 (0%) 94 96	23, 33, 50, 60	0
1	B	286/334 (85%)	-0.00	1 (0%) 94 96	24, 37, 58, 74	0
1	C	315/334 (94%)	-0.07	2 (0%) 90 93	19, 30, 50, 67	0
1	D	290/334 (86%)	-0.10	1 (0%) 94 96	20, 33, 52, 62	0
1	E	283/334 (84%)	-0.16	1 (0%) 93 95	20, 32, 49, 69	1 (0%)
1	F	309/334 (92%)	-0.02	3 (0%) 84 88	21, 34, 53, 70	1 (0%)
1	G	292/334 (87%)	-0.12	1 (0%) 94 96	20, 32, 50, 64	0
1	H	312/334 (93%)	0.00	2 (0%) 90 93	23, 35, 52, 62	0
All	All	2399/2672 (89%)	-0.07	12 (0%) 91 94	19, 33, 52, 74	2 (0%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	142	ALA	10.3
1	G	228	VAL	3.4
1	D	313	TRP	3.0
1	F	196	ALA	2.7
1	E	129	THR	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	402	6/6	0.70	0.14	0.63	36,39,41,43	0
3	GOL	H	402	6/6	0.71	0.15	0.19	41,48,50,56	0
3	GOL	E	402	6/6	0.86	0.13	-0.45	39,40,46,47	0
3	GOL	G	402	6/6	0.89	0.12	-0.62	31,37,41,42	0
3	GOL	C	402	6/6	0.85	0.12	-0.69	36,41,45,48	0
3	GOL	B	402	6/6	0.88	0.11	-1.22	35,39,41,43	0
2	NI	E	401	1/1	0.99	0.04	-	52,52,52,52	0
2	NI	F	401	1/1	0.98	0.04	-	56,56,56,56	0
2	NI	A	401	1/1	0.99	0.04	-	49,49,49,49	0
2	NI	D	401	1/1	0.97	0.07	-	64,64,64,64	0
2	NI	G	401	1/1	0.98	0.05	-	51,51,51,51	0
2	NI	H	401	1/1	0.98	0.07	-	55,55,55,55	0
2	NI	C	401	1/1	0.99	0.07	-	49,49,49,49	0
2	NI	B	401	1/1	0.97	0.06	-	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.