



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:28 PM GMT

PDB ID : 1XD8
Title : Crystal Structure of the Nitrogenase Fe protein Asp39Asn
Authors : Jang, S.B.; Jeong, M.S.; Seefeldt, L.C.; Peters, J.W.
Deposited on : 2004-09-05
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div>4%</div> <div>72%</div> <div>26%</div> <div>.</div> </div>
1	B	289	<div> <div>6%</div> <div>71%</div> <div>27%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitrogenase iron protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	289	Total	C	N	O	S	0	0	0
			2187	1364	370	432	21			
1	B	289	Total	C	N	O	S	0	0	0
			2187	1364	370	432	21			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASN	ASP	ENGINEERED	UNP P00459
B	39	ASN	ASP	ENGINEERED	UNP P00459

- Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nitrogenase iron protein 1



• Molecule 1: Nitrogenase iron protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.23Å 92.11Å 62.82Å 90.00° 101.05° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 19.70 – 2.70	Depositor EDS
% Data completeness (in resolution range)	88.0 (20.00-2.70) 92.8 (19.70-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.71Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.223 , 0.296 0.234 , 0.303	Depositor DCC
R_{free} test set	791 reflections (5.08%)	DCC
Wilson B-factor (Å ²)	62.3	Xtriage
Anisotropy	0.450	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 48.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 16355 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4382	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/2211	1.55	19/2977 (0.6%)
1	B	0.74	0/2211	1.55	19/2977 (0.6%)
All	All	0.74	0/4422	1.55	38/5954 (0.6%)

There are no bond length outliers.

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	ARG	NE-CZ-NH2	-20.73	109.94	120.30
1	B	100	ARG	NE-CZ-NH1	-20.67	109.96	120.30
1	B	100	ARG	NE-CZ-NH2	19.99	130.30	120.30
1	A	245	ARG	NE-CZ-NH1	19.58	130.09	120.30
1	A	245	ARG	NE-CZ-NH2	-17.62	111.49	120.30
1	A	100	ARG	NE-CZ-NH1	17.54	129.07	120.30
1	B	245	ARG	NE-CZ-NH2	16.87	128.73	120.30
1	B	245	ARG	NE-CZ-NH1	-16.27	112.16	120.30
1	B	100	ARG	CD-NE-CZ	12.43	141.00	123.60
1	A	245	ARG	CD-NE-CZ	11.87	140.21	123.60
1	B	245	ARG	CD-NE-CZ	10.38	138.13	123.60
1	A	46	ARG	NE-CZ-NH1	9.24	124.92	120.30
1	A	100	ARG	CD-NE-CZ	8.82	135.94	123.60
1	A	223	ARG	NE-CZ-NH1	8.63	124.61	120.30
1	B	46	ARG	NE-CZ-NH1	7.93	124.27	120.30
1	B	178	ARG	NE-CZ-NH1	6.90	123.75	120.30
1	A	287	GLU	N-CA-CB	-6.85	98.27	110.60
1	B	140	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	B	287	GLU	CA-CB-CG	6.25	127.15	113.40
1	B	287	GLU	N-CA-CB	-6.21	99.42	110.60
1	A	223	ARG	NE-CZ-NH2	-6.18	117.21	120.30
1	A	224	ARG	NE-CZ-NH1	5.90	123.25	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	140	ARG	NE-CZ-NH2	-5.89	117.35	120.30
1	A	124	TYR	CB-CG-CD2	-5.69	117.59	121.00
1	B	269	MET	CG-SD-CE	5.69	109.30	100.20
1	B	245	ARG	CA-CB-CG	5.65	125.82	113.40
1	B	223	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	287	GLU	CA-CB-CG	5.50	125.49	113.40
1	A	155	MET	CG-SD-CE	5.48	108.97	100.20
1	A	287	GLU	CB-CA-C	5.39	121.18	110.40
1	B	155	MET	CG-SD-CE	5.35	108.75	100.20
1	A	103	ILE	CB-CA-C	-5.30	100.99	111.60
1	A	122	VAL	CG1-CB-CG2	-5.27	102.47	110.90
1	B	224	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	B	287	GLU	CB-CA-C	5.10	120.61	110.40
1	B	286	ALA	CA-C-N	-5.08	106.02	117.20
1	A	286	ALA	CA-C-N	-5.07	106.05	117.20
1	B	191	ARG	NE-CZ-NH1	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2187	0	2200	37	0
1	B	2187	0	2200	39	0
2	A	8	0	0	1	0
All	All	4382	0	4400	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:290:SF4:FE1	1:B:97:CYS:HG	0.77	1.01

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:GLN:HE22	1:B:47:LEU:H	1.32	0.78
1:B:76:LEU:HD11	1:B:84:LYS:HB3	1.65	0.78
1:A:76:LEU:HD11	1:A:84:LYS:HB3	1.70	0.73
1:A:20:GLN:HE22	1:A:47:LEU:H	1.34	0.73
1:A:212:PRO:HG2	1:A:236:GLN:HE22	1.53	0.72
1:A:238:ASP:HA	1:A:241:ARG:HD2	1.72	0.71
1:B:212:PRO:HG2	1:B:236:GLN:HE22	1.58	0.69
1:B:162:ASN:HD21	1:B:259:ILE:H	1.45	0.62
1:A:5:CYS:SG	1:A:146:GLU:HB2	2.40	0.61
1:A:21:ASN:HD21	1:A:227:VAL:H	1.48	0.61
1:B:238:ASP:HA	1:B:241:ARG:HD2	1.82	0.61
1:A:64:ALA:HB1	1:A:69:ASP:HB2	1.83	0.61
1:B:209:HIS:HB2	1:B:243:LEU:HD13	1.82	0.60
1:B:5:CYS:SG	1:B:146:GLU:HB2	2.42	0.60
1:A:209:HIS:HB2	1:A:243:LEU:HD13	1.84	0.59
1:A:15:LYS:HE3	1:A:126:VAL:O	2.03	0.58
1:A:144:ALA:O	1:A:177:VAL:HG12	2.04	0.57
1:A:182:LEU:O	1:A:208:ILE:HG22	2.05	0.57
1:A:137:MET:HB3	1:A:138:PRO:HD3	1.85	0.57
1:A:95:VAL:HG23	1:B:171:TYR:OH	2.05	0.57
1:B:15:LYS:HE3	1:B:126:VAL:O	2.05	0.56
1:B:137:MET:HB3	1:B:138:PRO:HD3	1.86	0.56
1:A:162:ASN:HD21	1:A:259:ILE:H	1.52	0.56
1:A:173:ASN:H	1:A:173:ASN:ND2	2.04	0.55
1:B:268:LEU:HB3	1:B:274:MET:HB2	1.89	0.55
1:B:80:TYR:OH	1:B:228:ILE:HG22	2.07	0.55
1:B:193:ASP:O	1:B:197:ILE:HG13	2.07	0.54
1:A:193:ASP:O	1:A:197:ILE:HG13	2.08	0.54
1:B:217:VAL:O	1:B:221:GLU:HB2	2.08	0.54
1:B:64:ALA:HB1	1:B:69:ASP:HB2	1.88	0.54
1:B:21:ASN:HD21	1:B:227:VAL:H	1.53	0.54
1:B:33:VAL:HG22	1:B:121:PHE:HB2	1.90	0.53
1:B:144:ALA:O	1:B:177:VAL:HG12	2.09	0.53
1:B:3:ARG:HH12	1:B:248:VAL:HA	1.73	0.52
1:A:33:VAL:HG22	1:A:121:PHE:HB2	1.92	0.52
1:A:268:LEU:HB3	1:A:274:MET:HB2	1.92	0.52
1:B:182:LEU:O	1:B:208:ILE:HG22	2.10	0.51
1:A:3:ARG:HH12	1:A:248:VAL:HA	1.77	0.50
1:B:100:ARG:HG3	1:B:103:ILE:HD11	1.93	0.50
1:B:140:ARG:HD3	1:B:171:TYR:OH	2.11	0.50
1:B:186:SER:HB2	1:B:210:PHE:CZ	2.46	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:236:GLN:O	1:B:239:GLU:HB2	2.12	0.49
1:A:22:LEU:HD13	1:A:243:LEU:HG	1.96	0.48
1:B:187:ARG:HA	1:B:213:ARG:NH1	2.28	0.48
1:A:80:TYR:OH	1:A:228:ILE:HG22	2.15	0.46
1:A:252:LEU:O	1:A:252:LEU:HD13	2.15	0.46
1:A:171:TYR:OH	1:B:95:VAL:HG23	2.16	0.46
1:B:162:ASN:ND2	1:B:259:ILE:H	2.13	0.46
1:A:41:LYS:HD3	1:B:156:MET:HE1	1.98	0.45
1:A:78:ALA:HA	1:A:83:VAL:O	2.18	0.44
1:A:187:ARG:HA	1:A:213:ARG:NH1	2.33	0.44
1:B:173:ASN:ND2	1:B:173:ASN:H	2.16	0.43
1:B:162:ASN:HD21	1:B:259:ILE:N	2.15	0.43
1:A:217:VAL:O	1:A:221:GLU:HB2	2.19	0.43
1:B:214:ASP:HB2	1:B:236:GLN:NE2	2.33	0.43
1:A:60:MET:O	1:A:70:LEU:HD21	2.19	0.43
1:B:20:GLN:HE22	1:B:47:LEU:N	2.09	0.42
1:A:236:GLN:O	1:A:239:GLU:HB2	2.19	0.42
1:A:64:ALA:HB1	1:A:69:ASP:CB	2.49	0.42
1:B:244:ALA:O	1:B:248:VAL:HG23	2.20	0.42
1:B:78:ALA:HA	1:B:83:VAL:O	2.20	0.42
1:A:54:GLN:HG2	1:A:55:ASN:N	2.35	0.42
1:B:92:GLU:HB3	1:B:95:VAL:HG12	2.02	0.42
1:B:60:MET:O	1:B:70:LEU:HD21	2.20	0.42
1:A:186:SER:HB2	1:A:210:PHE:CZ	2.56	0.41
1:A:21:ASN:ND2	1:A:227:VAL:H	2.16	0.41
1:A:92:GLU:HB3	1:A:95:VAL:HG12	2.01	0.41
1:B:39:ASN:HA	1:B:40:PRO:HD3	1.83	0.41
1:A:140:ARG:HD3	1:A:171:TYR:OH	2.20	0.41
1:B:147:ILE:O	1:B:179:LEU:HD12	2.21	0.40
1:A:162:ASN:O	1:A:165:SER:HB2	2.21	0.40
1:A:147:ILE:HG21	1:A:168:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/289 (99%)	266 (93%)	19 (7%)	2 (1%)	26	55
1	B	287/289 (99%)	267 (93%)	18 (6%)	2 (1%)	26	55
All	All	574/578 (99%)	533 (93%)	37 (6%)	4 (1%)	26	55

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	80	TYR
1	A	80	TYR
1	B	190	ASP
1	A	190	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/233 (100%)	220 (94%)	13 (6%)	26	54
1	B	233/233 (100%)	220 (94%)	13 (6%)	26	54
All	All	466/466 (100%)	440 (94%)	26 (6%)	26	54

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	MET
1	A	52	LYS
1	A	56	THR
1	A	71	GLU
1	A	74	ASP
1	A	95	VAL
1	A	103	ILE
1	A	145	GLN
1	A	251	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	252	LEU
1	A	264	LEU
1	A	279	GLU
1	A	287	GLU
1	B	29	MET
1	B	52	LYS
1	B	56	THR
1	B	71	GLU
1	B	74	ASP
1	B	95	VAL
1	B	103	ILE
1	B	145	GLN
1	B	251	LYS
1	B	252	LEU
1	B	264	LEU
1	B	279	GLU
1	B	287	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	21	ASN
1	A	50	HIS
1	A	142	ASN
1	A	145	GLN
1	A	162	ASN
1	A	163	ASN
1	A	173	ASN
1	A	236	GLN
1	B	20	GLN
1	B	21	ASN
1	B	107	ASN
1	B	142	ASN
1	B	145	GLN
1	B	162	ASN
1	B	173	ASN
1	B	236	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SF4	A	290	1	0,12,12	0.00	-	0,24,24	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	A	290	1	-	0/0/48/48	4/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

All (4) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	290	SF4	FE1-FE2-S3-S4
2	A	290	SF4	FE3-FE4-S1-S2
2	A	290	SF4	FE2-FE4-S1-S3
2	A	290	SF4	FE1-FE3-S2-S4

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	290	SF4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/289 (100%)	0.14	11 (3%) 44 44	36, 75, 98, 100	0
1	B	289/289 (100%)	0.26	17 (5%) 26 24	41, 79, 99, 100	0
All	All	578/578 (100%)	0.20	28 (4%) 34 33	36, 77, 99, 100	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	4.4
1	A	289	VAL	3.6
1	B	233	LYS	3.4
1	A	286	ALA	3.2
1	B	74	ASP	3.1
1	A	287	GLU	3.1
1	B	116	GLU	3.0
1	B	226	THR	3.0
1	B	64	ALA	3.0
1	A	34	MET	2.8
1	A	116	GLU	2.8
1	B	65	GLY	2.7
1	B	121	PHE	2.6
1	A	1	ALA	2.5
1	A	35	ILE	2.5
1	B	63	GLU	2.5
1	B	52	LYS	2.3
1	A	23	VAL	2.3
1	B	62	ALA	2.2
1	B	190	ASP	2.1
1	A	190	ASP	2.1
1	A	32	LYS	2.1
1	B	227	VAL	2.1
1	B	118	ASP	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	115	TYR	2.0
1	B	30	GLY	2.0
1	A	226	THR	2.0
1	B	213	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SF4	A	290	8/8	0.99	0.08	-1.79	42,46,50,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.