



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 07:34 PM BST

PDB ID : 1XF7
Title : High Resolution NMR Structure of the Wilms' Tumor Suppressor Protein (WT1) Finger 3
Authors : Lachenmann, M.J.; Ladbury, J.E.; Dong, J.; Huang, K.; Carey, P.; Weiss, M.A.
Deposited on : 2004-09-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	29	

2 Ensemble composition and analysis ⓘ

This entry contains 40 models. Model 23 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:25 (24)	0.20	23

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters and 8 single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 7, 8, 10, 14, 23, 24, 25, 27, 33, 35, 36, 38
2	9, 17, 20, 21, 22, 32
3	4, 13, 16, 37
4	6, 15, 19
5	26, 34
6	11, 31
Single-model clusters	3; 12; 18; 28; 29; 30; 39; 40

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 489 atoms, of which 245 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Wilms' Tumor Protein.

Mol	Chain	Residues	Atoms						Trace
1	A	29	Total	C	H	N	O	S	0
			488	148	245	51	42	2	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	GLU	LYS	ENGINEERED	UNP P19544
A	29	LYS	THR	ENGINEERED	UNP P19544

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Wilms' Tumor Protein



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Wilms' Tumor Protein



4.2.2 Score per residue for model 2

- Molecule 1: Wilms' Tumor Protein



4.2.3 Score per residue for model 3

- Molecule 1: Wilms' Tumor Protein



4.2.4 Score per residue for model 4

- Molecule 1: Wilms' Tumor Protein




4.2.5 Score per residue for model 5

- Molecule 1: Wilms' Tumor Protein



4.2.6 Score per residue for model 6

- Molecule 1: Wilms' Tumor Protein



4.2.7 Score per residue for model 7

- Molecule 1: Wilms' Tumor Protein



4.2.8 Score per residue for model 8

- Molecule 1: Wilms' Tumor Protein




4.2.9 Score per residue for model 9

- Molecule 1: Wilms' Tumor Protein



4.2.10 Score per residue for model 10

- Molecule 1: Wilms' Tumor Protein



4.2.11 Score per residue for model 11

- Molecule 1: Wilms' Tumor Protein




4.2.12 Score per residue for model 12

- Molecule 1: Wilms' Tumor Protein




4.2.13 Score per residue for model 13

- Molecule 1: Wilms' Tumor Protein



4.2.14 Score per residue for model 14

- Molecule 1: Wilms' Tumor Protein




4.2.15 Score per residue for model 15

- Molecule 1: Wilms' Tumor Protein




4.2.16 Score per residue for model 16

- Molecule 1: Wilms' Tumor Protein



4.2.17 Score per residue for model 17

- Molecule 1: Wilms' Tumor Protein



4.2.18 Score per residue for model 18

- Molecule 1: Wilms' Tumor Protein



4.2.19 Score per residue for model 19

- Molecule 1: Wilms' Tumor Protein




4.2.20 Score per residue for model 20

- Molecule 1: Wilms' Tumor Protein



4.2.21 Score per residue for model 21

- Molecule 1: Wilms' Tumor Protein



4.2.22 Score per residue for model 22

- Molecule 1: Wilms' Tumor Protein



4.2.23 Score per residue for model 23 (medoid)

- Molecule 1: Wilms' Tumor Protein



4.2.24 Score per residue for model 24

- Molecule 1: Wilms' Tumor Protein



4.2.25 Score per residue for model 25

- Molecule 1: Wilms' Tumor Protein



4.2.26 Score per residue for model 26

- Molecule 1: Wilms' Tumor Protein



4.2.27 Score per residue for model 27

- Molecule 1: Wilms' Tumor Protein



4.2.28 Score per residue for model 28

- Molecule 1: Wilms' Tumor Protein




4.2.29 Score per residue for model 29

- Molecule 1: Wilms' Tumor Protein



4.2.30 Score per residue for model 30

- Molecule 1: Wilms' Tumor Protein



4.2.31 Score per residue for model 31

- Molecule 1: Wilms' Tumor Protein



4.2.32 Score per residue for model 32

- Molecule 1: Wilms' Tumor Protein




4.2.33 Score per residue for model 33

- Molecule 1: Wilms' Tumor Protein



4.2.34 Score per residue for model 34

- Molecule 1: Wilms' Tumor Protein



4.2.35 Score per residue for model 35

- Molecule 1: Wilms' Tumor Protein



4.2.36 Score per residue for model 36

- Molecule 1: Wilms' Tumor Protein



4.2.37 Score per residue for model 37

- Molecule 1: Wilms' Tumor Protein




4.2.38 Score per residue for model 38

- Molecule 1: Wilms' Tumor Protein



4.2.39 Score per residue for model 39

- Molecule 1: Wilms' Tumor Protein



4.2.40 Score per residue for model 40

- Molecule 1: Wilms' Tumor Protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DISTANCE GEOMETRY/SIMULATED ANNEALING FOLLOWED BY RESTRAINED MOLECULAR DYNAMICS*.

Of the 50 calculated structures, 40 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DGII	structure solution	Standalone
X-PLOR	refinement	3.1

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.25±0.01	0±0/211 (0.0±0.0%)	1.21±0.04	0±0/281 (0.1±0.2%)
All	All	1.25	0/8440 (0.0%)	1.21	9/11240 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.9±0.3
All	All	0	117

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	23	ARG	NE-CZ-NH2	-6.67	116.97	120.30	19	6
1	A	14	ARG	NE-CZ-NH2	-6.09	117.26	120.30	24	2
1	A	10	ARG	NE-CZ-NH2	-5.33	117.63	120.30	30	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	10	ARG	Sidechain	40
1	A	23	ARG	Sidechain	39
1	A	14	ARG	Sidechain	38

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	205	201	201	2±1
All	All	8240	8040	8040	87

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:12:PHE:CZ	1:A:21:HIS:CG	0.60	2.89	32	23
1:A:10:ARG:NH1	1:A:12:PHE:CE1	0.53	2.76	17	2
1:A:6:LYS:HD2	1:A:6:LYS:H	0.49	1.66	39	2
1:A:14:ARG:HD2	1:A:17:HIS:CE1	0.48	2.43	11	16
1:A:8:CYS:SG	1:A:10:ARG:HB2	0.48	2.47	29	3
1:A:14:ARG:CD	1:A:17:HIS:CE1	0.46	2.98	32	2
1:A:6:LYS:HD3	1:A:6:LYS:N	0.46	2.26	10	1
1:A:11:LYS:HB2	1:A:11:LYS:HZ2	0.46	1.70	12	1
1:A:3:PHE:CE1	1:A:15:SER:HB3	0.45	2.46	38	4
1:A:5:CYS:SG	1:A:7:THR:HB	0.45	2.52	40	7
1:A:8:CYS:O	1:A:9:GLN:HB2	0.44	2.12	39	11
1:A:7:THR:HG21	1:A:25:HIS:CG	0.44	2.48	19	4
1:A:6:LYS:H	1:A:6:LYS:HD3	0.43	1.72	15	1
1:A:6:LYS:CD	1:A:6:LYS:H	0.43	2.27	39	1
1:A:8:CYS:O	1:A:9:GLN:CB	0.43	2.67	16	1
1:A:11:LYS:H	1:A:11:LYS:CD	0.42	2.28	3	1
1:A:6:LYS:H	1:A:6:LYS:CD	0.42	2.26	16	1
1:A:20:THR:HA	1:A:23:ARG:CZ	0.42	2.44	13	1
1:A:11:LYS:H	1:A:11:LYS:HD2	0.41	1.74	3	1
1:A:4:GLN:HE21	1:A:11:LYS:NZ	0.41	2.13	8	1
1:A:21:HIS:CD2	1:A:22:THR:N	0.41	2.89	12	1
1:A:11:LYS:NZ	1:A:11:LYS:HB2	0.40	2.31	12	1
1:A:22:THR:C	1:A:24:THR:H	0.40	2.19	37	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	24/29 (83%)	19±1 (79±5%)	4±1 (16±6%)	1±1 (5±3%)	5	29
All	All	960/1160 (83%)	763 (79%)	153 (16%)	44 (5%)	5	29

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	9	GLN	30
1	A	23	ARG	9
1	A	22	THR	2
1	A	2	PRO	2
1	A	15	SER	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	24/28 (86%)	23±1 (94±4%)	1±1 (6±4%)	29	75
All	All	960/1120 (86%)	904 (94%)	56 (6%)	29	75

All 11 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	16	ASP	9
1	A	11	LYS	8
1	A	10	ARG	8
1	A	6	LYS	8
1	A	23	ARG	6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	19	LYS	6
1	A	14	ARG	4
1	A	2	PRO	3
1	A	9	GLN	2
1	A	25	HIS	1
1	A	22	THR	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided