



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 11:31 PM GMT

PDB ID : 1XFU
Title : Crystal structure of anthrax edema factor (EF) truncation mutant, EF-delta 64 in complex with calmodulin
Authors : Shen, Y.; Zhukovskaya, N.L.; Guo, Q.; Florian, J.; Tang, W.J.
Deposited on : 2004-09-15
Resolution : 3.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1005 (3.42-3.30)
Clashscore	102246	1076 (3.42-3.30)
Ramachandran outliers	100387	1059 (3.42-3.30)
Sidechain outliers	100360	1058 (3.42-3.30)
RSRZ outliers	91569	1010 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	747	<div> <div>6%</div> <div>24%</div> <div>58%</div> <div>14%</div> <div>..</div> </div>
1	B	747	<div> <div>6%</div> <div>24%</div> <div>58%</div> <div>15%</div> <div>..</div> </div>
1	C	747	<div> <div>7%</div> <div>24%</div> <div>58%</div> <div>14%</div> <div>..</div> </div>
1	D	747	<div> <div>6%</div> <div>24%</div> <div>58%</div> <div>15%</div> <div>..</div> </div>
1	E	747	<div> <div>5%</div> <div>24%</div> <div>59%</div> <div>14%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	747	<div><div></div><div>6%</div><div>24%</div><div>58%</div><div>15%</div><div></div><div></div></div>
2	O	149	<div><div></div><div>3%</div><div>19%</div><div>62%</div><div>15%</div><div></div><div></div></div>
2	P	149	<div><div></div><div>3%</div><div>21%</div><div>60%</div><div>14%</div><div></div><div></div></div>
2	Q	149	<div><div></div><div>4%</div><div>19%</div><div>62%</div><div>15%</div><div></div><div></div></div>
2	R	149	<div><div></div><div>3%</div><div>19%</div><div>62%</div><div>14%</div><div></div><div></div></div>
2	S	149	<div><div></div><div>2%</div><div>19%</div><div>62%</div><div>14%</div><div></div><div></div></div>
2	T	149	<div><div></div><div>%</div><div>21%</div><div>60%</div><div>14%</div><div></div><div></div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 42852 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calmodulin-sensitive adenylate cyclase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	B	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	C	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	D	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	E	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	F	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	MET	-	INITIATING METHIONINE	UNP P40136
A	55	HIS	-	EXPRESSION TAG	UNP P40136
A	56	HIS	-	EXPRESSION TAG	UNP P40136
A	57	HIS	-	EXPRESSION TAG	UNP P40136
A	58	HIS	-	EXPRESSION TAG	UNP P40136
A	59	HIS	-	EXPRESSION TAG	UNP P40136
A	60	HIS	-	EXPRESSION TAG	UNP P40136
A	61	ALA	-	CLONING ARTIFACT	UNP P40136
A	62	ALA	-	CLONING ARTIFACT	UNP P40136
A	63	ALA	-	CLONING ARTIFACT	UNP P40136
B	54	MET	-	INITIATING METHIONINE	UNP P40136
B	55	HIS	-	EXPRESSION TAG	UNP P40136
B	56	HIS	-	EXPRESSION TAG	UNP P40136
B	57	HIS	-	EXPRESSION TAG	UNP P40136
B	58	HIS	-	EXPRESSION TAG	UNP P40136
B	59	HIS	-	EXPRESSION TAG	UNP P40136
B	60	HIS	-	EXPRESSION TAG	UNP P40136

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	61	ALA	-	CLONING ARTIFACT	UNP P40136
B	62	ALA	-	CLONING ARTIFACT	UNP P40136
B	63	ALA	-	CLONING ARTIFACT	UNP P40136
C	54	MET	-	INITIATING METHIONINE	UNP P40136
C	55	HIS	-	EXPRESSION TAG	UNP P40136
C	56	HIS	-	EXPRESSION TAG	UNP P40136
C	57	HIS	-	EXPRESSION TAG	UNP P40136
C	58	HIS	-	EXPRESSION TAG	UNP P40136
C	59	HIS	-	EXPRESSION TAG	UNP P40136
C	60	HIS	-	EXPRESSION TAG	UNP P40136
C	61	ALA	-	CLONING ARTIFACT	UNP P40136
C	62	ALA	-	CLONING ARTIFACT	UNP P40136
C	63	ALA	-	CLONING ARTIFACT	UNP P40136
D	54	MET	-	INITIATING METHIONINE	UNP P40136
D	55	HIS	-	EXPRESSION TAG	UNP P40136
D	56	HIS	-	EXPRESSION TAG	UNP P40136
D	57	HIS	-	EXPRESSION TAG	UNP P40136
D	58	HIS	-	EXPRESSION TAG	UNP P40136
D	59	HIS	-	EXPRESSION TAG	UNP P40136
D	60	HIS	-	EXPRESSION TAG	UNP P40136
D	61	ALA	-	CLONING ARTIFACT	UNP P40136
D	62	ALA	-	CLONING ARTIFACT	UNP P40136
D	63	ALA	-	CLONING ARTIFACT	UNP P40136
E	54	MET	-	INITIATING METHIONINE	UNP P40136
E	55	HIS	-	EXPRESSION TAG	UNP P40136
E	56	HIS	-	EXPRESSION TAG	UNP P40136
E	57	HIS	-	EXPRESSION TAG	UNP P40136
E	58	HIS	-	EXPRESSION TAG	UNP P40136
E	59	HIS	-	EXPRESSION TAG	UNP P40136
E	60	HIS	-	EXPRESSION TAG	UNP P40136
E	61	ALA	-	CLONING ARTIFACT	UNP P40136
E	62	ALA	-	CLONING ARTIFACT	UNP P40136
E	63	ALA	-	CLONING ARTIFACT	UNP P40136
F	54	MET	-	INITIATING METHIONINE	UNP P40136
F	55	HIS	-	EXPRESSION TAG	UNP P40136
F	56	HIS	-	EXPRESSION TAG	UNP P40136
F	57	HIS	-	EXPRESSION TAG	UNP P40136
F	58	HIS	-	EXPRESSION TAG	UNP P40136
F	59	HIS	-	EXPRESSION TAG	UNP P40136
F	60	HIS	-	EXPRESSION TAG	UNP P40136
F	61	ALA	-	CLONING ARTIFACT	UNP P40136
F	62	ALA	-	CLONING ARTIFACT	UNP P40136

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	63	ALA	-	CLONING ARTIFACT	UNP P40136

- Molecule 2 is a protein called Calmodulin 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	146	Total	C	N	O	S	0	0	0
			1146	702	186	249	9			
2	P	146	Total	C	N	O	S	0	0	0
			1146	702	186	249	9			
2	Q	146	Total	C	N	O	S	0	0	0
			1146	702	186	249	9			
2	R	146	Total	C	N	O	S	0	0	0
			1146	702	186	249	9			
2	S	146	Total	C	N	O	S	0	0	0
			1146	702	186	249	9			
2	T	146	Total	C	N	O	S	0	0	0
			1146	702	186	249	9			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	F	1	Total	Mg	0	0
			1	1		


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	3	Total	Ca	0	0
			3	3		
4	Q	3	Total	Ca	0	0
			3	3		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	T	3	Total 3	Ca 3	0	0
4	O	3	Total 3	Ca 3	0	0
4	R	3	Total 3	Ca 3	0	0
4	S	3	Total 3	Ca 3	0	0






• Molecule 1: Calmodulin-sensitive adenylate cyclase




• Molecule 1: Calmodulin-sensitive adenylate cyclase










• Molecule 1: Calmodulin-sensitive adenylylate cyclase













• Molecule 2: Calmodulin 2



• Molecule 2: Calmodulin 2



• Molecule 2: Calmodulin 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	317.56Å 183.11Å 141.12Å 90.00° 90.05° 90.00°	Depositor
Resolution (Å)	29.99 – 3.35 40.47 – 3.30	Depositor EDS
% Data completeness (in resolution range)	95.8 (29.99-3.35) 94.8 (40.47-3.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 3.32Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.283 , 0.300 0.255 , 0.272	Depositor DCC
R_{free} test set	5597 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	100.0	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 42.4	EDS
Estimated twinning fraction	0.468 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.468 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.468 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.460 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.468 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 120891 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	42852	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/6104	0.79	7/8208 (0.1%)
1	B	0.53	0/6104	0.79	8/8208 (0.1%)
1	C	0.54	1/6104 (0.0%)	0.80	9/8208 (0.1%)
1	D	0.53	1/6104 (0.0%)	0.79	7/8208 (0.1%)
1	E	0.53	0/6104	0.79	7/8208 (0.1%)
1	F	0.54	1/6104 (0.0%)	0.79	7/8208 (0.1%)
2	O	0.56	0/1158	0.75	1/1553 (0.1%)
2	P	0.57	0/1158	0.75	1/1553 (0.1%)
2	Q	0.57	0/1158	0.75	1/1553 (0.1%)
2	R	0.57	0/1158	0.75	0/1553
2	S	0.57	0/1158	0.75	1/1553 (0.1%)
2	T	0.57	0/1158	0.75	1/1553 (0.1%)
All	All	0.54	3/43572 (0.0%)	0.79	50/58566 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	132	GLY	C-N	6.49	1.49	1.34
1	C	132	GLY	C-N	5.91	1.47	1.34
1	D	132	GLY	C-N	5.80	1.47	1.34

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	147	ARG	N-CA-C	6.70	129.09	111.00
1	F	147	ARG	N-CA-C	6.70	129.08	111.00
1	C	147	ARG	N-CA-C	6.69	129.05	111.00
1	A	147	ARG	N-CA-C	6.69	129.05	111.00
1	E	147	ARG	N-CA-C	6.68	129.04	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5992	0	6010	821	0
1	B	5992	0	6010	835	0
1	C	5992	0	6010	821	0
1	D	5992	0	6010	827	0
1	E	5992	0	6010	826	0
1	F	5992	0	6010	836	0
2	O	1146	0	1071	189	0
2	P	1146	0	1071	184	0
2	Q	1146	0	1071	187	0
2	R	1146	0	1071	193	0
2	S	1146	0	1071	193	0
2	T	1146	0	1071	191	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	O	3	0	0	0	0
4	P	3	0	0	0	0
4	Q	3	0	0	0	0
4	R	3	0	0	0	0
4	S	3	0	0	0	0
4	T	3	0	0	0	0
All	All	42852	0	42486	5969	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 70.

The worst 5 of 5969 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:ILE:HG13	1:A:171:TYR:HE1	1.02	1.19

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:165:GLN:HE21	1:E:251:PRO:HG2	1.07	1.17
1:E:154:ILE:HG13	1:E:171:TYR:HE1	1.01	1.15
1:F:154:ILE:HG13	1:F:171:TYR:HE1	1.03	1.14
1:B:165:GLN:HE21	1:B:251:PRO:HG2	1.06	1.14

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	733/747 (98%)	503 (69%)	145 (20%)	85 (12%)	0	3
1	B	733/747 (98%)	507 (69%)	142 (19%)	84 (12%)	0	3
1	C	733/747 (98%)	506 (69%)	145 (20%)	82 (11%)	0	3
1	D	733/747 (98%)	504 (69%)	144 (20%)	85 (12%)	0	3
1	E	733/747 (98%)	502 (68%)	146 (20%)	85 (12%)	0	3
1	F	733/747 (98%)	503 (69%)	146 (20%)	84 (12%)	0	3
2	O	144/149 (97%)	90 (62%)	39 (27%)	15 (10%)	1	4
2	P	144/149 (97%)	91 (63%)	39 (27%)	14 (10%)	1	5
2	Q	144/149 (97%)	90 (62%)	39 (27%)	15 (10%)	1	4
2	R	144/149 (97%)	90 (62%)	39 (27%)	15 (10%)	1	4
2	S	144/149 (97%)	92 (64%)	37 (26%)	15 (10%)	1	4
2	T	144/149 (97%)	92 (64%)	36 (25%)	16 (11%)	0	4
All	All	5262/5376 (98%)	3570 (68%)	1097 (21%)	595 (11%)	0	3

5 of 595 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	73	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	74	GLU
1	A	80	GLN
1	A	108	ASP
1	A	111	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	664/675 (98%)	581 (88%)	83 (12%)	6	24
1	B	664/675 (98%)	580 (87%)	84 (13%)	5	24
1	C	664/675 (98%)	577 (87%)	87 (13%)	5	22
1	D	664/675 (98%)	578 (87%)	86 (13%)	5	22
1	E	664/675 (98%)	580 (87%)	84 (13%)	5	24
1	F	664/675 (98%)	578 (87%)	86 (13%)	5	22
2	O	123/127 (97%)	104 (85%)	19 (15%)	3	16
2	P	123/127 (97%)	104 (85%)	19 (15%)	3	16
2	Q	123/127 (97%)	105 (85%)	18 (15%)	4	18
2	R	123/127 (97%)	104 (85%)	19 (15%)	3	16
2	S	123/127 (97%)	105 (85%)	18 (15%)	4	18
2	T	123/127 (97%)	105 (85%)	18 (15%)	4	18
All	All	4722/4812 (98%)	4101 (87%)	621 (13%)	5	22

5 of 621 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	284	LYS
1	E	115	LYS
2	R	55	VAL
1	D	377	GLN
1	D	507	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 218 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	377	GLN
1	E	239	HIS
2	R	107	HIS
1	D	480	ASN
1	D	655	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 24 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	735/747 (98%)	0.24	45 (6%) 25 25	24, 80, 134, 147	0
1	B	735/747 (98%)	0.25	45 (6%) 25 25	25, 80, 134, 148	0
1	C	735/747 (98%)	0.29	56 (7%) 17 17	25, 80, 134, 147	0
1	D	735/747 (98%)	0.23	48 (6%) 22 23	26, 80, 134, 147	0
1	E	735/747 (98%)	0.24	41 (5%) 28 27	26, 80, 135, 147	0
1	F	735/747 (98%)	0.24	44 (5%) 25 25	25, 81, 134, 147	0
2	O	146/149 (97%)	-0.03	4 (2%) 58 58	24, 65, 127, 137	0
2	P	146/149 (97%)	0.03	4 (2%) 58 58	24, 65, 126, 137	0
2	Q	146/149 (97%)	0.07	6 (4%) 41 40	23, 65, 126, 137	0
2	R	146/149 (97%)	-0.01	4 (2%) 58 58	24, 65, 126, 137	0
2	S	146/149 (97%)	-0.02	3 (2%) 67 67	24, 66, 126, 137	0
2	T	146/149 (97%)	-0.05	1 (0%) 89 89	23, 65, 126, 137	0
All	All	5286/5376 (98%)	0.21	301 (5%) 27 27	23, 77, 134, 148	0

The worst 5 of 301 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	225	ILE	13.6
1	A	230	ILE	11.6
1	F	230	ILE	10.4
1	E	226	ASP	10.3
1	E	225	ILE	9.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CA	R	707	1/1	0.93	0.22	0.47	79,79,79,79	0
4	CA	S	709	1/1	0.84	0.16	-0.04	78,78,78,78	0
4	CA	O	802	1/1	0.93	0.18	-0.18	44,44,44,44	0
4	CA	T	812	1/1	0.93	0.18	-0.26	43,43,43,43	0
4	CA	S	810	1/1	0.95	0.18	-0.27	45,45,45,45	0
4	CA	R	808	1/1	0.92	0.19	-0.32	41,41,41,41	0
4	CA	P	703	1/1	0.94	0.17	-0.36	71,71,71,71	0
4	CA	P	804	1/1	0.94	0.18	-0.57	43,43,43,43	0
4	CA	Q	806	1/1	0.88	0.17	-0.63	46,46,46,46	0
4	CA	T	811	1/1	0.97	0.16	-0.69	31,31,31,31	0
4	CA	T	711	1/1	0.94	0.14	-0.79	77,77,77,77	0
4	CA	O	701	1/1	0.97	0.12	-0.92	83,83,83,83	0
4	CA	O	801	1/1	0.96	0.15	-1.20	29,29,29,29	0
4	CA	Q	805	1/1	0.96	0.13	-1.23	32,32,32,32	0
4	CA	S	809	1/1	0.96	0.11	-1.46	25,25,25,25	0
4	CA	Q	705	1/1	0.94	0.07	-1.64	79,79,79,79	0
4	CA	P	803	1/1	0.97	0.12	-1.80	33,33,33,33	0
4	CA	R	807	1/1	0.98	0.12	-1.94	33,33,33,33	0
3	MG	B	901	1/1	0.93	0.16	-	13,13,13,13	0
3	MG	F	905	1/1	0.98	0.23	-	16,16,16,16	0
3	MG	E	904	1/1	0.97	0.16	-	14,14,14,14	0
3	MG	D	903	1/1	0.97	0.19	-	19,19,19,19	0
3	MG	C	902	1/1	0.90	0.13	-	14,14,14,14	0
3	MG	A	900	1/1	0.98	0.27	-	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.