



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:36 AM GMT

PDB ID : 2XKQ
Title : CRYSTAL STRUCTURE OF STREPTOCOCCUS SUIS DPR WITH MANGANESE
Authors : Haikarainen, T.; Thanassoulas, A.; Stavros, P.; Nounesis, G.; Haataja, S.; Papageorgiou, A.C.
Deposited on : 2010-07-12
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>8%</div> </div> </div>
1	B	165	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>7%</div> <div>8%</div> </div> </div>
1	C	165	<div> <div></div> <div> <div></div> <div>84%</div> <div>5%</div> <div>10%</div> </div> </div>
1	D	165	<div> <div></div> <div> <div></div> <div>88%</div> <div></div> <div>8%</div> </div> </div>
1	E	165	<div> <div>%</div> <div> <div></div> <div>89%</div> <div></div> <div>8%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	165	<div><div><div>%</div><div><div></div><div></div><div></div></div><div>89%</div><div><div></div><div></div><div></div></div><div>8%</div></div></div>
1	G	165	<div><div><div>%</div><div><div></div><div></div><div></div></div><div>84%</div><div><div></div><div></div><div></div></div><div>8%</div><div>8%</div></div></div>
1	H	165	<div><div><div></div><div></div><div></div></div><div>85%</div><div><div></div><div></div><div></div></div><div>6%</div><div>8%</div></div>
1	I	165	<div><div><div>%</div><div><div></div><div></div><div></div></div><div>87%</div><div><div></div><div></div><div></div></div><div>9%</div></div></div>
1	J	165	<div><div><div>%</div><div><div></div><div></div><div></div></div><div>87%</div><div><div></div><div></div><div></div></div><div>5%</div><div>8%</div></div></div>
1	K	165	<div><div><div></div><div></div><div></div></div><div>85%</div><div><div></div><div></div><div></div></div><div>7%</div><div>8%</div></div>
1	L	165	<div><div><div>%</div><div><div></div><div></div><div></div></div><div>82%</div><div><div></div><div></div><div></div></div><div>8%</div><div>8%</div></div></div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 15132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA PROTECTION DURING STARVATION PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	0	0
			1209	771	199	233	6			
1	B	151	Total	C	N	O	S	0	1	0
			1211	772	199	234	6			
1	C	149	Total	C	N	O	S	0	1	0
			1198	763	197	232	6			
1	D	152	Total	C	N	O	S	0	1	0
			1217	775	200	236	6			
1	E	151	Total	C	N	O	S	0	1	0
			1207	769	198	234	6			
1	F	151	Total	C	N	O	S	0	0	0
			1209	771	199	233	6			
1	G	151	Total	C	N	O	S	0	0	0
			1209	771	199	233	6			
1	H	151	Total	C	N	O	S	0	1	0
			1208	769	199	234	6			
1	I	150	Total	C	N	O	S	0	2	0
			1205	767	198	234	6			
1	J	152	Total	C	N	O	S	0	1	0
			1216	774	199	237	6			
1	K	152	Total	C	N	O	S	0	2	0
			1219	776	200	237	6			
1	L	151	Total	C	N	O	S	0	2	0
			1213	773	199	235	6			

- Molecule 2 is MANGANESE (III) ION (three-letter code: MN3) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Mn	0	0
			1	1		
2	J	1	Total	Mn	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Mn 1	0	0
2	K	1	Total 1	Mn 1	0	0
2	E	1	Total 1	Mn 1	0	0
2	H	1	Total 1	Mn 1	0	0
2	B	1	Total 1	Mn 1	0	0
2	I	1	Total 1	Mn 1	0	0
2	C	1	Total 1	Mn 1	0	0
2	A	1	Total 1	Mn 1	0	0
2	L	1	Total 1	Mn 1	0	0
2	F	1	Total 1	Mn 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	I	1	Total 1	Cl 1	0	0
3	A	1	Total 1	Cl 1	0	0
3	D	1	Total 1	Cl 1	0	0
3	C	1	Total 1	Cl 1	0	0

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	D	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	E	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	K	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total	Ca	0	0
			1	1		
5	B	2	Total	Ca	0	0
			2	2		
5	D	1	Total	Ca	0	0
			1	1		
5	F	2	Total	Ca	0	0
			2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	34	Total	O	0	0
			34	34		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	59	Total 59	O 59	0	0
6	C	29	Total 29	O 29	0	0
6	D	58	Total 58	O 58	0	0
6	E	31	Total 31	O 31	0	0
6	F	60	Total 60	O 60	0	0
6	G	30	Total 30	O 30	0	0
6	H	42	Total 42	O 42	0	0
6	I	43	Total 43	O 43	0	0
6	J	49	Total 49	O 49	0	0
6	K	54	Total 54	O 54	0	0
6	L	40	Total 40	O 40	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN





- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN








- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.22Å 139.47Å 143.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.00 – 2.40 44.24 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.00-2.40) 99.8 (44.24-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.4.0078	Depositor
R, R_{free}	0.183 , 0.245 0.207 , 0.265	Depositor DCC
R_{free} test set	4190 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.9	EDS
Estimated twinning fraction	0.024 for -h,l,k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	4 of 83898 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15132	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EPE, MN3, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/1234	0.64	0/1666
1	B	0.78	0/1240	0.69	0/1674
1	C	0.67	0/1227	0.62	0/1656
1	D	0.75	0/1246	0.65	0/1682
1	E	0.67	0/1236	0.62	0/1670
1	F	0.78	0/1234	0.66	0/1666
1	G	0.66	0/1234	0.66	0/1666
1	H	0.70	0/1237	0.65	0/1670
1	I	0.71	0/1238	0.67	0/1671
1	J	0.70	0/1245	0.62	0/1682
1	K	0.72	0/1252	0.66	0/1690
1	L	0.70	0/1246	0.68	0/1682
All	All	0.71	0/14869	0.65	0/20075

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1209	0	1174	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1211	0	1175	6	0
1	C	1198	0	1159	4	0
1	D	1217	0	1180	2	0
1	E	1207	0	1164	1	0
1	F	1209	0	1174	5	0
1	G	1209	0	1174	4	0
1	H	1208	0	1166	5	0
1	I	1205	0	1165	2	0
1	J	1216	0	1170	3	0
1	K	1219	0	1181	3	0
1	L	1213	0	1176	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	I	1	0	0	0	0
4	B	15	0	17	0	0
4	D	15	0	17	1	0
4	E	15	0	17	0	0
4	K	15	0	17	0	0
5	B	2	0	0	0	0
5	D	1	0	0	0	0
5	F	2	0	0	0	0
5	G	1	0	0	0	0
6	A	34	0	0	0	0
6	B	59	0	0	1	0
6	C	29	0	0	1	0
6	D	58	0	0	1	0
6	E	31	0	0	0	0
6	F	60	0	0	0	0
6	G	30	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	H	42	0	0	2	0
6	I	43	0	0	0	0
6	J	49	0	0	0	0
6	K	54	0	0	1	0
6	L	40	0	0	1	0
All	All	15132	0	14126	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:74:ASP:O	1:J:78:GLU:HG3	1.77	0.84
1:F:117:ARG:HH11	1:F:117:ARG:HG3	1.46	0.80
1:A:24:ASP:HB3	1:A:135:VAL:HG11	1.66	0.78
1:K:75:GLU:HG3	6:K:2021:HOH:O	1.93	0.69
1:G:78:GLU:OE1	1:L:157:LYS:NZ	2.33	0.61
1:C:24:ASP:HB2	1:C:139:GLU:OE2	2.01	0.61
1:G:59:HIS:HB3	1:G:60:PRO:HD3	1.85	0.57
1:B:130:GLN:NE2	1:B:152:LYS:HE2	2.20	0.57
1:L:46:VAL:HG13	1:L:114:GLN:HB3	1.88	0.56
1:C:47:HIS:CE1	1:C:59:HIS:CE1	2.95	0.55
1:K:59:HIS:HB3	1:K:60:PRO:HD3	1.89	0.55
1:F:90:THR:HG22	1:H:102:GLU:HB3	1.91	0.52
1:G:119:VAL:HG21	1:G:163:GLN:HE21	1.74	0.52
1:H:119:VAL:HG21	1:H:163:GLN:HE21	1.75	0.51
1:B:67:GLU:HB2	6:B:2019:HOH:O	2.10	0.51
1:A:109:VAL:HG12	1:A:114:GLN:HG2	1.91	0.50
1:B:99:GLN:HG3	1:B:124:TYR:OH	2.13	0.48
1:J:47:HIS:CE1	1:J:59:HIS:CE1	3.03	0.47
1:H:44:HIS:HD2	6:H:2004:HOH:O	1.96	0.47
1:L:138:GLU:HA	1:L:138:GLU:OE2	2.15	0.47
1:F:117:ARG:HH11	1:F:117:ARG:CG	2.20	0.47
1:C:60:PRO:O	1:C:63:ASP:HB2	2.15	0.46
1:K:78:GLU:HA	1:K:81:ILE:HD12	1.98	0.45
1:C:44:HIS:HD2	6:C:2002:HOH:O	2.00	0.45
1:A:60:PRO:O	1:A:63:ASP:HB2	2.16	0.45
1:B:47:HIS:CE1	1:B:59:HIS:CE1	3.05	0.45
1:G:155:ILE:HD13	1:G:155:ILE:HA	1.88	0.44
1:F:90:THR:HB	6:H:2032:HOH:O	2.16	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:VAL:HG21	1:A:163:GLN:HE21	1.83	0.44
1:H:152:LYS:O	1:H:156:GLU:HG3	2.18	0.43
1:B:119:VAL:HG21	1:B:163:GLN:HE21	1.82	0.43
1:E:110:THR:O	1:E:114:GLN:HG2	2.19	0.43
1:I:119:VAL:HG21	1:I:163:GLN:HE21	1.83	0.43
1:D:163:GLN:NE2	6:D:2054:HOH:O	2.38	0.43
1:F:117:ARG:NH1	1:F:117:ARG:HG3	2.26	0.43
1:L:130:GLN:NE2	1:L:152:LYS:HE2	2.34	0.43
1:J:78:GLU:HA	1:J:81:ILE:HD12	2.02	0.42
1:I:60:PRO:O	1:I:63:ASP:HB2	2.20	0.42
1:L:112:GLU:HG3	6:L:2029:HOH:O	2.20	0.41
1:H:68:GLU:OE2	1:H:154[B]:SER:OG	2.34	0.41
1:B:60:PRO:O	1:B:63:ASP:HB2	2.20	0.41
1:L:47:HIS:CE1	1:L:59:HIS:CE1	3.09	0.41
1:L:149:ASN:HA	1:L:149:ASN:HD22	1.77	0.41
1:D:57:ILE:HD12	4:D:1173:EPE:O1S	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/165 (90%)	147 (99%)	2 (1%)	0	100	100
1	B	150/165 (91%)	149 (99%)	1 (1%)	0	100	100
1	C	148/165 (90%)	146 (99%)	2 (1%)	0	100	100
1	D	151/165 (92%)	150 (99%)	1 (1%)	0	100	100
1	E	150/165 (91%)	148 (99%)	2 (1%)	0	100	100
1	F	149/165 (90%)	148 (99%)	1 (1%)	0	100	100
1	G	149/165 (90%)	147 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	150/165 (91%)	148 (99%)	2 (1%)	0	100	100
1	I	150/165 (91%)	149 (99%)	1 (1%)	0	100	100
1	J	151/165 (92%)	150 (99%)	1 (1%)	0	100	100
1	K	152/165 (92%)	151 (99%)	1 (1%)	0	100	100
1	L	151/165 (92%)	149 (99%)	2 (1%)	0	100	100
All	All	1800/1980 (91%)	1782 (99%)	18 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/141 (92%)	123 (95%)	6 (5%)	32	50
1	B	130/141 (92%)	125 (96%)	5 (4%)	40	60
1	C	129/141 (92%)	124 (96%)	5 (4%)	39	59
1	D	131/141 (93%)	126 (96%)	5 (4%)	40	60
1	E	129/141 (92%)	127 (98%)	2 (2%)	70	86
1	F	129/141 (92%)	126 (98%)	3 (2%)	58	78
1	G	129/141 (92%)	122 (95%)	7 (5%)	27	43
1	H	129/141 (92%)	125 (97%)	4 (3%)	47	69
1	I	130/141 (92%)	126 (97%)	4 (3%)	47	69
1	J	130/141 (92%)	127 (98%)	3 (2%)	58	78
1	K	132/141 (94%)	124 (94%)	8 (6%)	23	36
1	L	131/141 (93%)	124 (95%)	7 (5%)	28	44
All	All	1558/1692 (92%)	1499 (96%)	59 (4%)	40	60

All (59) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	HIS
1	A	73	LEU
1	A	101	LYS
1	A	131	LYS
1	A	146	ASP
1	A	160	TRP
1	B	44	HIS
1	B	63	ASP
1	B	99	GLN
1	B	101	LYS
1	B	111	ILE
1	C	24	ASP
1	C	44	HIS
1	C	56	MET
1	C	117	ARG
1	C	149	ASN
1	D	44	HIS
1	D	95	SER
1	D	101	LYS
1	D	131	LYS
1	D	146	ASP
1	E	44	HIS
1	E	112	GLU
1	F	44	HIS
1	F	101	LYS
1	F	117	ARG
1	G	44	HIS
1	G	92	LYS
1	G	101	LYS
1	G	146	ASP
1	G	149	ASN
1	G	154	SER
1	G	160	TRP
1	H	44	HIS
1	H	95	SER
1	H	101	LYS
1	H	138	GLU
1	I	44	HIS
1	I	63	ASP
1	I	117	ARG
1	I	160	TRP
1	J	44	HIS
1	J	117	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	J	146	ASP
1	K	44	HIS
1	K	67	GLU
1	K	95[A]	SER
1	K	95[B]	SER
1	K	101	LYS
1	K	112	GLU
1	K	146	ASP
1	K	149	ASN
1	L	22	LEU
1	L	44	HIS
1	L	60	PRO
1	L	131	LYS
1	L	135	VAL
1	L	138	GLU
1	L	149	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (44) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	ASN
1	A	149	ASN
1	A	163	GLN
1	B	44	HIS
1	B	130	GLN
1	B	149	ASN
1	B	163	GLN
1	C	44	HIS
1	C	130	GLN
1	C	149	ASN
1	C	163	GLN
1	D	130	GLN
1	D	149	ASN
1	E	130	GLN
1	E	149	ASN
1	E	163	GLN
1	F	130	GLN
1	F	149	ASN
1	F	163	GLN
1	G	149	ASN
1	G	163	GLN
1	H	44	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	99	GLN
1	H	108	ASN
1	H	130	GLN
1	H	149	ASN
1	H	163	GLN
1	I	44	HIS
1	I	130	GLN
1	I	149	ASN
1	I	163	GLN
1	J	149	ASN
1	J	163	GLN
1	K	44	HIS
1	K	99	GLN
1	K	108	ASN
1	K	130	GLN
1	K	149	ASN
1	K	163	GLN
1	L	44	HIS
1	L	99	GLN
1	L	130	GLN
1	L	149	ASN
1	L	163	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 26 ligands modelled in this entry, 22 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EPE	B	1173	-	14,15,15	0.49	0	18,20,20	1.77	4 (22%)
4	EPE	D	1173	-	14,15,15	0.55	0	18,20,20	2.07	7 (38%)
4	EPE	E	1173	-	14,15,15	0.53	0	18,20,20	2.52	5 (27%)
4	EPE	K	1173	-	14,15,15	0.43	0	18,20,20	2.53	6 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	B	1173	-	-	0/9/19/19	0/1/1/1
4	EPE	D	1173	-	-	0/9/19/19	0/1/1/1
4	EPE	E	1173	-	-	0/9/19/19	0/1/1/1
4	EPE	K	1173	-	-	0/9/19/19	0/1/1/1

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	K	1173	EPE	C9-N1-C6	-2.17	105.71	111.27
4	B	1173	EPE	O1S-S-C10	-2.13	105.08	106.91
4	E	1173	EPE	C7-N4-C5	2.04	116.49	111.27
4	E	1173	EPE	C7-N4-C3	2.34	117.26	111.27
4	B	1173	EPE	C7-N4-C5	2.49	117.65	111.27
4	D	1173	EPE	C5-N4-C3	2.54	114.39	108.90
4	D	1173	EPE	C3-C2-N1	2.54	115.18	110.63
4	D	1173	EPE	C7-N4-C3	2.69	118.17	111.27
4	D	1173	EPE	C7-N4-C5	2.69	118.18	111.27
4	D	1173	EPE	C5-C6-N1	2.94	115.89	110.63
4	B	1173	EPE	C7-N4-C3	3.09	119.20	111.27
4	E	1173	EPE	C6-N1-C2	3.10	115.62	108.90
4	K	1173	EPE	C7-N4-C5	3.14	119.31	111.27
4	K	1173	EPE	C7-N4-C3	3.28	119.68	111.27
4	D	1173	EPE	C6-N1-C2	3.83	117.20	108.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1173	EPE	O2S-S-C10	3.95	110.28	106.91
4	E	1173	EPE	C5-N4-C3	4.38	118.38	108.90
4	B	1173	EPE	C5-N4-C3	4.39	118.40	108.90
4	K	1173	EPE	O1S-S-C10	4.75	110.95	106.91
4	K	1173	EPE	O2S-S-C10	4.95	111.13	106.91
4	K	1173	EPE	C5-N4-C3	5.49	120.78	108.90
4	E	1173	EPE	O2S-S-C10	8.01	113.74	106.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1173	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/165 (91%)	0.07	1 (0%) 89 88	31, 41, 56, 66	0
1	B	151/165 (91%)	0.03	1 (0%) 89 88	25, 34, 48, 68	0
1	C	149/165 (90%)	-0.04	0 100 100	34, 42, 58, 73	0
1	D	152/165 (92%)	0.05	0 100 100	25, 33, 48, 60	0
1	E	151/165 (91%)	0.11	2 (1%) 79 79	34, 45, 61, 77	0
1	F	151/165 (91%)	0.05	2 (1%) 79 79	24, 33, 49, 56	0
1	G	151/165 (91%)	0.29	2 (1%) 79 79	33, 43, 60, 76	0
1	H	151/165 (91%)	0.06	0 100 100	31, 40, 57, 70	0
1	I	150/165 (90%)	0.11	2 (1%) 79 79	29, 40, 59, 79	0
1	J	152/165 (92%)	0.08	2 (1%) 79 79	25, 38, 55, 76	0
1	K	152/165 (92%)	0.04	0 100 100	28, 36, 49, 56	0
1	L	151/165 (91%)	0.04	1 (0%) 89 88	29, 38, 51, 63	0
All	All	1812/1980 (91%)	0.08	13 (0%) 89 88	24, 39, 57, 79	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	23	ALA	4.4
1	B	23	ALA	4.0
1	I	95[A]	SER	3.1
1	L	95[A]	SER	2.9
1	E	23	ALA	2.4
1	F	23	ALA	2.4
1	J	21	SER	2.3
1	E	91	LEU	2.3
1	I	96	GLU	2.2
1	A	22	LEU	2.2
1	F	96	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	22	LEU	2.1
1	J	96	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EPE	E	1173	15/15	0.96	0.22	1.10	17,27,33,33	0
4	EPE	D	1173	15/15	0.95	0.15	-1.17	17,23,28,28	0
3	CL	A	1174	1/1	0.98	0.10	-2.26	20,20,20,20	0
4	EPE	K	1173	15/15	0.96	0.14	-2.30	18,25,33,33	0
2	MN3	K	1174	1/1	0.99	0.06	-2.75	21,21,21,21	0
5	CA	G	1174	1/1	0.93	0.07	-3.40	26,26,26,26	0
2	MN3	B	1174	1/1	0.97	0.09	-3.40	14,14,14,14	0
2	MN3	J	1173	1/1	0.97	0.09	-3.70	17,17,17,17	0
2	MN3	F	1173	1/1	0.99	0.11	-3.81	13,13,13,13	0
4	EPE	B	1173	15/15	0.97	0.12	-3.94	16,26,34,35	0
3	CL	D	1176	1/1	0.96	0.11	-4.67	15,15,15,15	0
2	MN3	C	1173	1/1	0.98	0.05	-4.86	30,30,30,30	0
2	MN3	H	1173	1/1	0.98	0.07	-6.01	24,24,24,24	0
2	MN3	A	1173	1/1	0.98	0.04	-7.18	20,20,20,20	0
2	MN3	G	1173	1/1	0.93	0.04	-7.27	34,34,34,34	0
2	MN3	E	1174	1/1	0.98	0.03	-7.30	29,29,29,29	0
2	MN3	L	1173	1/1	0.96	0.04	-7.42	23,23,23,23	0
2	MN3	D	1174	1/1	0.97	0.06	-7.62	20,20,20,20	0
2	MN3	I	1173	1/1	0.97	0.04	-9.65	26,26,26,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	CA	F	1174	1/1	1.00	0.04	-13.03	13,13,13,13	0
5	CA	D	1175	1/1	0.84	0.15	-	29,29,29,29	0
3	CL	C	1174	1/1	0.91	0.08	-	33,33,33,33	0
5	CA	B	1175	1/1	0.91	0.07	-	42,42,42,42	0
3	CL	I	1174	1/1	0.90	0.14	-	25,25,25,25	0
5	CA	B	1176	1/1	0.81	0.08	-	58,58,58,58	0
5	CA	F	1175	1/1	0.94	0.06	-	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.