



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:35 AM GMT

PDB ID : 2XMR
Title : CRYSTAL STRUCTURE OF HUMAN NDRG2 PROTEIN PROVIDES INSIGHT INTO ITS ROLE AS A TUMOR SUPPRESSOR
Authors : Hwang, J.; Kim, Y.; Lee, H.; Kim, M.H.
Deposited on : 2010-07-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	<div> <div>11%</div> <div>92%</div> <div>6%</div> </div>
1	B	281	<div> <div>14%</div> <div>94%</div> <div>6%</div> </div>
1	C	281	<div> <div>14%</div> <div>94%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	1306	-	-	X	-
3	ACT	B	1305	-	-	X	-
3	ACT	C	1305	-	-	X	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7032 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN NDRG2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2197	1405	363	416	13			
1	B	281	Total	C	N	O	S	0	0	0
			2197	1405	363	416	13			
1	C	281	Total	C	N	O	S	0	1	0
			2208	1411	367	417	13			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	ALA	LYS	ENGINEERED MUTATION	UNP Q9UN36
A	47	ALA	LYS	ENGINEERED MUTATION	UNP Q9UN36
B	45	ALA	LYS	ENGINEERED MUTATION	UNP Q9UN36
B	47	ALA	LYS	ENGINEERED MUTATION	UNP Q9UN36
C	45	ALA	LYS	ENGINEERED MUTATION	UNP Q9UN36
C	47	ALA	LYS	ENGINEERED MUTATION	UNP Q9UN36

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	132	Total	O	0	0
			132	132		
5	B	152	Total	O	0	0
			152	152		
5	C	115	Total	O	0	0
			115	115		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PROTEIN NDRG2



• Molecule 1: PROTEIN NDRG2



• Molecule 1: PROTEIN NDRG2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.23Å 88.05Å 126.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 29.93 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.5 (30.00-2.00) 96.5 (29.93-2.00)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.180 , 0.232 0.195 , 0.247	Depositor DCC
R_{free} test set	3217 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	25.8	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.9	EDS
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 63288 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7032	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	1/2255 (0.0%)	0.67	2/3072 (0.1%)
1	B	0.57	0/2255	0.64	0/3072
1	C	0.53	0/2266	0.61	0/3086
All	All	0.57	1/6776 (0.0%)	0.64	2/9230 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	218	LEU	CG-CD1	-7.71	1.23	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	LEU	CB-CG-CD1	6.23	121.60	111.00
1	A	224	ASN	N-CA-C	-5.49	96.17	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	223	TYR	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	224	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2197	0	2131	26	0
1	B	2197	0	2131	16	0
1	C	2208	0	2143	17	0
2	A	1	0	0	0	0
3	A	4	0	3	4	0
3	B	4	0	3	2	0
3	C	4	0	3	5	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
4	C	6	0	8	0	0
5	A	132	0	0	2	0
5	B	152	0	0	0	0
5	C	115	0	0	3	0
All	All	7032	0	6438	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:83:ARG:HE	3:B:1305:ACT:H1	1.00	1.09
1:A:83:ARG:HE	3:A:1306:ACT:H1	1.18	1.01
1:C:85:HIS:NE2	3:C:1305:ACT:H1	1.80	0.97
1:A:25:LYS:HZ2	1:A:26:THR:H	1.13	0.94
1:A:282:GLN:H	1:A:282:GLN:HE21	1.16	0.93
1:A:119:GLN:HE22	1:B:298:GLN:HE22	1.10	0.93
1:B:83:ARG:NE	3:B:1305:ACT:H1	1.86	0.90
1:C:282:GLN:H	1:C:282:GLN:HE21	1.20	0.84
1:B:282:GLN:H	1:B:282:GLN:HE21	1.33	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:83:ARG:HE	3:C:1305:ACT:CH3	1.99	0.75
1:A:240:ARG:HG2	1:A:240:ARG:HH11	1.53	0.73
1:A:119:GLN:NE2	1:B:298:GLN:HE22	1.86	0.73
1:A:83:ARG:NE	3:A:1306:ACT:H1	2.01	0.71
1:A:282:GLN:NE2	1:A:282:GLN:H	1.91	0.66
1:C:282:GLN:H	1:C:282:GLN:NE2	1.92	0.66
1:C:83:ARG:HE	3:C:1305:ACT:H2	1.61	0.66
1:A:240:ARG:CG	1:A:240:ARG:HH11	2.09	0.65
1:A:85:HIS:NE2	3:A:1306:ACT:H2	2.15	0.61
1:C:37:THR:HG22	5:C:2025:HOH:O	2.04	0.57
1:C:83:ARG:HE	3:C:1305:ACT:H3	1.69	0.57
1:A:85:HIS:NE2	3:A:1306:ACT:CH3	2.68	0.56
1:A:282:GLN:N	1:A:282:GLN:HE21	1.96	0.56
1:B:282:GLN:H	1:B:282:GLN:NE2	2.00	0.56
1:C:25:LYS:HG3	1:C:27:HIS:CE1	2.41	0.56
1:B:300:MET:HE2	1:B:302:TYR:CE2	2.42	0.55
1:B:300:MET:HE2	1:B:302:TYR:HE2	1.72	0.55
1:A:110:LEU:HD23	1:A:113:MET:HE1	1.88	0.54
1:A:25:LYS:HE3	5:C:2036:HOH:O	2.08	0.52
1:C:25:LYS:HG3	1:C:27:HIS:HE1	1.75	0.51
1:A:119:GLN:HE22	1:B:298:GLN:NE2	1.92	0.50
1:C:282:GLN:HE21	1:C:282:GLN:N	2.00	0.49
1:A:240:ARG:NH1	1:A:240:ARG:CG	2.72	0.47
1:B:300:MET:CE	1:B:302:TYR:HE2	2.26	0.47
1:A:25:LYS:NZ	1:A:26:THR:H	1.97	0.46
1:A:110:LEU:HD23	1:A:113:MET:CE	2.47	0.45
1:A:242:PRO:HG2	1:A:300:MET:HG2	1.99	0.45
1:C:230:ASN:HA	1:C:230:ASN:HD22	1.67	0.44
1:B:71:PHE:CE2	1:B:73:ASP:HB2	2.53	0.44
1:A:115:PRO:HB2	1:B:303:MET:CE	2.48	0.44
1:B:277:SER:HB3	1:B:285:GLN:HE21	1.83	0.43
1:C:277:SER:HB3	1:C:285:GLN:OE1	2.19	0.43
1:A:25:LYS:HA	1:A:25:LYS:HZ3	1.84	0.43
1:B:74:MET:O	1:B:78:ILE:HG12	2.18	0.42
1:A:25:LYS:HB3	1:A:27:HIS:CE1	2.53	0.42
1:B:265:ASP:HA	1:B:266:PRO:HD2	1.86	0.42
1:B:295:TYR:HB2	1:C:172:LEU:HG	2.00	0.42
1:A:181:GLU:HG2	5:A:2088:HOH:O	2.19	0.42
1:C:27:HIS:HD2	1:C:120:TYR:OH	2.03	0.41
1:C:41:TYR:HB2	3:C:1305:ACT:H3	2.02	0.41
1:A:115:PRO:O	1:A:119:GLN:HG3	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:277:SER:CB	1:A:285:GLN:HG3	2.51	0.41
1:C:247:VAL:HG21	1:C:254:GLU:HA	2.02	0.41
1:C:204:ARG:HD2	5:C:2077:HOH:O	2.19	0.41
1:A:147:ASP:CG	5:A:2062:HOH:O	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/281 (99%)	276 (99%)	3 (1%)	0	100	100
1	B	279/281 (99%)	278 (100%)	1 (0%)	0	100	100
1	C	280/281 (100%)	280 (100%)	0	0	100	100
All	All	838/843 (99%)	834 (100%)	4 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/238 (100%)	229 (96%)	9 (4%)	40	36
1	B	238/238 (100%)	235 (99%)	3 (1%)	76	79
1	C	239/238 (100%)	235 (98%)	4 (2%)	68	71

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	715/714 (100%)	699 (98%)	16 (2%)	60	62

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LYS
1	A	43	THR
1	A	100	LEU
1	A	147	ASP
1	A	205	ASN
1	A	218	LEU
1	A	224	ASN
1	A	240	ARG
1	A	282	GLN
1	B	35	SER
1	B	103	GLN
1	B	282	GLN
1	C	198	GLU
1	C	230	ASN
1	C	259	GLU
1	C	282	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	HIS
1	A	75	GLN
1	A	119	GLN
1	A	224	ASN
1	A	280	GLN
1	A	282	GLN
1	A	298	GLN
1	B	27	HIS
1	B	75	GLN
1	B	103	GLN
1	B	230	ASN
1	B	280	GLN
1	B	282	GLN
1	B	285	GLN
1	C	27	HIS
1	C	75	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	103	GLN
1	C	224	ASN
1	C	230	ASN
1	C	280	GLN
1	C	282	GLN
1	C	298	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	A	1306	-	1,3,3	1.49	0	0,3,3	0.00	-
4	GOL	A	1307	-	5,5,5	0.28	0	5,5,5	0.53	0
3	ACT	B	1305	-	1,3,3	0.68	0	0,3,3	0.00	-
4	GOL	B	1306	-	5,5,5	0.33	0	5,5,5	0.22	0
3	ACT	C	1305	-	1,3,3	0.39	0	0,3,3	0.00	-
4	GOL	C	1306	-	5,5,5	0.29	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	1306	-	-	0/0/0/0	0/0/0/0
4	GOL	A	1307	-	-	0/4/4/4	0/0/0/0
3	ACT	B	1305	-	-	0/0/0/0	0/0/0/0
4	GOL	B	1306	-	-	0/4/4/4	0/0/0/0
3	ACT	C	1305	-	-	0/0/0/0	0/0/0/0
4	GOL	C	1306	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1306	ACT	4	0
3	B	1305	ACT	2	0
3	C	1305	ACT	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/281 (100%)	0.40	32 (11%) 7 7	17, 25, 42, 61	1 (0%)
1	B	281/281 (100%)	0.49	39 (13%) 4 4	18, 25, 42, 60	1 (0%)
1	C	281/281 (100%)	0.58	38 (13%) 4 4	17, 27, 42, 59	1 (0%)
All	All	843/843 (100%)	0.49	109 (12%) 5 5	17, 26, 43, 61	3 (0%)

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	236	ASP	7.4
1	C	234	GLY	7.3
1	B	234	GLY	6.1
1	C	236	ASP	6.0
1	B	232	GLU	5.9
1	C	237	ILE	5.9
1	B	235	GLY	5.6
1	A	236	ASP	5.5
1	A	234	GLY	5.4
1	C	131	VAL	4.8
1	B	24	ALA	4.6
1	C	235	GLY	4.5
1	A	233	ARG	4.4
1	B	233	ARG	4.1
1	B	129	VAL	4.0
1	B	237	ILE	4.0
1	B	238	THR	3.8
1	C	232	GLU	3.6
1	C	191	GLU	3.6
1	A	135	ALA	3.6
1	A	130	GLY	3.5
1	C	153	VAL	3.5
1	B	43	THR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	129	VAL	3.4
1	C	25	LYS	3.4
1	C	155	ILE	3.4
1	A	24	ALA	3.3
1	A	129	VAL	3.3
1	C	24	ALA	3.3
1	B	153	VAL	3.3
1	C	233	ARG	3.3
1	C	137	ILE	3.2
1	B	44	PRO	3.2
1	A	131	VAL	3.2
1	C	205	ASN	3.2
1	A	194	SER	3.1
1	A	155	ILE	3.1
1	B	155	ILE	3.1
1	C	134	GLY	3.1
1	A	100	LEU	3.0
1	C	100	LEU	3.0
1	B	65	PHE	3.0
1	C	133	ALA	3.0
1	B	231	PHE	3.0
1	C	127	ILE	2.9
1	B	154	LEU	2.9
1	B	131	VAL	2.9
1	C	194	SER	2.9
1	B	54	TYR	2.9
1	A	54	TYR	2.9
1	A	232	GLU	2.9
1	C	54	TYR	2.8
1	B	130	GLY	2.8
1	A	137	ILE	2.8
1	A	44	PRO	2.7
1	C	52	LEU	2.7
1	C	64	CYS	2.7
1	A	132	GLY	2.7
1	B	135	ALA	2.7
1	A	237	ILE	2.7
1	B	191	GLU	2.7
1	B	53	THR	2.6
1	A	235	GLY	2.6
1	C	132	GLY	2.6
1	C	130	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	154	LEU	2.6
1	C	195	GLY	2.6
1	B	134	GLY	2.5
1	A	238	THR	2.5
1	A	52	LEU	2.5
1	B	52	LEU	2.5
1	A	240	ARG	2.5
1	C	138	LEU	2.5
1	C	135	ALA	2.5
1	A	127	ILE	2.5
1	A	153	VAL	2.5
1	B	230	ASN	2.4
1	C	65	PHE	2.4
1	A	72	GLU	2.4
1	C	196	ASN	2.4
1	C	86	VAL	2.4
1	C	99	PRO	2.4
1	A	198	GLU	2.4
1	C	72	GLU	2.4
1	B	30	GLU	2.3
1	B	137	ILE	2.3
1	A	196	ASN	2.3
1	A	53	THR	2.3
1	C	53	THR	2.3
1	C	57	VAL	2.3
1	A	231	PHE	2.3
1	B	64	CYS	2.3
1	A	195	GLY	2.3
1	B	86	VAL	2.2
1	B	262	SER	2.2
1	B	25	LYS	2.2
1	B	133	ALA	2.2
1	A	259	GLU	2.2
1	C	198	GLU	2.2
1	C	154	LEU	2.1
1	B	138	LEU	2.1
1	B	263	LYS	2.1
1	B	127	ILE	2.1
1	B	250	GLN	2.1
1	B	72	GLU	2.1
1	B	59	LEU	2.1
1	B	132	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	43	THR	2.0
1	A	230	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ACT	C	1305	4/4	0.86	0.18	2.37	27,29,30,30	0
3	ACT	A	1306	4/4	0.93	0.13	0.78	16,20,21,22	0
4	GOL	C	1306	6/6	0.94	0.13	-0.30	33,37,38,40	0
4	GOL	A	1307	6/6	0.96	0.10	-0.38	28,30,31,33	0
4	GOL	B	1306	6/6	0.96	0.10	-0.75	24,28,31,35	0
3	ACT	B	1305	4/4	0.95	0.10	-0.94	18,20,21,22	0
2	CA	A	1305	1/1	0.99	0.07	-1.16	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.