



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2016 – 11:12 AM GMT

PDB ID : 4XMU
Title : Crystal Structure of Met260Ala mutant of E. coli Aminopeptidase N in complex with L-Alanine
Authors : Addlagatta, A.; Gumpena, R.; Kishor, C.
Deposited on : 2015-01-15
Resolution : 2.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1643 (2.94-2.90)
Clashscore	102246	1871 (2.94-2.90)
Ramachandran outliers	100387	1824 (2.94-2.90)
Sidechain outliers	100360	1826 (2.94-2.90)
RSRZ outliers	91569	1650 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	866	<div> <div>2%</div> <div>92%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	906	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MLI	A	907	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	866	6951	4402	1204	1318	27	0	2	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	260	ALA	MET	engineered mutation	UNP P04825

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is ALANINE (three-letter code: ALA) (formula: C₃H₇NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			6	3	1	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Na	0	0
			3	3		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	3	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	195	Total	O	0	0
			195	195		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminopeptidase N



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.54Å 120.54Å 170.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.92 – 2.91 24.92 – 2.91	Depositor EDS
% Data completeness (in resolution range)	99.7 (24.92-2.91) 100.0 (24.92-2.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.25	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 2.89Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.146 , 0.210 0.155 , 0.212	Depositor DCC
R_{free} test set	1619 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 41.4	EDS
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 32028 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7169	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, MLI, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/7112	0.87	17/9658 (0.2%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	548	ASP	CB-CG-OD1	8.09	125.58	118.30
1	A	346	ARG	NE-CZ-NH1	8.03	124.31	120.30
1	A	307	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	A	438	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	A	452	ASP	CB-CG-OD1	6.19	123.87	118.30
1	A	293	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	A	633	ASP	CB-CG-OD1	5.72	123.45	118.30
1	A	548	ASP	CB-CG-OD2	-5.71	113.16	118.30
1	A	728	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	A	452	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	A	246	ASP	CB-CG-OD1	5.25	123.02	118.30
1	A	89	ARG	NE-CZ-NH1	5.23	122.91	120.30
1	A	736	ASP	CB-CG-OD1	5.21	122.99	118.30
1	A	556	ARG	NE-CZ-NH1	5.19	122.89	120.30
1	A	728	ARG	NE-CZ-NH1	5.09	122.84	120.30
1	A	600	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	A	337	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6951	0	6792	16	0
2	A	1	0	0	0	0
3	A	6	0	4	0	0
4	A	3	0	0	0	0
5	A	6	0	8	0	0
6	A	7	0	2	0	0
7	A	195	0	0	1	0
All	All	7169	0	6806	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:468:ARG:HD3	7:A:1170:HOH:O	2.02	0.58
1:A:339:VAL:HG21	1:A:611:ALA:HB1	1.87	0.56
1:A:629:PHE:O	1:A:835:ARG:NH2	2.43	0.51
1:A:106:LEU:HD22	1:A:110:TYR:CD2	2.46	0.51
1:A:65:HIS:HB2	1:A:93:LYS:HB3	1.93	0.50
1:A:369:VAL:HG22	1:A:375:PHE:CE2	2.50	0.47
1:A:739:TRP:CG	1:A:745:VAL:HG11	2.50	0.46
1:A:369:VAL:HG22	1:A:375:PHE:CD2	2.51	0.46
1:A:364:ILE:HD13	1:A:387:VAL:HG21	1.98	0.46
1:A:193:ASP:OD2	1:A:217:ARG:HD2	2.17	0.45
1:A:342:ILE:HG23	1:A:565:TRP:CE2	2.52	0.45
1:A:189:LEU:HD23	1:A:189:LEU:C	2.39	0.44
1:A:265:ASN:HB2	1:A:268:LEU:O	2.18	0.43
1:A:223:ALA:N	1:A:224:PRO:CD	2.82	0.42
1:A:776:MET:O	1:A:782:ILE:HD11	2.20	0.41
1:A:631:ILE:HD12	1:A:631:ILE:N	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	866/866 (100%)	840 (97%)	23 (3%)	3 (0%)	46 78

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	204	ARG
1	A	264[A]	GLU
1	A	264[B]	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	742/740 (100%)	719 (97%)	23 (3%)	47 81

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	ASP
1	A	24	ASP
1	A	48	SER
1	A	107	GLU
1	A	151	LYS
1	A	161	ARG
1	A	164	GLN
1	A	172	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	204	ARG
1	A	286	LYS
1	A	317	SER
1	A	327	ASP
1	A	356	ASP
1	A	501	LYS
1	A	517	GLN
1	A	547	SER
1	A	590	SER
1	A	607	LYS
1	A	608	ILE
1	A	680	LEU
1	A	775	THR
1	A	800	GLU
1	A	860	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ALA	A	902	2	0,5,5	0.00	-	2,6,6	0.32	0
5	GOL	A	906	-	5,5,5	0.38	0	5,5,5	0.61	0
6	MLI	A	907	-	0,6,6	0.00	-	0,7,7	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ALA	A	902	2	-	0/0/4/4	0/0/0/0
5	GOL	A	906	-	-	0/4/4/4	0/0/0/0
6	MLI	A	907	-	-	0/0/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	866/866 (100%)	-0.50	17 (1%) 68 64	15, 26, 44, 75	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	78	GLU	4.5
1	A	504	HIS	4.1
1	A	5	PRO	4.1
1	A	169	ASN	3.9
1	A	503	GLY	3.7
1	A	168	GLU	3.4
1	A	855	GLU	3.1
1	A	856	ASN	2.8
1	A	505	PRO	2.7
1	A	760	ASN	2.6
1	A	502	GLY	2.5
1	A	607	LYS	2.2
1	A	398	GLU	2.2
1	A	46	GLY	2.1
1	A	666	SER	2.1
1	A	673	GLU	2.1
1	A	606	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	GOL	A	906	6/6	0.90	0.25	3.66	51,62,64,66	0
6	MLI	A	907	7/7	0.92	0.23	2.92	38,45,47,50	0
4	NA	A	904	1/1	0.92	0.20	1.49	23,23,23,23	0
4	NA	A	903	1/1	0.89	0.13	0.92	21,21,21,21	0
3	ALA	A	902	6/6	0.98	0.15	0.29	34,36,38,39	0
4	NA	A	905	1/1	0.95	0.11	-0.82	25,25,25,25	0
2	ZN	A	901	1/1	0.99	0.03	-3.11	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.