



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 11:33 PM GMT

PDB ID : 1XSE  
Title : Crystal Structure of Guinea Pig 11beta-Hydroxysteroid Dehydrogenase Type 1  
Authors : Ogg, D.; Elleby, B.; Norstrom, C.; Stefansson, K.; Abrahmsen, L.; Oppermann, U.; Svensson, S.  
Deposited on : 2004-10-19  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	
1	B	295	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4655 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 11beta-hydroxysteroid dehydrogenase type 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	0	0
			2105	1351	351	388	15			
1	B	274	Total	C	N	O	S	0	0	0
			2105	1351	351	388	15			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MET	-	CLONING ARTIFACT	UNP Q6QLL4
A	4	GLY	-	CLONING ARTIFACT	UNP Q6QLL4
A	5	SER	-	CLONING ARTIFACT	UNP Q6QLL4
A	6	SER	-	CLONING ARTIFACT	UNP Q6QLL4
A	7	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
A	8	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
A	9	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
A	10	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
A	11	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
A	12	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
A	13	SER	-	CLONING ARTIFACT	UNP Q6QLL4
A	14	SER	-	CLONING ARTIFACT	UNP Q6QLL4
A	15	GLY	-	CLONING ARTIFACT	UNP Q6QLL4
A	16	LEU	-	CLONING ARTIFACT	UNP Q6QLL4
A	17	VAL	-	CLONING ARTIFACT	UNP Q6QLL4
A	18	PRO	-	CLONING ARTIFACT	UNP Q6QLL4
A	19	ARG	-	CLONING ARTIFACT	UNP Q6QLL4
A	20	GLY	-	CLONING ARTIFACT	UNP Q6QLL4
A	21	SER	-	CLONING ARTIFACT	UNP Q6QLL4
A	22	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
A	23	MET	-	CLONING ARTIFACT	UNP Q6QLL4
B	3	MET	-	CLONING ARTIFACT	UNP Q6QLL4
B	4	GLY	-	CLONING ARTIFACT	UNP Q6QLL4
B	5	SER	-	CLONING ARTIFACT	UNP Q6QLL4
B	6	SER	-	CLONING ARTIFACT	UNP Q6QLL4

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	7	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
B	8	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
B	9	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
B	10	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
B	11	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
B	12	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
B	13	SER	-	CLONING ARTIFACT	UNP Q6QLL4
B	14	SER	-	CLONING ARTIFACT	UNP Q6QLL4
B	15	GLY	-	CLONING ARTIFACT	UNP Q6QLL4
B	16	LEU	-	CLONING ARTIFACT	UNP Q6QLL4
B	17	VAL	-	CLONING ARTIFACT	UNP Q6QLL4
B	18	PRO	-	CLONING ARTIFACT	UNP Q6QLL4
B	19	ARG	-	CLONING ARTIFACT	UNP Q6QLL4
B	20	GLY	-	CLONING ARTIFACT	UNP Q6QLL4
B	21	SER	-	CLONING ARTIFACT	UNP Q6QLL4
B	22	HIS	-	CLONING ARTIFACT	UNP Q6QLL4
B	23	MET	-	CLONING ARTIFACT	UNP Q6QLL4

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	235	Total 235	O 235	0	0
3	B	114	Total 114	O 114	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.39 Å   118.39 Å   184.77 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	10.00 – 2.50 9.98 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.2 (10.00-2.50) 99.7 (9.98-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.26 (at 2.50 Å)	Xtriage
Refinement program	REFMAC 5.2.0001	Depositor
R, $R_{free}$	0.192   ,   0.267 0.253   ,   0.326	Depositor DCC
$R_{free}$ test set	1162 reflections (5.14%)	DCC
Wilson B-factor (Å <sup>2</sup> )	56.4	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 78.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 22609 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4655	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.75	0/2145	0.84	5/2897 (0.2%)
1	B	0.64	1/2145 (0.0%)	0.77	3/2897 (0.1%)
All	All	0.70	1/4290 (0.0%)	0.81	8/5794 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	80	GLU	CG-CD	6.45	1.61	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	295	LEU	CA-CB-CG	7.42	132.37	115.30
1	A	252	ARG	NE-CZ-NH2	-7.11	116.75	120.30
1	A	204	ASN	N-CA-C	7.05	130.04	111.00
1	B	204	ASN	N-CA-C	6.35	128.14	111.00
1	A	198	ARG	NE-CZ-NH2	-6.17	117.22	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2105	0	2135	63	0
1	B	2105	0	2135	56	0
2	A	48	0	26	4	0
2	B	48	0	26	3	0
3	A	235	0	0	5	0
3	B	114	0	0	6	0
All	All	4655	0	4322	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:VAL:HB	3:A:1518:HOH:O	1.25	1.30
1:B:122:LEU:H	1:B:143:ASN:HD21	1.06	1.00
1:A:140:MET:HE1	1:A:144:PHE:CD2	2.12	0.84
1:A:122:LEU:H	1:A:143:ASN:HD21	1.28	0.80
1:B:220:THR:HG22	1:B:223:ALA:H	1.49	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/295 (92%)	260 (96%)	10 (4%)	2 (1%)	26	46
1	B	272/295 (92%)	263 (97%)	9 (3%)	0	100	100
All	All	544/590 (92%)	523 (96%)	19 (4%)	2 (0%)	39	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	65	ALA
1	A	270	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/242 (93%)	204 (91%)	20 (9%)	12	23
1	B	224/242 (93%)	199 (89%)	25 (11%)	7	14
All	All	448/484 (93%)	403 (90%)	45 (10%)	9	18

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	36	LYS
1	B	122	LEU
1	B	274	LYS
1	B	63	VAL
1	B	126	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	72	GLN
1	B	119	ASN
1	B	130	HIS
1	A	143	ASN
1	B	120	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NDP	A	1300	-	42,52,52	1.93	6 (14%)	55,80,80	1.98	10 (18%)
2	NDP	B	2300	-	42,52,52	1.90	6 (14%)	55,80,80	1.76	10 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1300	-	-	0/30/77/77	0/5/5/5
2	NDP	B	2300	-	-	0/30/77/77	0/5/5/5

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1300	NDP	C4N-C5N	-4.90	1.38	1.49
2	B	2300	NDP	C4N-C5N	-4.24	1.39	1.49
2	B	2300	NDP	C2N-C3N	2.18	1.40	1.34
2	A	1300	NDP	C2N-C3N	2.28	1.40	1.34
2	B	2300	NDP	C5A-C4A	2.74	1.46	1.40

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1300	NDP	N3A-C2A-N1A	-10.55	120.82	128.89
2	B	2300	NDP	N3A-C2A-N1A	-7.83	122.90	128.89
2	B	2300	NDP	C4A-C5A-N7A	-3.41	106.35	109.48
2	B	2300	NDP	PN-O3-PA	-3.38	123.25	132.73
2	A	1300	NDP	PN-O3-PA	-3.00	124.30	132.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1300	NDP	4	0
2	B	2300	NDP	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	274/295 (92%)	0.48	10 (3%) 46 51	48, 55, 63, 79	0
1	B	274/295 (92%)	1.13	64 (23%) 1 1	49, 56, 64, 78	0
All	All	548/590 (92%)	0.81	74 (13%) 4 4	48, 56, 63, 79	0

The worst 5 of 74 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	297	GLY	12.8
1	B	294	LYS	8.6
1	B	295	LEU	8.5
1	B	104	ALA	7.2
1	B	24	ASN	7.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NDP	A	1300	48/48	0.96	0.18	0.19	50,54,59,61	0
2	NDP	B	2300	48/48	0.94	0.13	-0.90	52,55,61,62	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.