



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 23, 2016 – 08:20 AM EST

PDB ID : 1XV9
Title : crystal structure of CAR/RXR heterodimer bound with SRC1 peptide, fatty acid, and 5b-pregnane-3,20-dione.
Authors : Xu, R.X.; Lambert, M.H.; Wisely, B.B.; Warren, E.N.; Weinert, E.E.; Waitt, G.M.; Williams, J.D.; Moore, L.B.; Willson, T.M.; Moore, J.T.
Deposited on : 2004-10-27
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028320
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028320


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	<div> <div>3%</div> <div>42%</div> <div>51%</div> <div>6%</div> <div>.</div> </div>
1	C	236	<div> <div>38%</div> <div>54%</div> <div>6%</div> <div>.</div> </div>
2	B	246	<div> <div>4%</div> <div>41%</div> <div>54%</div> <div>.</div> </div>
2	D	246	<div> <div>4%</div> <div>46%</div> <div>51%</div> <div>.</div> </div>
3	E	13	<div> <div>23%</div> <div>31%</div> <div>38%</div> <div>8%</div> <div>23%</div> </div>
3	F	13	<div> <div>8%</div> <div>46%</div> <div>46%</div> <div>8%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	13	
3	H	13	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CI2	D	1001[A]	-	-	-	X
4	CI2	D	1001[B]	-	-	-	X
5	F15	A	1003	-	-	-	X
5	F15	C	1002	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8296 atoms, of which 122 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinoic acid receptor RXR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	0	0	0
			1824	1166	311	336	11			
1	C	232	Total	C	N	O	S	0	0	0
			1824	1166	311	336	11			

- Molecule 2 is a protein called Orphan nuclear receptor NR1H3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	246	Total	C	N	O	S	0	0	0
			1990	1287	341	351	11			
2	D	246	Total	C	N	O	S	0	0	0
			1990	1287	341	351	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	VAL	DELETION	UNP Q14994
B	?	-	SER	DELETION	UNP Q14994
B	?	-	PRO	DELETION	UNP Q14994
B	?	-	THR	DELETION	UNP Q14994
D	?	-	VAL	DELETION	UNP Q14994
D	?	-	SER	DELETION	UNP Q14994
D	?	-	PRO	DELETION	UNP Q14994
D	?	-	THR	DELETION	UNP Q14994

- Molecule 3 is a protein called nuclear receptor coactivator 1 isoform 1.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	10	Total	C	N	O	0	0	0
			86	55	18	13			
3	F	12	Total	C	N	O	0	0	0
			102	63	23	16			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	12	Total	C	N	O	0	0	0
			106	66	23	17			
3	H	12	Total	C	N	O	0	0	0
			106	66	23	17			

- Molecule 4 is (5BETA)-PREGNANE-3,20-DIONE (three-letter code: C12) (formula: $C_{21}H_{32}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	H	O	0	1
			110	42	64	4		

- Molecule 5 is PENTADECANOIC ACID (three-letter code: F15) (formula: $C_{15}H_{30}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	H	O	0	0
			46	15	29	2		
5	A	1	Total	C	H	O	0	0
			46	15	29	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	15	Total	O	0	0
			15	15		
6	B	13	Total	O	0	0
			13	13		
6	C	15	Total	O	0	0
			15	15		
6	D	21	Total	O	0	0
			21	21		
6	E	1	Total	O	0	0
			1	1		
6	H	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Retinoic acid receptor RXR-alpha




• Molecule 1: Retinoic acid receptor RXR-alpha




• Molecule 2: Orphan nuclear receptor NR1I3







• Molecule 2: Orphan nuclear receptor NR1I3



• Molecule 3: nuclear receptor coactivator 1 isoform 1



• Molecule 3: nuclear receptor coactivator 1 isoform 1



• Molecule 3: nuclear receptor coactivator 1 isoform 1



• Molecule 3: nuclear receptor coactivator 1 isoform 1



E685	R686	H687	K688	I689	L690	H691	R692	L693	L694	Q695	E696	GLY

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.44Å 128.44Å 214.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.70 49.37 – 2.70	Depositor EDS
% Data completeness (in resolution range)	86.4 (50.00-2.70) 89.6 (49.37-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 2.69Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.181 , 0.239 0.188 , 0.239	Depositor DCC
R_{free} test set	2891 reflections (6.00%)	DCC
Wilson B-factor (Å ²)	63.4	Xtriage
Anisotropy	0.781	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 65.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	0.477 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8296	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: Cl2, F15

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/1862	0.67	0/2523
1	C	0.38	0/1862	0.68	0/2523
2	B	0.38	0/2036	0.65	0/2758
2	D	0.39	0/2036	0.67	0/2758
3	E	0.37	0/87	0.62	0/116
3	F	0.39	0/103	0.69	0/135
3	G	0.31	0/107	0.62	0/142
3	H	0.37	0/107	0.54	0/142
All	All	0.38	0/8200	0.66	0/11097

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1824	0	1849	127	0
1	C	1824	0	1849	147	0
2	B	1990	0	2023	134	0
2	D	1990	0	2023	126	0
3	E	86	0	86	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	102	0	102	6	0
3	G	106	0	105	9	0
3	H	106	0	105	9	0
4	D	46	64	64	7	0
5	A	17	29	29	2	0
5	C	17	29	29	4	0
6	A	15	0	0	0	0
6	B	13	0	0	3	0
6	C	15	0	0	3	0
6	D	21	0	0	1	0
6	E	1	0	0	1	0
6	H	1	0	0	0	0
All	All	8174	122	8264	530	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 530 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:317:SER:HA	1:C:325:LEU:HD12	1.41	1.02
2:B:307:PHE:O	2:B:311:LYS:HG3	1.68	0.94
1:C:426:ARG:HG2	2:D:317:ALA:O	1.72	0.88
2:B:166:THR:O	2:B:170:LEU:HG	1.73	0.88
1:C:297:GLN:O	1:C:301:LEU:HG	1.73	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/236 (98%)	180 (78%)	38 (16%)	12 (5%)	2	4
1	C	230/236 (98%)	182 (79%)	35 (15%)	13 (6%)	2	3
2	B	244/246 (99%)	209 (86%)	29 (12%)	6 (2%)	7	18
2	D	244/246 (99%)	213 (87%)	26 (11%)	5 (2%)	9	24
3	E	8/13 (62%)	6 (75%)	2 (25%)	0	100	100
3	F	10/13 (77%)	4 (40%)	6 (60%)	0	100	100
3	G	10/13 (77%)	7 (70%)	3 (30%)	0	100	100
3	H	10/13 (77%)	6 (60%)	4 (40%)	0	100	100
All	All	986/1016 (97%)	807 (82%)	143 (14%)	36 (4%)	4	9

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	449	THR
2	B	107	SER
2	B	198	ALA
1	C	412	PRO
1	C	449	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/204 (98%)	191 (96%)	9 (4%)	34	65
1	C	200/204 (98%)	192 (96%)	8 (4%)	38	69
2	B	219/219 (100%)	205 (94%)	14 (6%)	22	47
2	D	219/219 (100%)	211 (96%)	8 (4%)	41	72
3	E	9/12 (75%)	8 (89%)	1 (11%)	8	17
3	F	10/12 (83%)	10 (100%)	0	100	100
3	G	11/12 (92%)	10 (91%)	1 (9%)	12	26
3	H	11/12 (92%)	11 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	879/894 (98%)	838 (95%)	41 (5%)	32 63

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	250	ARG
2	B	323	ASN
2	D	296	ILE
2	B	259	TYR
2	B	304	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	323	ASN
1	C	275	GLN
2	D	331	GLN
2	B	344	GLN
1	C	267	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	F15	A	1003	-	13,16,16	0.91	0	13,16,16	1.13	1 (7%)
5	F15	C	1002	-	13,16,16	0.90	0	13,16,16	0.82	0
4	CI2	D	1001[A]	-	26,26,26	1.18	4 (15%)	42,42,42	1.56	7 (16%)
4	CI2	D	1001[B]	-	26,26,26	1.19	3 (11%)	42,42,42	1.31	6 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	F15	A	1003	-	-	0/12/14/14	0/0/0/0
5	F15	C	1002	-	-	0/12/14/14	0/0/0/0
4	CI2	D	1001[A]	-	-	0/4/62/62	0/4/4/4
4	CI2	D	1001[B]	-	-	0/4/62/62	0/4/4/4

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1001[A]	CI2	C22-C4	-2.28	1.52	1.56
4	D	1001[B]	CI2	C15-C13	2.09	1.54	1.50
4	D	1001[B]	CI2	C8-C19	2.10	1.57	1.53
4	D	1001[A]	CI2	C21-C20	2.15	1.58	1.53
4	D	1001[A]	CI2	C20-C19	2.43	1.58	1.53

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1001[A]	CI2	C10-C11-C12	-3.45	106.11	111.06
4	D	1001[A]	CI2	C4-C22-C7	-3.01	96.60	99.75
4	D	1001[B]	CI2	C4-C22-C7	-2.67	96.95	99.75
4	D	1001[B]	CI2	C18-C17-C11	-2.28	106.24	110.28
4	D	1001[A]	CI2	C23-C22-C21	-2.23	106.69	110.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1003	F15	2	0
5	C	1002	F15	4	0
4	D	1001[A]	CI2	5	0
4	D	1001[B]	CI2	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	232/236 (98%)	0.30	8 (3%) 49 49	45, 87, 131, 154	0
1	C	232/236 (98%)	0.16	0 100 100	44, 82, 111, 130	0
2	B	246/246 (100%)	0.36	9 (3%) 45 45	55, 82, 140, 177	0
2	D	246/246 (100%)	0.39	10 (4%) 41 41	56, 82, 135, 173	0
3	E	10/13 (76%)	0.83	3 (30%) 1 0	112, 123, 125, 127	0
3	F	12/13 (92%)	0.55	1 (8%) 14 11	99, 105, 132, 134	0
3	G	12/13 (92%)	0.75	1 (8%) 14 11	115, 123, 132, 144	0
3	H	12/13 (92%)	1.30	3 (25%) 1 1	130, 134, 145, 148	0
All	All	1002/1016 (98%)	0.33	35 (3%) 48 48	44, 85, 134, 177	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	103	PRO	9.4
1	A	256	LEU	6.6
2	B	104	VAL	6.1
2	D	105	GLN	5.6
2	D	301	ARG	5.5

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	F15	C	1002	17/17	0.87	0.47	7.65	95,99,107,107	0
4	CI2	D	1001[A]	23/23	0.93	0.47	7.50	83,86,87,87	55
4	CI2	D	1001[B]	23/23	0.93	0.47	5.68	98,100,100,101	55
5	F15	A	1003	17/17	0.89	0.35	4.20	71,81,83,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.