



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:39 AM GMT

PDB ID : 2XVE
Title : CRYSTAL STRUCTURE OF BACTERIAL FLAVIN-CONTAINING
MONOOXYGENASE
Authors : Cho, H.J.; Kang, B.S.
Deposited on : 2010-10-26
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	<div> <div>7%</div> <div>90%</div> <div>6%</div> <div>.</div> </div>
1	B	464	<div> <div>6%</div> <div>89%</div> <div>7%</div> <div>.</div> </div>
1	C	464	<div> <div>10%</div> <div>88%</div> <div>8%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NO3	A	465	-	X	-	X
2	NO3	B	466	-	X	-	X
2	NO3	B	467	-	X	-	-
2	NO3	C	466	-	X	-	X
4	GOL	B	468	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 11860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FLAVIN-CONTAINING MONOOXYGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	445	Total	C	N	O	S	0	5	0
			3652	2347	600	685	20			
1	B	446	Total	C	N	O	S	0	7	0
			3686	2368	606	692	20			
1	C	444	Total	C	N	O	S	0	4	0
			3637	2336	598	683	20			

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	N	O	0	0
			4	1	3		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	240	Total	O	0	0
			240	240		
5	B	323	Total	O	0	0
			323	323		
5	C	182	Total	O	0	0
			182	182		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: FLAVIN-CONTAINING MONOOXYGENASE



• Molecule 1: FLAVIN-CONTAINING MONOOXYGENASE



• Molecule 1: FLAVIN-CONTAINING MONOOXYGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	159.22Å 68.99Å 140.53Å 90.00° 90.18° 90.00°	Depositor
Resolution (Å)	50.01 – 1.99 32.10 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.7 (50.01-1.99) 82.2 (32.10-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	55.86 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.170 , 0.197 0.192 , 0.215	Depositor DCC
R_{free} test set	4254 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	35.6	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.4	EDS
Estimated twinning fraction	0.013 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 85461 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11860	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FAD, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	1/3781 (0.0%)	0.47	0/5138
1	B	0.41	0/3822	0.49	0/5191
1	C	0.39	0/3763	0.47	0/5115
All	All	0.41	1/11366 (0.0%)	0.47	0/15444

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	320	TYR	CD1-CE1	-5.76	1.30	1.39

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3652	0	3413	15	0
1	B	3686	0	3454	20	0
1	C	3637	0	3395	19	0
2	A	4	0	0	0	0
2	B	8	0	0	0	0
2	C	4	0	0	0	0
3	B	53	0	31	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	53	0	31	0	0
4	B	12	0	16	1	0
4	C	6	0	8	0	0
5	A	240	0	0	1	0
5	B	323	0	0	4	0
5	C	182	0	0	0	0
All	All	11860	0	10348	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:46:GLN:HE22	1:C:65:SER:H	1.28	0.82
1:B:46:GLN:HE22	1:B:65:SER:H	1.24	0.82
1:B:46:GLN:NE2	1:B:65:SER:H	1.93	0.67
1:C:60:GLU:OE2	1:C:127:ARG:NH2	2.27	0.66
1:C:5:ILE:HG13	1:C:157:TYR:HB2	1.79	0.64
1:B:290[B]:ASN:HD21	1:B:292:ARG:HH11	1.48	0.61
1:C:124:THR:HG23	1:C:142:VAL:HG13	1.83	0.61
1:B:229:ARG:HG2	1:B:246:ASN:HB3	1.83	0.61
1:C:76:LYS:HE2	1:C:94:ILE:HG13	1.84	0.60
1:A:76:LYS:HE3	1:A:95:ALA:O	2.02	0.59
1:C:46:GLN:NE2	1:C:65:SER:H	1.98	0.57
1:B:353:MET:O	1:B:357:GLU:HG2	2.06	0.56
1:A:60:GLU:OE2	1:A:127:ARG:NH2	2.38	0.55
1:B:290[B]:ASN:H	1:B:290[B]:ASN:HD22	1.53	0.55
1:A:76:LYS:HE2	1:A:94:ILE:HG13	1.89	0.53
1:A:57[B]:GLU:CD	1:A:57[B]:GLU:H	2.12	0.53
1:B:73:ASN:OD1	4:B:468:GOL:H2	2.07	0.53
1:B:60:GLU:OE2	1:B:127:ARG:NH2	2.38	0.52
1:B:73:ASN:HB2	5:B:2068:HOH:O	2.10	0.51
1:C:124:THR:CG2	1:C:142:VAL:HG13	2.43	0.49
1:A:164:HIS:HD2	5:A:2098:HOH:O	1.96	0.48
1:C:175:GLY:HA2	1:C:178:LYS:HE2	1.96	0.47
1:C:199:THR:HG23	1:C:223:LYS:HB3	1.97	0.47
1:B:73:ASN:CG	5:B:2068:HOH:O	2.53	0.46
1:A:108:LYS:HA	1:A:111:VAL:HG12	1.98	0.46
1:B:126:VAL:HA	1:B:142:VAL:HG12	1.99	0.45
1:B:3:THR:HB	1:B:32:PRO:HB3	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:201:LEU:HB2	1:C:265:VAL:HG21	1.99	0.45
1:B:237:TRP:HB3	1:B:241:TRP:HB3	2.00	0.44
1:C:17:GLN:HE21	1:C:159:VAL:HG11	1.82	0.44
1:A:102:VAL:HA	1:A:440:LEU:HD12	1.99	0.44
1:C:191:ASP:O	1:C:194:GLU:HG2	2.16	0.44
1:A:124:THR:HG23	1:A:142:VAL:HG13	1.99	0.44
1:B:216:CYS:HB2	1:B:224:LEU:HD21	1.99	0.44
1:A:211:ASP:O	1:A:215:GLN:HG2	2.18	0.44
1:A:17:GLN:HE21	1:A:159:VAL:HG11	1.82	0.43
1:C:104:TRP:O	1:C:108:LYS:HG2	2.18	0.43
1:A:199:THR:HG23	1:A:223:LYS:HB3	2.01	0.43
1:C:71:TRP:HA	1:C:99:PRO:HA	2.00	0.43
1:B:164:HIS:HD2	5:B:2148:HOH:O	2.01	0.43
1:B:399:GLU:HG2	1:B:403:HIS:CE1	2.54	0.43
1:B:168:PRO:HB2	1:B:170:VAL:CG2	2.49	0.42
1:A:124:THR:CG2	1:A:142:VAL:HG13	2.50	0.42
1:C:326:ASP:HB3	1:C:385:TYR:CE1	2.55	0.42
1:C:211:ASP:O	1:C:215:GLN:HG2	2.20	0.42
1:C:173:PHE:HB2	1:C:176:PHE:CE1	2.55	0.42
1:C:231:ALA:HA	1:C:232:PRO:HD3	1.95	0.41
1:B:73:ASN:HB3	5:B:2179:HOH:O	2.20	0.41
1:A:168:PRO:HB3	1:A:185:HIS:CE1	2.56	0.41
1:C:75:PRO:HB3	1:C:416:TYR:CD2	2.55	0.41
1:A:31:ILE:HG22	1:A:337:MET:CE	2.51	0.41
1:B:326:ASP:HB3	1:B:385:TYR:CE1	2.55	0.41
1:B:366:GLU:HA	1:B:369:TYR:CE2	2.56	0.41
1:A:78:CYS:HB3	1:A:397:PHE:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/464 (97%)	436 (97%)	12 (3%)	0	100	100
1	B	451/464 (97%)	435 (96%)	16 (4%)	0	100	100
1	C	446/464 (96%)	434 (97%)	12 (3%)	0	100	100
All	All	1345/1392 (97%)	1305 (97%)	40 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/405 (95%)	380 (99%)	4 (1%)	82	85
1	B	389/405 (96%)	386 (99%)	3 (1%)	86	89
1	C	382/405 (94%)	378 (99%)	4 (1%)	82	85
All	All	1155/1215 (95%)	1144 (99%)	11 (1%)	82	85

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	TRP
1	A	142	VAL
1	A	184	LEU
1	A	387	SER
1	B	43	TRP
1	B	69	TYR
1	B	184	LEU
1	C	43	TRP
1	C	69	TYR
1	C	184	LEU
1	C	297	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	40	GLN
1	A	48	ASN
1	A	164	HIS
1	A	282	ASN
1	A	297	ASN
1	A	377	GLN
1	A	414	HIS
1	B	17	GLN
1	B	25	GLN
1	B	40	GLN
1	B	46	GLN
1	B	48	ASN
1	B	132	ASN
1	B	164	HIS
1	B	297	ASN
1	C	17	GLN
1	C	22	GLN
1	C	40	GLN
1	C	46	GLN
1	C	48	ASN
1	C	164	HIS
1	C	297	ASN
1	C	403	HIS
1	C	429	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NO3	A	465	-	3,3,3	3.24	3 (100%)	3,3,3	0.10	0
3	FAD	B	465	-	48,58,58	1.22	6 (12%)	54,89,89	2.14	7 (12%)
2	NO3	B	466	-	3,3,3	3.25	3 (100%)	3,3,3	0.14	0
2	NO3	B	467	-	3,3,3	3.29	3 (100%)	3,3,3	0.10	0
4	GOL	B	468	-	5,5,5	0.29	0	5,5,5	0.26	0
4	GOL	B	469	-	5,5,5	0.31	0	5,5,5	0.28	0
3	FAD	C	465	-	48,58,58	1.21	5 (10%)	54,89,89	2.15	7 (12%)
2	NO3	C	466	-	3,3,3	3.28	3 (100%)	3,3,3	0.11	0
4	GOL	C	467	-	5,5,5	0.33	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NO3	A	465	-	-	0/0/0/0	0/0/0/0
3	FAD	B	465	-	-	0/30/50/50	0/6/6/6
2	NO3	B	466	-	-	0/0/0/0	0/0/0/0
2	NO3	B	467	-	-	0/0/0/0	0/0/0/0
4	GOL	B	468	-	-	0/4/4/4	0/0/0/0
4	GOL	B	469	-	-	0/4/4/4	0/0/0/0
3	FAD	C	465	-	-	0/30/50/50	0/6/6/6
2	NO3	C	466	-	-	0/0/0/0	0/0/0/0
4	GOL	C	467	-	-	0/4/4/4	0/0/0/0

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	465	FAD	C5X-N5	2.03	1.38	1.35
3	B	465	FAD	C2A-N1A	2.40	1.38	1.33
3	C	465	FAD	C2A-N1A	2.52	1.38	1.33
3	C	465	FAD	C1'-N10	2.73	1.51	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	465	FAD	C1'-N10	2.77	1.51	1.48
3	B	465	FAD	C4-N3	2.85	1.38	1.33
2	A	465	NO3	O3-N	2.87	1.40	1.25
2	B	466	NO3	O3-N	2.89	1.40	1.25
2	A	465	NO3	O2-N	2.89	1.40	1.25
2	C	466	NO3	O3-N	2.89	1.40	1.25
2	B	466	NO3	O2-N	2.89	1.40	1.25
2	B	467	NO3	O2-N	2.90	1.40	1.25
3	C	465	FAD	C4-N3	2.90	1.38	1.33
2	B	467	NO3	O3-N	2.93	1.40	1.25
2	C	466	NO3	O2-N	2.94	1.40	1.25
3	C	465	FAD	C4X-N5	3.18	1.38	1.33
3	B	465	FAD	C4X-N5	3.23	1.38	1.33
3	C	465	FAD	C2A-N3A	3.64	1.38	1.32
3	B	465	FAD	C2A-N3A	3.68	1.38	1.32
2	B	466	NO3	O1-N	3.87	1.40	1.24
2	A	465	NO3	O1-N	3.87	1.40	1.24
2	C	466	NO3	O1-N	3.90	1.40	1.24
2	B	467	NO3	O1-N	3.94	1.40	1.24

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	465	FAD	N3A-C2A-N1A	-12.07	119.65	128.89
3	B	465	FAD	N3A-C2A-N1A	-11.98	119.72	128.89
3	B	465	FAD	P-O3P-PA	-3.71	122.30	132.73
3	C	465	FAD	P-O3P-PA	-3.34	123.36	132.73
3	C	465	FAD	C4X-C4-N3	-2.77	119.80	123.59
3	B	465	FAD	C4X-C4-N3	-2.75	119.82	123.59
3	C	465	FAD	C1'-N10-C9A	2.05	121.17	118.86
3	B	465	FAD	C1'-N10-C9A	2.06	121.17	118.86
3	B	465	FAD	C5X-C9A-N10	2.68	119.66	117.62
3	C	465	FAD	C5X-C9A-N10	2.73	119.69	117.62
3	C	465	FAD	C4X-N5-C5X	3.06	120.28	116.76
3	B	465	FAD	C4X-N5-C5X	3.12	120.35	116.76
3	B	465	FAD	C4-N3-C2	5.48	119.99	115.25
3	C	465	FAD	C4-N3-C2	5.51	120.01	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	468	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/464 (95%)	0.47	34 (7%) 17 18	21, 37, 55, 63	0
1	B	446/464 (96%)	0.39	28 (6%) 23 24	20, 33, 46, 55	0
1	C	444/464 (95%)	0.72	47 (10%) 8 9	29, 42, 68, 82	0
All	All	1335/1392 (95%)	0.52	109 (8%) 14 15	20, 37, 63, 82	0

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	446	ASP	5.2
1	B	2	ALA	5.0
1	C	2	ALA	4.6
1	A	231	ALA	4.2
1	A	232	PRO	4.1
1	A	445	SER	4.0
1	C	212	ILE	3.8
1	C	445	SER	3.7
1	C	325	PHE	3.7
1	B	313	ILE	3.7
1	C	313	ILE	3.7
1	C	429	HIS	3.7
1	A	91	GLY	3.6
1	B	134[A]	ASP	3.5
1	C	436	LEU	3.5
1	A	134	ASP	3.5
1	C	239	GLU	3.4
1	A	325	PHE	3.4
1	C	329	ALA	3.3
1	C	222	LYS	3.3
1	B	446	ASP	3.3
1	B	325	PHE	3.2
1	C	248	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	159	VAL	3.1
1	B	212	ILE	3.0
1	A	341	PRO	3.0
1	A	28	GLY	3.0
1	C	231	ALA	3.0
1	C	256	TYR	2.9
1	A	234	GLY	2.9
1	A	441	GLU	2.9
1	C	71	TRP	2.9
1	C	258	ALA	2.9
1	C	364	THR	2.8
1	C	159	VAL	2.8
1	B	160	CYS	2.8
1	C	269	ILE	2.8
1	A	344	SER	2.8
1	C	259	ASP	2.8
1	B	70	LEU	2.7
1	C	184	LEU	2.7
1	A	402	HIS	2.7
1	A	442	ALA	2.7
1	A	230	THR	2.7
1	B	8	LEU	2.7
1	B	162	THR	2.6
1	A	235	TYR	2.6
1	C	431	PRO	2.6
1	B	209	ALA	2.6
1	C	232	PRO	2.6
1	C	262	SER	2.6
1	C	252	THR	2.6
1	A	29	ALA	2.5
1	C	444	LEU	2.5
1	C	229	ARG	2.5
1	A	160	CYS	2.5
1	A	314	GLY	2.5
1	C	365	ALA	2.5
1	B	445	SER	2.5
1	C	67	TYR	2.5
1	C	264	LYS	2.4
1	B	314	GLY	2.4
1	C	421	THR	2.4
1	B	47	TRP	2.4
1	B	161	CYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	14	GLY	2.4
1	C	66	MET	2.4
1	B	73	ASN	2.3
1	B	9	GLY	2.3
1	C	327	ALA	2.3
1	B	322	PHE	2.3
1	B	321	SER	2.3
1	B	447	LYS	2.3
1	A	31	ILE	2.3
1	A	136	GLN	2.3
1	C	302	VAL	2.3
1	C	246	ASN	2.3
1	B	312	TYR	2.3
1	C	430	THR	2.3
1	C	237	TRP	2.3
1	A	313	ILE	2.3
1	B	6	ALA	2.3
1	C	161	CYS	2.2
1	C	442	ALA	2.2
1	C	245	PRO	2.2
1	A	220	GLY	2.2
1	B	441	GLU	2.2
1	B	165	PHE	2.2
1	A	347	GLU	2.2
1	A	338	GLY	2.2
1	C	234	GLY	2.2
1	A	47	TRP	2.2
1	B	258	ALA	2.2
1	C	241	TRP	2.2
1	A	256	TYR	2.2
1	A	405	LYS	2.1
1	A	152[A]	SER	2.1
1	A	233	MET	2.1
1	C	70	LEU	2.1
1	C	402	HIS	2.1
1	C	13	SER	2.1
1	A	217	TYR	2.1
1	A	444	LEU	2.1
1	C	228	TYR	2.1
1	A	260	GLY	2.1
1	C	14	GLY	2.1
1	C	260	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	67	TYR	2.0
1	B	13	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	B	468	6/6	0.81	0.62	10.44	84,84,84,84	0
2	NO3	A	465	4/4	0.96	0.24	5.78	50,50,51,51	0
2	NO3	B	466	4/4	0.93	0.31	4.47	54,55,55,55	0
2	NO3	C	466	4/4	0.97	0.24	4.25	60,61,61,61	0
3	FAD	B	465	53/53	0.97	0.26	1.79	29,31,33,35	0
3	FAD	C	465	53/53	0.95	0.21	1.10	37,41,43,43	0
2	NO3	B	467	4/4	0.94	0.17	0.50	63,63,63,63	0
4	GOL	C	467	6/6	0.86	0.24	-	81,81,81,81	0
4	GOL	B	469	6/6	0.85	0.61	-	85,85,85,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.