



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:32 PM GMT

PDB ID : 4XWO
Title : Structure of Get3 bound to the transmembrane domain of Sec22
Authors : Mateja, A.; Paduch, M.; Chang, H.-Y.; Szydlowska, A.; Kossiakoff, A.A.;
Hegde, R.S.; Keenan, R.J.
Deposited on : 2015-01-29
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	 2% 61% 20% 17%
1	B	354	 4% 61% 22% 16%
1	G	354	 2% 64% 19% 17%
1	H	354	 4% 63% 21% 15%
1	M	354	 4% 64% 21% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	N	354	
1	S	354	
1	T	354	
2	C	230	
2	E	230	
2	I	230	
2	K	230	
2	O	230	
2	Q	230	
2	U	230	
2	W	230	
3	D	217	
3	F	217	
3	J	217	
3	L	217	
3	P	217	
3	R	217	
3	V	217	
3	X	217	
4	a	41	
4	g	41	
4	m	41	
4	s	41	

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 46697 atoms, of which 184 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase GET3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	0
			2319	1472	384	446	17			
1	B	297	Total	C	N	O	S	0	0	0
			2336	1483	386	451	16			
1	G	294	Total	C	N	O	S	0	0	0
			2323	1473	386	447	17			
1	H	300	Total	C	N	O	S	0	0	0
			2371	1502	394	459	16			
1	M	304	Total	C	N	O	S	0	0	0
			2405	1521	400	468	16			
1	N	300	Total	C	N	O	S	0	0	0
			2375	1505	393	461	16			
1	S	296	Total	C	N	O	S	0	0	0
			2330	1477	387	450	16			
1	T	306	Total	C	N	O	S	0	0	0
			2402	1523	398	465	16			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	ASN	ASP	engineered mutation	UNP Q12154
B	57	ASN	ASP	engineered mutation	UNP Q12154
G	57	ASN	ASP	engineered mutation	UNP Q12154
H	57	ASN	ASP	engineered mutation	UNP Q12154
M	57	ASN	ASP	engineered mutation	UNP Q12154
N	57	ASN	ASP	engineered mutation	UNP Q12154
S	57	ASN	ASP	engineered mutation	UNP Q12154
T	57	ASN	ASP	engineered mutation	UNP Q12154

- Molecule 2 is a protein called Antibody heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	218	Total	C	N	O	S	0	0	0
			1641	1038	278	319	6			
2	E	222	Total	C	N	O	S	0	0	0
			1664	1050	282	326	6			
2	I	220	Total	C	N	O	S	0	0	0
			1655	1046	279	324	6			
2	K	216	Total	C	N	O	S	0	0	0
			1628	1031	276	315	6			
2	O	218	Total	C	N	O	S	0	0	0
			1635	1034	277	318	6			
2	Q	221	Total	C	N	O	S	0	0	0
			1655	1045	281	323	6			
2	U	214	Total	C	N	O	S	0	0	0
			1616	1024	273	313	6			
2	W	221	Total	C	N	O	S	0	0	0
			1655	1045	281	323	6			

- Molecule 3 is a protein called Antibody light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	216	Total	C	N	O	S	0	0	0
			1658	1038	276	338	6			
3	F	215	Total	C	N	O	S	0	0	0
			1650	1034	275	335	6			
3	J	215	Total	C	N	O	S	0	0	0
			1650	1034	275	335	6			
3	L	215	Total	C	N	O	S	0	0	0
			1650	1034	275	335	6			
3	P	216	Total	C	N	O	S	0	0	0
			1658	1038	276	338	6			
3	R	216	Total	C	N	O	S	0	0	0
			1658	1038	276	338	6			
3	V	215	Total	C	N	O	S	0	0	0
			1650	1034	275	335	6			
3	X	216	Total	C	N	O	S	0	0	0
			1658	1038	276	338	6			

- Molecule 4 is a protein called Sec22.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	a	20	Total	C	N	O	0	0	0
			100	60	20	20			
4	g	14	Total	C	N	O	0	0	0
			70	42	14	14			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	m	19	Total	C	N	O	0	0	0
			95	57	19	19			
4	s	21	Total	C	N	O	0	0	0
			105	63	21	21			


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total	Mg	0	0
			1	1		
5	H	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		
5	A	1	Total	Mg	0	0
			1	1		
5	T	1	Total	Mg	0	0
			1	1		
5	N	1	Total	Mg	0	0
			1	1		
5	S	1	Total	Mg	0	0
			1	1		
5	M	1	Total	Mg	0	0
			1	1		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	1	Total	Zn	0	0
			1	1		
6	A	1	Total	Zn	0	0
			1	1		
6	S	1	Total	Zn	0	0
			1	1		
6	M	1	Total	Zn	0	0
			1	1		

- Molecule 7 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	A	1	Total 38	C 10	H 11	N 5	O 10	P 2	0	1
7	B	1	Total 39	C 10	H 12	N 5	O 10	P 2	0	1
7	G	1	Total 38	C 10	H 11	N 5	O 10	P 2	0	1
7	H	1	Total 39	C 10	H 12	N 5	O 10	P 2	0	1
7	M	1	Total 38	C 10	H 11	N 5	O 10	P 2	0	1
7	N	1	Total 39	C 10	H 12	N 5	O 10	P 2	0	1
7	S	1	Total 38	C 10	H 11	N 5	O 10	P 2	0	1
7	T	1	Total 39	C 10	H 12	N 5	O 10	P 2	0	1

- Molecule 8 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	A	1	Total	C	H	N	O	P	0	1
			42	10	11	5	13	3		
8	B	1	Total	C	H	N	O	P	0	1
			43	10	12	5	13	3		
8	G	1	Total	C	H	N	O	P	0	1
			42	10	11	5	13	3		
8	H	1	Total	C	H	N	O	P	0	1
			43	10	12	5	13	3		
8	M	1	Total	C	H	N	O	P	0	1
			42	10	11	5	13	3		
8	N	1	Total	C	H	N	O	P	0	1
			43	10	12	5	13	3		
8	S	1	Total	C	H	N	O	P	0	1
			42	10	11	5	13	3		
8	T	1	Total	C	H	N	O	P	0	1
			43	10	12	5	13	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	20	Total	O	0	0
			20	20		
9	B	17	Total	O	0	0
			17	17		
9	C	16	Total	O	0	0
			16	16		
9	D	19	Total	O	0	0
			19	19		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	E	14	Total 14	O 14	0	0
9	F	23	Total 23	O 23	0	0
9	G	20	Total 20	O 20	0	0
9	H	25	Total 25	O 25	0	0
9	I	25	Total 25	O 25	0	0
9	J	34	Total 34	O 34	0	0
9	K	15	Total 15	O 15	0	0
9	L	19	Total 19	O 19	0	0
9	M	20	Total 20	O 20	0	0
9	N	29	Total 29	O 29	0	0
9	O	12	Total 12	O 12	0	0
9	P	15	Total 15	O 15	0	0
9	Q	13	Total 13	O 13	0	0
9	R	9	Total 9	O 9	0	0
9	S	16	Total 16	O 16	0	0
9	T	22	Total 22	O 22	0	0
9	U	8	Total 8	O 8	0	0
9	V	9	Total 9	O 9	0	0
9	W	16	Total 16	O 16	0	0
9	X	9	Total 9	O 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATPase GET3




• Molecule 1: ATPase GET3




• Molecule 1: ATPase GET3







• Molecule 1: ATPase GET3




• Molecule 1: ATPase GET3




• Molecule 1: ATPase GET3







• Molecule 1: ATPase GET3



• Molecule 1: ATPase GET3




• Molecule 2: Antibody heavy chain








- Molecule 2: Antibody heavy chain




- Molecule 2: Antibody heavy chain



- Molecule 2: Antibody heavy chain




- Molecule 2: Antibody heavy chain




- Molecule 2: Antibody heavy chain







• Molecule 2: Antibody heavy chain




• Molecule 2: Antibody heavy chain




• Molecule 3: Antibody light chain




• Molecule 3: Antibody light chain




- Molecule 3: Antibody light chain




- Molecule 3: Antibody light chain




- Molecule 3: Antibody light chain



- Molecule 3: Antibody light chain



- Molecule 3: Antibody light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	112.58Å 119.45Å 147.57Å 71.89° 89.86° 66.61°	Depositor
Resolution (Å)	39.72 – 2.75 39.72 – 2.75	Depositor EDS
% Data completeness (in resolution range)	90.6 (39.72-2.75) 74.7 (39.72-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 2.77Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.196 , 0.247 0.196 , 0.246	Depositor DCC
R_{free} test set	7268 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	49.5	Xtriage
Anisotropy	0.416	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 43.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 156379 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	46697	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.80 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.9625e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, MG, ZN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2356	0.43	0/3175
1	B	0.27	0/2373	0.45	0/3199
1	G	0.27	0/2360	0.44	0/3180
1	H	0.27	0/2409	0.45	0/3247
1	M	0.28	0/2444	0.47	0/3296
1	N	0.28	0/2413	0.46	0/3252
1	S	0.25	0/2368	0.43	0/3192
1	T	0.25	0/2440	0.44	0/3289
2	C	0.28	0/1682	0.47	0/2292
2	E	0.27	0/1706	0.47	0/2326
2	I	0.29	0/1696	0.48	0/2312
2	K	0.26	0/1669	0.46	0/2275
2	O	0.26	0/1676	0.44	0/2285
2	Q	0.26	0/1697	0.48	0/2314
2	U	0.23	0/1657	0.42	0/2260
2	W	0.25	0/1697	0.47	0/2314
3	D	0.27	0/1694	0.47	0/2299
3	F	0.28	0/1686	0.46	0/2288
3	J	0.29	0/1686	0.47	0/2288
3	L	0.26	0/1686	0.46	0/2288
3	P	0.25	0/1694	0.44	0/2299
3	R	0.26	0/1694	0.44	0/2299
3	V	0.23	0/1686	0.41	0/2288
3	X	0.25	0/1694	0.45	0/2299
All	All	0.26	0/46163	0.45	0/62556

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2319	0	2316	77	0
1	B	2336	0	2325	92	0
1	G	2323	0	2319	65	0
1	H	2371	0	2354	66	1
1	M	2405	0	2388	60	1
1	N	2375	0	2362	64	0
1	S	2330	0	2324	65	1
1	T	2402	0	2390	61	0
2	C	1641	0	1604	31	0
2	E	1664	0	1625	38	0
2	I	1655	0	1616	30	0
2	K	1628	0	1595	32	0
2	O	1635	0	1591	33	0
2	Q	1655	0	1619	58	1
2	U	1616	0	1579	40	0
2	W	1655	0	1619	53	2
3	D	1658	0	1611	31	0
3	F	1650	0	1607	44	0
3	J	1650	0	1607	36	0
3	L	1650	0	1607	40	0
3	P	1658	0	1611	46	0
3	R	1658	0	1611	35	0
3	V	1650	0	1608	44	0
3	X	1658	0	1611	39	0
4	a	100	0	22	0	0
4	g	70	0	16	0	0
4	m	95	0	21	0	0
4	s	105	0	23	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	G	1	0	0	0	0
5	H	1	0	0	0	0
5	M	1	0	0	0	0
5	N	1	0	0	0	0
5	S	1	0	0	0	0
5	T	1	0	0	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	G	1	0	0	0	0
6	M	1	0	0	0	0
6	S	1	0	0	0	0
7	A	27	11	12	0	0
7	B	27	12	12	1	0
7	G	27	11	12	0	0
7	H	27	12	12	1	0
7	M	27	11	9	0	0
7	N	27	12	12	0	0
7	S	27	11	11	0	0
7	T	27	12	12	1	0
8	A	31	11	12	2	0
8	B	31	12	12	2	0
8	G	31	11	12	1	0
8	H	31	12	12	2	0
8	M	31	11	10	1	0
8	N	31	12	12	0	0
8	S	31	11	11	0	0
8	T	31	12	12	2	0
9	A	20	0	0	2	0
9	B	17	0	0	0	0
9	C	16	0	0	0	0
9	D	19	0	0	1	0
9	E	14	0	0	3	0
9	F	23	0	0	1	0
9	G	20	0	0	6	0
9	H	25	0	0	1	0
9	I	25	0	0	2	0
9	J	34	0	0	2	0
9	K	15	0	0	2	0
9	L	19	0	0	1	0
9	M	20	0	0	1	0
9	N	29	0	0	1	0
9	O	12	0	0	0	0
9	P	15	0	0	0	0
9	Q	13	0	0	4	0
9	R	9	0	0	0	0
9	S	16	0	0	3	0
9	T	22	0	0	2	0
9	U	8	0	0	2	0
9	V	9	0	0	1	0
9	W	16	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	X	9	0	0	2	0
All	All	46513	184	44766	1043	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 1043 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:145:ARG:HG2	3:F:145:ARG:HH11	1.20	1.05
2:Q:209:HIS:H	2:Q:215:LYS:HZ1	1.07	1.02
1:G:137:ASP:OD2	1:H:175:ARG:NH1	1.97	0.98
1:N:69:LYS:O	1:N:75:ARG:NH1	1.94	0.98
3:L:145:ARG:HG2	3:L:145:ARG:HH11	1.31	0.94

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:Q:141:SER:OG	1:S:188:GLU:OE2[1_564]	1.82	0.38
1:H:48:ASN:OD1	2:W:24:SER:OG[1_455]	2.13	0.07
1:M:188:GLU:OE1	2:W:138:LYS:NZ[1_455]	2.15	0.05

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/354 (80%)	277 (98%)	6 (2%)	0	100	100
1	B	287/354 (81%)	280 (98%)	7 (2%)	0	100	100
1	G	284/354 (80%)	278 (98%)	5 (2%)	1 (0%)	39	72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	290/354 (82%)	285 (98%)	4 (1%)	1 (0%)	46	77
1	M	296/354 (84%)	289 (98%)	6 (2%)	1 (0%)	46	77
1	N	290/354 (82%)	282 (97%)	4 (1%)	4 (1%)	14	38
1	S	288/354 (81%)	284 (99%)	3 (1%)	1 (0%)	46	77
1	T	298/354 (84%)	293 (98%)	4 (1%)	1 (0%)	46	77
2	C	214/230 (93%)	204 (95%)	9 (4%)	1 (0%)	34	67
2	E	220/230 (96%)	207 (94%)	12 (6%)	1 (0%)	34	67
2	I	216/230 (94%)	207 (96%)	8 (4%)	1 (0%)	34	67
2	K	212/230 (92%)	205 (97%)	7 (3%)	0	100	100
2	O	214/230 (93%)	204 (95%)	10 (5%)	0	100	100
2	Q	219/230 (95%)	205 (94%)	12 (6%)	2 (1%)	21	52
2	U	210/230 (91%)	199 (95%)	9 (4%)	2 (1%)	19	48
2	W	219/230 (95%)	206 (94%)	12 (6%)	1 (0%)	34	67
3	D	214/217 (99%)	209 (98%)	4 (2%)	1 (0%)	34	67
3	F	213/217 (98%)	207 (97%)	5 (2%)	1 (0%)	34	67
3	J	213/217 (98%)	208 (98%)	5 (2%)	0	100	100
3	L	213/217 (98%)	206 (97%)	5 (2%)	2 (1%)	21	52
3	P	214/217 (99%)	208 (97%)	5 (2%)	1 (0%)	34	67
3	R	214/217 (99%)	208 (97%)	6 (3%)	0	100	100
3	V	213/217 (98%)	205 (96%)	6 (3%)	2 (1%)	21	52
3	X	214/217 (99%)	208 (97%)	5 (2%)	1 (0%)	34	67
All	All	5748/6408 (90%)	5564 (97%)	159 (3%)	25 (0%)	39	72

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	206	SER
3	F	206	SER
2	I	59	SER
3	L	206	SER
1	N	283	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/309 (85%)	257 (98%)	6 (2%)	58	87
1	B	263/309 (85%)	258 (98%)	5 (2%)	65	89
1	G	264/309 (85%)	262 (99%)	2 (1%)	86	96
1	H	269/309 (87%)	263 (98%)	6 (2%)	60	87
1	M	273/309 (88%)	266 (97%)	7 (3%)	54	84
1	N	270/309 (87%)	267 (99%)	3 (1%)	80	94
1	S	263/309 (85%)	262 (100%)	1 (0%)	93	98
1	T	270/309 (87%)	268 (99%)	2 (1%)	88	97
2	C	182/193 (94%)	180 (99%)	2 (1%)	80	94
2	E	185/193 (96%)	184 (100%)	1 (0%)	92	97
2	I	184/193 (95%)	184 (100%)	0	100	100
2	K	181/193 (94%)	179 (99%)	2 (1%)	80	94
2	O	180/193 (93%)	178 (99%)	2 (1%)	80	94
2	Q	184/193 (95%)	181 (98%)	3 (2%)	70	91
2	U	179/193 (93%)	178 (99%)	1 (1%)	90	97
2	W	184/193 (95%)	181 (98%)	3 (2%)	70	91
3	D	191/192 (100%)	189 (99%)	2 (1%)	82	95
3	F	190/192 (99%)	188 (99%)	2 (1%)	80	94
3	J	190/192 (99%)	189 (100%)	1 (0%)	92	97
3	L	190/192 (99%)	188 (99%)	2 (1%)	80	94
3	P	191/192 (100%)	189 (99%)	2 (1%)	82	95
3	R	191/192 (100%)	189 (99%)	2 (1%)	82	95
3	V	190/192 (99%)	190 (100%)	0	100	100
3	X	191/192 (100%)	190 (100%)	1 (0%)	92	97
All	All	5118/5552 (92%)	5060 (99%)	58 (1%)	80	94

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	J	111	ARG
1	M	126	LEU
2	U	207	VAL
2	K	138	LYS
3	L	98	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
3	R	127	GLN
3	R	140	ASN
1	T	154	GLN
1	N	301	GLN
2	Q	173	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 12 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	ADP	A	403[B]	5	22,29,29	4.05	7 (31%)	27,45,45	2.72	3 (11%)
8	ATP	A	404[A]	5	24,33,33	2.27	9 (37%)	31,52,52	2.20	7 (22%)
7	ADP	B	402[B]	1,5	22,29,29	4.07	8 (36%)	27,45,45	2.80	4 (14%)
8	ATP	B	403[A]	5	24,33,33	2.28	9 (37%)	31,52,52	2.21	6 (19%)
7	ADP	G	403[B]	5	22,29,29	4.04	6 (27%)	27,45,45	2.81	4 (14%)
8	ATP	G	404[A]	5	24,33,33	2.25	9 (37%)	31,52,52	2.33	8 (25%)
7	ADP	H	402[B]	1,5	22,29,29	4.06	8 (36%)	27,45,45	2.81	4 (14%)
8	ATP	H	403[A]	5	24,33,33	2.25	9 (37%)	31,52,52	2.22	7 (22%)
7	ADP	M	403[B]	1,5	22,29,29	4.14	5 (22%)	27,45,45	2.99	5 (18%)
8	ATP	M	404[A]	1,5	24,33,33	2.45	8 (33%)	31,52,52	2.44	12 (38%)
7	ADP	N	402[B]	5	22,29,29	4.08	7 (31%)	27,45,45	2.79	3 (11%)
8	ATP	N	403[A]	5	24,33,33	2.29	9 (37%)	31,52,52	2.26	7 (22%)
7	ADP	S	403[B]	5	22,29,29	4.05	7 (31%)	27,45,45	2.79	4 (14%)
8	ATP	S	404[A]	5	24,33,33	2.31	9 (37%)	31,52,52	2.26	10 (32%)
7	ADP	T	402[B]	1,5	22,29,29	4.09	7 (31%)	27,45,45	2.75	3 (11%)
8	ATP	T	403[A]	1,5	24,33,33	2.28	9 (37%)	31,52,52	2.23	11 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ADP	A	403[B]	5	-	0/12/32/32	0/3/3/3
8	ATP	A	404[A]	5	-	0/18/38/38	0/3/3/3
7	ADP	B	402[B]	1,5	-	0/12/32/32	0/3/3/3
8	ATP	B	403[A]	5	-	0/18/38/38	0/3/3/3
7	ADP	G	403[B]	5	-	0/12/32/32	0/3/3/3
8	ATP	G	404[A]	5	-	0/18/38/38	0/3/3/3
7	ADP	H	402[B]	1,5	-	0/12/32/32	0/3/3/3
8	ATP	H	403[A]	5	-	0/18/38/38	0/3/3/3
7	ADP	M	403[B]	1,5	-	0/12/32/32	0/3/3/3
8	ATP	M	404[A]	1,5	-	0/18/38/38	0/3/3/3
7	ADP	N	402[B]	5	-	0/12/32/32	0/3/3/3
8	ATP	N	403[A]	5	-	0/18/38/38	0/3/3/3
7	ADP	S	403[B]	5	-	0/12/32/32	0/3/3/3
8	ATP	S	404[A]	5	-	0/18/38/38	0/3/3/3
7	ADP	T	402[B]	1,5	-	0/12/32/32	0/3/3/3
8	ATP	T	403[A]	1,5	-	0/18/38/38	0/3/3/3

The worst 5 of 126 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	G	403[B]	ADP	O4'-C4'	-5.60	1.32	1.45
7	N	402[B]	ADP	O4'-C4'	-5.55	1.32	1.45
7	A	403[B]	ADP	O4'-C4'	-5.53	1.32	1.45
7	T	402[B]	ADP	O4'-C4'	-5.52	1.32	1.45
7	S	403[B]	ADP	O4'-C4'	-5.51	1.32	1.45

The worst 5 of 98 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	403[B]	ADP	N3-C2-N1	-11.83	119.83	128.89
7	N	402[B]	ADP	N3-C2-N1	-11.75	119.90	128.89
7	S	403[B]	ADP	N3-C2-N1	-11.72	119.92	128.89
7	T	402[B]	ADP	N3-C2-N1	-11.72	119.92	128.89
7	H	402[B]	ADP	N3-C2-N1	-11.56	120.04	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	404[A]	ATP	2	0
7	B	402[B]	ADP	1	0
8	B	403[A]	ATP	2	0
8	G	404[A]	ATP	1	0
7	H	402[B]	ADP	1	0
8	H	403[A]	ATP	2	0
8	M	404[A]	ATP	1	0
7	T	402[B]	ADP	1	0
8	T	403[A]	ATP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/354 (82%)	-0.02	6 (2%) 68 63	39, 68, 136, 194	0
1	B	297/354 (83%)	0.08	14 (4%) 35 28	29, 56, 148, 213	0
1	G	294/354 (83%)	-0.10	7 (2%) 62 56	28, 56, 134, 178	0
1	H	300/354 (84%)	0.04	14 (4%) 35 28	30, 59, 143, 244	0
1	M	304/354 (85%)	-0.05	13 (4%) 39 32	32, 57, 152, 226	0
1	N	300/354 (84%)	-0.02	9 (3%) 54 47	28, 52, 136, 183	0
1	S	296/354 (83%)	-0.01	7 (2%) 62 56	48, 77, 141, 197	0
1	T	306/354 (86%)	0.07	16 (5%) 31 24	43, 64, 139, 201	0
2	C	218/230 (94%)	-0.44	5 (2%) 64 57	32, 55, 96, 150	0
2	E	222/230 (96%)	-0.52	3 (1%) 78 73	33, 53, 98, 150	0
2	I	220/230 (95%)	-0.49	4 (1%) 71 66	28, 49, 95, 190	0
2	K	216/230 (93%)	-0.38	2 (0%) 85 82	42, 62, 119, 165	0
2	O	218/230 (94%)	-0.27	3 (1%) 78 73	36, 64, 142, 199	0
2	Q	221/230 (96%)	-0.29	5 (2%) 64 57	35, 63, 132, 223	0
2	U	214/230 (93%)	0.92	39 (18%) 2 1	55, 112, 204, 273	0
2	W	221/230 (96%)	0.11	10 (4%) 37 30	50, 82, 133, 172	0
3	D	216/217 (99%)	-0.41	1 (0%) 91 90	39, 62, 91, 165	0
3	F	215/217 (99%)	-0.52	1 (0%) 91 90	35, 52, 78, 129	0
3	J	215/217 (99%)	-0.47	1 (0%) 91 90	29, 50, 81, 178	0
3	L	215/217 (99%)	-0.42	1 (0%) 91 90	42, 71, 100, 182	0
3	P	216/217 (99%)	-0.02	12 (5%) 28 21	42, 88, 149, 231	0
3	R	216/217 (99%)	-0.25	5 (2%) 64 57	42, 69, 108, 167	0
3	V	215/217 (99%)	0.99	43 (20%) 1 1	74, 141, 202, 241	0
3	X	216/217 (99%)	0.33	18 (8%) 14 9	53, 86, 166, 200	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
4	a	0/41	-	-	-	-
4	g	0/41	-	-	-	-
4	m	0/41	-	-	-	-
4	s	0/41	-	-	-	-
All	All	5864/6572 (89%)	-0.08	239 (4%) 41 34	28, 65, 151, 273	0

The worst 5 of 239 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	U	225	CYS	17.4
1	M	214	GLY	12.6
3	V	157	LEU	11.6
2	U	207	VAL	10.9
3	V	156	ALA	9.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	ADP	N	402[B]	27/27	0.96	0.23	0.38	41,49,61,64	39
8	ATP	N	403[A]	31/31	0.96	0.23	0.23	41,49,60,64	43
5	MG	H	401	1/1	0.93	0.23	-0.11	43,43,43,43	0
8	ATP	M	404[A]	31/31	0.95	0.18	-0.12	34,42,49,52	42
7	ADP	M	403[B]	27/27	0.95	0.17	-0.17	34,42,49,52	38
8	ATP	A	404[A]	31/31	0.96	0.19	-0.18	44,50,59,60	42
7	ADP	G	403[B]	27/27	0.97	0.17	-0.21	38,45,56,60	38

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	ADP	A	403[B]	27/27	0.96	0.18	-0.24	44,50,60,60	38
8	ATP	G	404[A]	31/31	0.97	0.17	-0.25	36,44,54,60	42
8	ATP	H	403[A]	31/31	0.97	0.19	-0.29	35,46,57,62	43
7	ADP	B	402[B]	27/27	0.96	0.18	-0.30	34,42,55,62	39
8	ATP	B	403[A]	31/31	0.96	0.19	-0.31	34,42,53,62	43
8	ATP	S	404[A]	31/31	0.95	0.18	-0.36	49,57,68,72	42
7	ADP	H	402[B]	27/27	0.97	0.18	-0.39	36,45,62,62	39
7	ADP	S	403[B]	27/27	0.96	0.17	-0.42	49,57,69,71	38
8	ATP	T	403[A]	31/31	0.96	0.19	-0.46	48,56,68,73	43
7	ADP	T	402[B]	27/27	0.96	0.17	-0.66	48,56,70,73	39
6	ZN	G	402	1/1	0.99	0.06	-	67,67,67,67	0
5	MG	T	401	1/1	0.98	0.23	-	55,55,55,55	0
5	MG	G	401	1/1	0.96	0.18	-	32,32,32,32	0
5	MG	B	401	1/1	0.96	0.24	-	54,54,54,54	0
5	MG	S	401	1/1	0.90	0.15	-	53,53,53,53	0
6	ZN	M	402	1/1	0.94	0.06	-	72,72,72,72	0
5	MG	M	401	1/1	0.98	0.07	-	50,50,50,50	0
6	ZN	S	402	1/1	0.97	0.05	-	97,97,97,97	0
6	ZN	A	402	1/1	0.97	0.04	-	65,65,65,65	0
5	MG	A	401	1/1	0.93	0.18	-	57,57,57,57	0
5	MG	N	401	1/1	0.97	0.15	-	50,50,50,50	0

6.5 Other polymers

There are no such residues in this entry.