



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:38 PM GMT

PDB ID : 1Y0X  
Title : Thyroxine-Thyroid Hormone Receptor Interactions  
Authors : Sandler, B.; Webb, P.; Apriletti, J.W.; Huber, B.R.; Togashi, M.; Cunha Lima, S.T.; Juric, S.; Nilsson, S.; Wagner, R.; Fletterick, R.J.; Baxter, J.D.  
Deposited on : 2004-11-16  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	281	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	T44	X	500	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Thyroid hormone receptor beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	250	Total	C	N	O	S	0	0	0
			1934	1243	325	350	16			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
X	181	MET	-	EXPRESSION TAG	UNP P10828
X	182	GLY	-	EXPRESSION TAG	UNP P10828
X	183	SER	-	EXPRESSION TAG	UNP P10828
X	184	SER	-	EXPRESSION TAG	UNP P10828
X	185	HIS	-	EXPRESSION TAG	UNP P10828
X	186	HIS	-	EXPRESSION TAG	UNP P10828
X	187	HIS	-	EXPRESSION TAG	UNP P10828
X	188	HIS	-	EXPRESSION TAG	UNP P10828
X	189	HIS	-	EXPRESSION TAG	UNP P10828
X	190	HIS	-	EXPRESSION TAG	UNP P10828
X	191	SER	-	EXPRESSION TAG	UNP P10828
X	192	SER	-	EXPRESSION TAG	UNP P10828
X	193	GLY	-	EXPRESSION TAG	UNP P10828
X	194	LEU	-	EXPRESSION TAG	UNP P10828
X	195	VAL	-	EXPRESSION TAG	UNP P10828
X	196	PRO	-	EXPRESSION TAG	UNP P10828
X	197	ARG	-	EXPRESSION TAG	UNP P10828
X	198	GLY	-	EXPRESSION TAG	UNP P10828
X	199	SER	-	EXPRESSION TAG	UNP P10828
X	200	HIS	-	EXPRESSION TAG	UNP P10828
X	201	MET	-	EXPRESSION TAG	UNP P10828

- Molecule 2 is CACODYLATE ION (three-letter code: CAC) (formula:  $C_2H_6AsO_2$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	X	1	Total 1	As 1	0	0
2	X	1	Total 1	As 1	0	0
2	X	1	Total 1	As 1	0	0
2	X	1	Total 1	As 1	0	0
2	X	1	Total 1	As 1	0	0
2	X	1	Total 1	As 1	0	0

- Molecule 3 is 3,5,3',5'-TETRAIODO-L-THYRONINE (three-letter code: T44) (formula:  $C_{15}H_{11}I_4NO_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	X	1	Total	C	I	N	O	0
			24	15	4	1	4	


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	X	10	Total	O	0	0
			10	10		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thyroid hormone receptor beta-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.79 Å 68.79 Å 130.40 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	59.76 – 3.10 29.79 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (59.76-3.10) 99.3 (29.79-2.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 2.80 Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.220 , 0.266 0.272 , 0.316	Depositor DCC
$R_{free}$ test set	736 reflections (11.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.8	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 32.4	EDS
Estimated twinning fraction	0.031 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 9192 reflections	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	1974	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, T44

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	X	1.88	44/1977 (2.2%)	1.50	23/2681 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	1

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	220	GLU	CD-OE2	9.06	1.35	1.25
1	X	267	GLU	CG-CD	8.02	1.64	1.51
1	X	403	PHE	CD1-CE1	-7.55	1.24	1.39
1	X	248	GLU	CG-CD	7.51	1.63	1.51
1	X	299	GLU	CD-OE1	7.41	1.33	1.25
1	X	267	GLU	CD-OE2	7.30	1.33	1.25
1	X	438	ARG	NE-CZ	-6.75	1.24	1.33
1	X	245	PHE	CE2-CZ	6.58	1.49	1.37
1	X	449	GLU	CA-CB	6.52	1.68	1.53
1	X	223	LYS	CA-CB	6.43	1.68	1.53
1	X	403	PHE	CD2-CE2	-6.20	1.26	1.39
1	X	433	ALA	CA-CB	-6.18	1.39	1.52
1	X	248	GLU	CD-OE2	6.17	1.32	1.25
1	X	457	GLU	CD-OE1	6.06	1.32	1.25
1	X	249	ASP	CB-CG	6.04	1.64	1.51
1	X	295	GLU	CD-OE2	6.01	1.32	1.25
1	X	245	PHE	CD1-CE1	5.90	1.51	1.39
1	X	406	TYR	CD2-CE2	-5.89	1.30	1.39

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	299	GLU	CG-CD	5.87	1.60	1.51
1	X	418	TRP	CD2-CE2	-5.81	1.34	1.41
1	X	457	GLU	CD-OE2	5.78	1.32	1.25
1	X	387	ALA	CA-CB	5.75	1.64	1.52
1	X	217	GLU	CA-CB	5.75	1.66	1.53
1	X	245	PHE	CG-CD1	5.62	1.47	1.38
1	X	316	ARG	CB-CG	-5.40	1.38	1.52
1	X	409	TYR	CD1-CE1	5.39	1.47	1.39
1	X	264	VAL	CA-CB	5.34	1.66	1.54
1	X	460	GLU	C-O	5.28	1.33	1.23
1	X	207	SER	CA-CB	-5.27	1.45	1.52
1	X	455	PHE	CE1-CZ	-5.25	1.27	1.37
1	X	208	ILE	C-O	5.24	1.33	1.23
1	X	333	GLU	CG-CD	5.23	1.59	1.51
1	X	239	TRP	CB-CG	5.21	1.59	1.50
1	X	395	TYR	CD1-CE1	5.19	1.47	1.39
1	X	393	GLU	CD-OE2	5.18	1.31	1.25
1	X	319	VAL	CA-CB	-5.15	1.44	1.54
1	X	404	GLU	CD-OE2	-5.15	1.20	1.25
1	X	441	HIS	CB-CG	5.14	1.59	1.50
1	X	406	TYR	CG-CD1	-5.12	1.32	1.39
1	X	392	ILE	CA-CB	-5.10	1.43	1.54
1	X	272	PHE	CB-CG	-5.09	1.42	1.51
1	X	225	VAL	CB-CG1	5.08	1.63	1.52
1	X	370	VAL	CB-CG1	-5.07	1.42	1.52
1	X	250	ILE	CA-CB	-5.05	1.43	1.54

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	282	ARG	NE-CZ-NH2	-10.80	114.90	120.30
1	X	320	ARG	NE-CZ-NH2	-10.45	115.08	120.30
1	X	316	ARG	NE-CZ-NH2	7.57	124.08	120.30
1	X	419	PRO	CA-N-CD	-7.00	101.71	111.50
1	X	300	ASP	CB-CG-OD2	6.77	124.39	118.30
1	X	429	ARG	NE-CZ-NH1	6.73	123.67	120.30
1	X	320	ARG	NE-CZ-NH1	6.71	123.66	120.30
1	X	265	ASP	CB-CG-OD2	6.63	124.27	118.30
1	X	438	ARG	NE-CZ-NH2	-6.45	117.07	120.30
1	X	282	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	X	316	ARG	NE-CZ-NH1	-5.80	117.40	120.30
1	X	288	LYS	CD-CE-NZ	-5.79	98.39	111.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	281	THR	OG1-CB-CG2	-5.62	97.07	110.00
1	X	351	ASP	CB-CG-OD2	5.56	123.31	118.30
1	X	421	LEU	CA-CB-CG	5.56	128.09	115.30
1	X	367	ASP	CB-CG-OD1	5.49	123.24	118.30
1	X	423	MET	CG-SD-CE	-5.44	91.50	100.20
1	X	409	TYR	CB-CG-CD1	5.37	124.22	121.00
1	X	454	LEU	CB-CG-CD2	-5.37	101.87	111.00
1	X	413	HIS	CA-CB-CG	-5.28	104.62	113.60
1	X	243	ARG	NE-CZ-NH1	5.22	122.91	120.30
1	X	379	MET	CG-SD-CE	-5.04	92.13	100.20
1	X	225	VAL	CA-CB-CG2	-5.04	103.34	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	459	PHE	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	1934	0	1893	59	2
2	X	6	0	0	4	0
3	X	24	0	9	9	0
4	X	10	0	0	1	0
All	All	1974	0	1902	59	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:298:CYS:HB2	2:X:503:CAC:AS	2.03	1.19

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:416:HIS:O	1:X:420:LYS:HG3	1.68	0.92
1:X:298:CYS:HG	2:X:502:CAC:AS	1.61	0.91
1:X:277:THR:O	1:X:281:THR:HG23	1.81	0.78
1:X:250:ILE:O	1:X:340:GLN:NE2	2.16	0.77
1:X:451:PHE:HB3	1:X:456:LEU:HD21	1.68	0.75
1:X:264:VAL:HG11	1:X:446:CYS:SG	2.26	0.75
1:X:331:ASN:C	1:X:333:GLU:H	1.92	0.72
1:X:313:MET:HE1	3:X:500:T44:I5'	2.62	0.70
1:X:313:MET:CE	3:X:500:T44:I5'	3.12	0.68
1:X:331:ASN:O	1:X:333:GLU:N	2.27	0.67
1:X:435:HIS:CE1	3:X:500:T44:I5'	3.18	0.66
1:X:451:PHE:CB	1:X:456:LEU:HD21	2.25	0.66
1:X:276:ILE:O	1:X:280:ILE:HG12	1.96	0.65
1:X:337:THR:OG1	1:X:340:GLN:HG3	1.97	0.64
1:X:453:PRO:O	1:X:457:GLU:HG3	1.99	0.63
1:X:279:ALA:HB2	1:X:331:ASN:OD1	2.00	0.62
1:X:214:PRO:HB2	1:X:219:TRP:NE1	2.15	0.61
1:X:353:ILE:HD13	3:X:500:T44:I3	2.72	0.59
1:X:331:ASN:C	1:X:333:GLU:N	2.57	0.57
1:X:225:VAL:HG21	1:X:399:PHE:HE2	1.68	0.57
1:X:451:PHE:HB3	1:X:456:LEU:CD2	2.36	0.56
1:X:391:ARG:CZ	2:X:504:CAC:AS	3.14	0.55
1:X:444:VAL:CG2	1:X:445:GLU:N	2.71	0.53
1:X:276:ILE:HG13	3:X:500:T44:I5	2.80	0.51
1:X:294:CYS:CB	2:X:501:CAC:AS	3.19	0.51
1:X:276:ILE:HD11	3:X:500:T44:I5'	2.80	0.51
1:X:213:GLU:HB3	1:X:214:PRO:CD	2.41	0.51
1:X:313:MET:HE3	3:X:500:T44:I5'	2.80	0.50
1:X:205:GLN:HE22	1:X:212:PRO:HB3	1.76	0.50
1:X:214:PRO:HB2	1:X:219:TRP:CD1	2.47	0.50
1:X:199:SER:O	1:X:201:MET:N	2.44	0.50
1:X:233:ASN:O	1:X:234:ALA:C	2.50	0.49
1:X:239:TRP:CD2	1:X:320:ARG:HD2	2.47	0.49
1:X:218:GLU:OE1	1:X:405:HIS:ND1	2.43	0.49
1:X:372:LEU:HD13	1:X:403:PHE:HA	1.95	0.49
1:X:250:ILE:HG12	1:X:267:GLU:OE2	2.13	0.48
1:X:235:GLN:O	1:X:236:GLY:C	2.52	0.48
1:X:277:THR:N	1:X:278:PRO:HD2	2.30	0.47
1:X:341:LEU:HD11	1:X:346:LEU:HD12	1.95	0.47
1:X:250:ILE:HA	1:X:267:GLU:OE2	2.14	0.47
1:X:415:THR:O	1:X:420:LYS:HE3	2.15	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:444:VAL:HG23	1:X:445:GLU:N	2.31	0.45
1:X:225:VAL:HG21	1:X:399:PHE:CE2	2.51	0.45
1:X:328:LEU:CD1	1:X:354:PHE:CE1	3.01	0.44
1:X:250:ILE:HG23	1:X:267:GLU:HG2	1.99	0.44
1:X:370:VAL:O	1:X:374:GLN:HG3	2.16	0.44
1:X:451:PHE:N	1:X:451:PHE:CD1	2.86	0.43
1:X:328:LEU:HD12	1:X:354:PHE:CE1	2.54	0.43
1:X:410:ARG:O	1:X:411:LYS:C	2.54	0.43
1:X:342:LYS:HG3	1:X:347:GLY:HA2	2.01	0.41
1:X:435:HIS:NE2	3:X:500:T44:I5'	3.23	0.41
1:X:225:VAL:HG12	1:X:371:ALA:HB1	2.02	0.41
1:X:429:ARG:HD3	4:X:5:HOH:O	2.19	0.41
1:X:277:THR:N	1:X:278:PRO:CD	2.84	0.41
1:X:205:GLN:NE2	1:X:212:PRO:HB3	2.35	0.41
1:X:246:LEU:HA	1:X:247:PRO:HD3	1.99	0.41
1:X:353:ILE:CD1	3:X:500:T44:I3	3.39	0.40
1:X:328:LEU:HG	1:X:354:PHE:CZ	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:201:MET:O	1:X:457:GLU:OE1[3_664]	1.28	0.92
1:X:201:MET:O	1:X:457:GLU:CD[3_664]	1.86	0.34

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	246/281 (88%)	229 (93%)	9 (4%)	8 (3%)	5 26

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	200	HIS
1	X	202	GLU
1	X	236	GLY
1	X	248	GLU
1	X	332	GLY
1	X	234	ALA
1	X	445	GLU
1	X	448	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	206/246 (84%)	177 (86%)	29 (14%)	4 18

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	205	GLN
1	X	238	HIS
1	X	240	LYS
1	X	241	GLN
1	X	242	LYS
1	X	243	ARG
1	X	248	GLU
1	X	267	GLU
1	X	270	SER
1	X	276	ILE
1	X	295	GLU
1	X	298	CYS
1	X	302	ILE
1	X	306	LYS
1	X	320	ARG
1	X	331	ASN
1	X	334	MET
1	X	359	SER
1	X	362	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	X	373	LEU
1	X	381	SER
1	X	398	SER
1	X	411	LYS
1	X	415	THR
1	X	419	PRO
1	X	444	VAL
1	X	446	CYS
1	X	452	PRO
1	X	453	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	205	GLN
1	X	235	GLN
1	X	241	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 6 are modelled with single atom - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	T44	X	500	-	22,25,25	1.33	4 (18%)	30,36,36	1.96	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	T44	X	500	-	-	0/8/12/12	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	500	T44	C3'-I3'	-2.37	2.03	2.10
3	X	500	T44	O4-C4	-2.21	1.34	1.39
3	X	500	T44	C4-C5	2.08	1.47	1.40
3	X	500	T44	C4'-C5'	2.72	1.46	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	500	T44	O4'-C4'-C3'	-3.17	111.77	120.12
3	X	500	T44	C6'-C5'-I5'	-2.76	113.38	118.60
3	X	500	T44	C4'-C3'-I3'	-2.09	115.46	119.49
3	X	500	T44	C1-C7-C8	2.24	119.52	114.34
3	X	500	T44	O4-C1'-C6'	2.70	127.23	119.04
3	X	500	T44	C4'-C5'-I5'	3.18	125.61	119.49
3	X	500	T44	O4'-C4'-C5'	6.01	135.95	120.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	X	500	T44	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	X	250/281 (88%)	-0.06	3 (1%)	81 64	32, 52, 78, 99	2 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	199	SER	4.6
1	X	249	ASP	2.3
1	X	238	HIS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	T44	X	500	24/24	0.95	0.18	-1.09	37,51,54,72	0
2	CAC	X	502	1/5	0.97	0.09	-	97,97,97,97	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CAC	X	506	1/5	0.79	0.24	-	61,61,61,61	1
2	CAC	X	501	1/5	0.69	0.15	-	82,82,82,82	1
2	CAC	X	505	1/5	0.86	0.22	-	51,51,51,51	1
2	CAC	X	503	1/5	0.03	2.03	-	174,174,174,174	1
2	CAC	X	504	1/5	0.92	0.07	-	65,65,65,65	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.