



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:46 AM GMT

PDB ID : 2Y84
Title : DNTR INDUCER BINDING DOMAIN
Authors : Devesse, L.; Smirnova, I.; Lonneborg, R.; Kapp, U.; Brzezinski, P.; Leonard, G.A.; Dian, C.
Deposited on : 2011-02-03
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	<div> <div>65%</div> <div>28%</div> <div>• 7%</div> </div>
1	B	228	<div> <div>70%</div> <div>23%</div> <div>• 7%</div> </div>
1	C	228	<div> <div>67%</div> <div>26%</div> <div>• 7%</div> </div>
1	D	228	<div> <div>64%</div> <div>27%</div> <div>• 7%</div> </div>
1	E	228	<div> <div>62%</div> <div>28%</div> <div>• 7%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	228	
1	G	228	
1	H	228	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	D	1302	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSR-TYPE REGULATORY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	4	0
			1717	1103	306	297	11			
1	B	213	Total	C	N	O	S	0	0	0
			1680	1082	296	291	11			
1	C	213	Total	C	N	O	S	0	0	0
			1672	1076	299	286	11			
1	D	212	Total	C	N	O	S	0	1	0
			1679	1081	299	288	11			
1	E	212	Total	C	N	O	S	0	1	0
			1690	1088	302	289	11			
1	F	214	Total	C	N	O	S	0	0	0
			1678	1081	293	293	11			
1	G	213	Total	C	N	O	S	0	1	0
			1694	1089	303	291	11			
1	H	211	Total	C	N	O	S	0	0	0
			1669	1076	296	286	11			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
A	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
A	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
A	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
A	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
A	306	HIS	-	EXPRESSION TAG	UNP Q7WT50
A	307	HIS	-	EXPRESSION TAG	UNP Q7WT50
B	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
B	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
B	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
B	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
B	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
B	306	HIS	-	EXPRESSION TAG	UNP Q7WT50

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	307	HIS	-	EXPRESSION TAG	UNP Q7WT50
C	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
C	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
C	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
C	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
C	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
C	306	HIS	-	EXPRESSION TAG	UNP Q7WT50
C	307	HIS	-	EXPRESSION TAG	UNP Q7WT50
D	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
D	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
D	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
D	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
D	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
D	306	HIS	-	EXPRESSION TAG	UNP Q7WT50
D	307	HIS	-	EXPRESSION TAG	UNP Q7WT50
E	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
E	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
E	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
E	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
E	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
E	306	HIS	-	EXPRESSION TAG	UNP Q7WT50
E	307	HIS	-	EXPRESSION TAG	UNP Q7WT50
F	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
F	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
F	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
F	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
F	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
F	306	HIS	-	EXPRESSION TAG	UNP Q7WT50
F	307	HIS	-	EXPRESSION TAG	UNP Q7WT50
G	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
G	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
G	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
G	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
G	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
G	306	HIS	-	EXPRESSION TAG	UNP Q7WT50
G	307	HIS	-	EXPRESSION TAG	UNP Q7WT50
H	80	MET	LEU	ENGINEERED MUTATION	UNP Q7WT50
H	302	HIS	-	EXPRESSION TAG	UNP Q7WT50
H	303	HIS	-	EXPRESSION TAG	UNP Q7WT50
H	304	HIS	-	EXPRESSION TAG	UNP Q7WT50
H	305	HIS	-	EXPRESSION TAG	UNP Q7WT50
H	306	HIS	-	EXPRESSION TAG	UNP Q7WT50

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	307	HIS	-	EXPRESSION TAG	UNP Q7WT50

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	E	1	Total C O 6 3 3	0	0
2	H	1	Total C O 6 3 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	13	Total O 13 13	0	0
3	B	16	Total O 16 16	0	0
3	C	18	Total O 18 18	0	0
3	D	10	Total O 10 10	0	0
3	E	20	Total O 20 20	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	8	Total 8	O 8	0	0
3	G	10	Total 10	O 10	0	0
3	H	9	Total 9	O 9	0	0

S299
E300
ALA
HIS
HIS
HIS
HIS
HIS
HIS

• Molecule 1: LYSR-TYPE REGULATORY PROTEIN

Chain E: 62% 28% 7%

MET GLN THR ALA LEU THR THR ARG ASP S89 P92 F93 T96 M103 T104 D105 I106 G107 E108 M109 Y110 F111 M112 P113 P114 L115 M116 L119 A120 Q121 I126 L131 N137 L138 K139 E140 D141 M142 V147 A150 L151 G152 L153 E156 T159 Q163 R168
H169 A170 Y171 S181 A182 M174 F175 H179 P180 K183 S187 H196 L202 N203 H206 V209 M222 R223 F229 I242 V245 Q247 R248 F249 A250 V255 F256 T261 H264 P265 D270 I271 N274 L275 F276 K280 V289 L290 R291 V295

S299
E300
ALA
HIS
HIS
HIS
HIS
HIS
HIS

• Molecule 1: LYSR-TYPE REGULATORY PROTEIN

Chain F: 2% 64% 26% 6%

MET GLN THR ALA LEU THR THR ARG D88 S89 F90 D91 A94 T98 M103 G205 E108 M112 P113 P114 P115 M116 L119 A123 P124 Q127 L131 M134 M137 E140 D141 V147 A150 E156 T159 Q163 F167 R168 H169 A170 F175
D178 H179 P180 S181 A182 K183 M186 Q190 F191 T192 A201 L202 T204 G205 H206 D210 I218 R223 L236 I242 V245 R248 F249 V255 P256 L259 T260 D270 L271 L275 F276 H277 H278 A279 K280 D284 N287 R288 R291 V295

E296
L297
F298
S299
E300
A301
HIS
HIS
HIS
HIS
HIS

• Molecule 1: LYSR-TYPE REGULATORY PROTEIN

Chain G: 2% 62% 29% 7%


MET GLN THR ALA LEU THR THR ARG ASP S89 F90 D91 P92 T96 I106 P113 P114 M116 L119 A120 Q121 I126 N134 K139 E143 L151 E156 L157 Q158 F161 F162 Q163 R164 R168 H169 R170 Y171 V172 C173 M174 F175 P185 L188 K189 Q190
F191 T192 E193 H196 V197 G198 T204 G205 H206 G207 E208 V209 D210 G211 L212 L213 E214 I218 K219 M222 R223 F229 I230 A231 I232 I235 L236 I242 V245 Q247 R248 F249 E254 V255 F256 F257 G258 L259 P265 L268 P269 M174 D270 I271 N274 L275 F276 G277

H278 A279 K280 V289 L290 R291 Q292 F294 A301 HIS HIS HIS HIS HIS

• Molecule 1: LYSR-TYPE REGULATORY PROTEIN

Chain H: 2% 68% 23% 7%

MET GLN THR ALA LEU THR THR ARG ASP SER F90 F93 T96 M103 E108 P113 P114 A120 Q121 I126 P133 L138 K139 E140 D141 M142 V147 A150 L153 F161 F162 R165 L166 F167 R168 H169 C173 P185 H196 V197 G198



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.20Å 111.50Å 213.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.23 – 2.80 48.23 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.23-2.80) 99.3 (48.23-2.80)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 2.81Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.218 , 0.244 0.208 , 0.237	Depositor DCC
R_{free} test set	2437 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	36.5	Xtriage
Anisotropy	0.275	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 48177 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	13607	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/1772	0.41	0/2402
1	B	0.23	0/1726	0.42	0/2343
1	C	0.24	0/1718	0.43	0/2333
1	D	0.24	0/1728	0.44	0/2345
1	E	0.28	0/1736	0.45	0/2353
1	F	0.22	0/1724	0.43	0/2342
1	G	0.22	0/1741	0.42	0/2361
1	H	0.24	0/1715	0.43	0/2327
All	All	0.24	0/13860	0.43	0/18806

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1717	0	1715	36	0
1	B	1680	0	1665	40	0
1	C	1672	0	1652	42	0
1	D	1679	0	1672	56	0
1	E	1690	0	1689	53	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1678	0	1652	43	0
1	G	1694	0	1692	53	0
1	H	1669	0	1661	40	0
2	D	12	0	16	0	0
2	E	6	0	8	0	0
2	H	6	0	8	0	0
3	A	13	0	0	0	0
3	B	16	0	0	0	0
3	C	18	0	0	0	0
3	D	10	0	0	1	0
3	E	20	0	0	0	0
3	F	8	0	0	2	0
3	G	10	0	0	0	0
3	H	9	0	0	0	0
All	All	13607	0	13430	342	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 342 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:170:ARG:HG3	1:G:170:ARG:HH11	1.31	0.93
1:F:255:VAL:HG13	1:F:256:PRO:HD3	1.52	0.90
1:F:170:ARG:HG2	1:F:270:ASP:HB3	1.54	0.88
1:B:300:GLU:HG3	1:B:300:GLU:O	1.75	0.86
1:G:168:ARG:HH11	1:G:270:ASP:HB2	1.42	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/228 (94%)	212 (99%)	3 (1%)	0	100	100
1	B	211/228 (92%)	204 (97%)	7 (3%)	0	100	100
1	C	211/228 (92%)	206 (98%)	5 (2%)	0	100	100
1	D	211/228 (92%)	204 (97%)	7 (3%)	0	100	100
1	E	211/228 (92%)	206 (98%)	5 (2%)	0	100	100
1	F	212/228 (93%)	209 (99%)	3 (1%)	0	100	100
1	G	212/228 (93%)	209 (99%)	3 (1%)	0	100	100
1	H	209/228 (92%)	204 (98%)	5 (2%)	0	100	100
All	All	1692/1824 (93%)	1654 (98%)	38 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/196 (94%)	174 (95%)	10 (5%)	27	60
1	B	177/196 (90%)	171 (97%)	6 (3%)	44	78
1	C	174/196 (89%)	165 (95%)	9 (5%)	29	62
1	D	178/196 (91%)	167 (94%)	11 (6%)	23	54
1	E	179/196 (91%)	167 (93%)	12 (7%)	20	50
1	F	176/196 (90%)	161 (92%)	15 (8%)	13	36
1	G	179/196 (91%)	166 (93%)	13 (7%)	17	44
1	H	176/196 (90%)	166 (94%)	10 (6%)	25	58
All	All	1423/1568 (91%)	1337 (94%)	86 (6%)	23	56

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	139	LYS
1	F	131	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	165	ARG
1	E	156	GLU
1	E	170	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	247	GLN
1	F	203	ASN
1	H	203	ASN
1	F	127	GLN
1	F	137	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	D	1301	-	5,5,5	0.35	0	5,5,5	0.21	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	D	1302	-	5,5,5	0.35	0	5,5,5	0.35	0
2	GOL	E	1301	-	5,5,5	0.35	0	5,5,5	0.27	0
2	GOL	H	1301	-	5,5,5	0.36	0	5,5,5	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	D	1301	-	-	0/4/4/4	0/0/0/0
2	GOL	D	1302	-	-	0/4/4/4	0/0/0/0
2	GOL	E	1301	-	-	0/4/4/4	0/0/0/0
2	GOL	H	1301	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/228 (93%)	-0.30	0 100 100	13, 23, 44, 60	0
1	B	213/228 (93%)	-0.31	1 (0%) 91 88	13, 26, 49, 63	0
1	C	213/228 (93%)	-0.32	0 100 100	17, 28, 45, 62	1 (0%)
1	D	212/228 (92%)	-0.05	3 (1%) 78 69	13, 28, 52, 68	0
1	E	212/228 (92%)	-0.29	1 (0%) 91 88	13, 26, 44, 59	0
1	F	214/228 (93%)	-0.12	4 (1%) 70 59	18, 34, 57, 72	1 (0%)
1	G	213/228 (93%)	0.01	4 (1%) 70 59	15, 33, 63, 77	0
1	H	211/228 (92%)	0.07	4 (1%) 70 59	20, 37, 57, 71	0
All	All	1701/1824 (93%)	-0.16	17 (0%) 84 77	13, 29, 53, 77	2 (0%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	89	SER	3.4
1	F	89	SER	3.2
1	F	88	ASP	3.1
1	D	89	SER	2.9
1	B	301	ALA	2.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	D	1302	6/6	0.91	0.22	2.60	35,38,40,42	0
2	GOL	D	1301	6/6	0.89	0.22	0.31	38,41,43,47	0
2	GOL	E	1301	6/6	0.97	0.16	0.29	25,28,31,31	0
2	GOL	H	1301	6/6	0.93	0.16	-1.23	26,30,32,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.