



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 11:41 PM GMT

PDB ID : 1YA3
Title : Crystal structure of the human mineralocorticoid receptor ligand-binding domain bound to progesterone and harboring the S810L mutation responsible for a severe form of hypertension
Authors : Fagart, J.; Huyet, J.; Pinon, G.M.; Rochel, M.; Mayer, C.; Rafestin-Oblin, M.E.
Deposited on : 2004-12-17
Resolution : 2.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1406 (2.36-2.32)
Clashscore	102246	1509 (2.36-2.32)
Ramachandran outliers	100387	1490 (2.36-2.32)
Sidechain outliers	100360	1491 (2.36-2.32)
RSRZ outliers	91569	1412 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	<div> <div>7%</div> <div>52% 33% 6% 9%</div> </div>
1	B	255	<div> <div>4%</div> <div>49% 37% • 10%</div> </div>
1	C	255	<div> <div>4%</div> <div>45% 38% 7% 9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	STR	B	2001	-	-	-	X
2	STR	C	3001	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5764 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mineralocorticoid receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	0	0
			1890	1235	301	340	14			
1	B	230	Total	C	N	O	S	0	0	0
			1860	1215	296	335	14			
1	C	232	Total	C	N	O	S	0	0	0
			1888	1231	301	342	14			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	730	SER	-	CLONING ARTIFACT	UNP P08235
A	810	LEU	SER	ENGINEERED	UNP P08235
A	910	ALA	CYS	ENGINEERED	UNP P08235
B	730	SER	-	CLONING ARTIFACT	UNP P08235
B	810	LEU	SER	ENGINEERED	UNP P08235
B	910	ALA	CYS	ENGINEERED	UNP P08235
C	730	SER	-	CLONING ARTIFACT	UNP P08235
C	810	LEU	SER	ENGINEERED	UNP P08235
C	910	ALA	CYS	ENGINEERED	UNP P08235

- Molecule 2 is PROGESTERONE (three-letter code: STR) (formula: $C_{21}H_{30}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			23	21	2		
2	B	1	Total	C	O	0	0
			23	21	2		
2	C	1	Total	C	O	0	0
			23	21	2		


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	22	Total	O	0	0
			22	22		
3	B	10	Total	O	0	0
			10	10		
3	C	25	Total	O	0	0
			25	25		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Mineralocorticoid receptor





4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	120.72Å 120.72Å 42.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	18.72 – 2.34 18.72 – 2.34	Depositor EDS
% Data completeness (in resolution range)	81.7 (18.72-2.34) 76.1 (18.72-2.34)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.35Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.237 , 0.284 0.229 , 0.283	Depositor DCC
R_{free} test set	2186 reflections (9.97%)	DCC
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.774	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 64.3	EDS
Estimated twinning fraction	0.043 for -h,-k,l 0.047 for h,-h-k,-l 0.024 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 23881 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5764	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.49 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.9363e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: STR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	2/1934 (0.1%)	0.71	2/2619 (0.1%)
1	B	0.42	1/1904 (0.1%)	0.63	2/2581 (0.1%)
1	C	0.49	2/1931 (0.1%)	0.66	2/2611 (0.1%)
All	All	0.51	5/5769 (0.1%)	0.66	6/7811 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	973	SER	C-N	-15.79	1.04	1.33
1	A	982	HIS	C-O	-7.57	1.08	1.23
1	B	982	HIS	C-O	-7.05	1.09	1.23
1	C	982	HIS	C-O	7.02	1.36	1.23
1	C	956	PHE	C-N	-6.42	1.22	1.34

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	982	HIS	CA-C-O	8.22	137.37	120.10
1	C	846	TYR	N-CA-C	7.20	130.45	111.00
1	A	846	TYR	N-CA-C	5.63	126.22	111.00
1	B	846	TYR	N-CA-C	5.20	125.03	111.00
1	B	982	HIS	CA-C-O	5.16	130.94	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1890	0	1872	103	1
1	B	1860	0	1834	84	0
1	C	1888	0	1867	128	2
2	A	23	0	30	1	0
2	B	23	0	30	2	0
2	C	23	0	30	12	0
3	A	22	0	0	0	0
3	B	10	0	0	1	0
3	C	25	0	0	1	0
All	All	5764	0	5663	315	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 315 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:785:LYS:O	1:C:791:LYS:HD3	1.19	1.32
1:C:771:ARG:HD2	1:C:957:PRO:HG3	1.04	1.02
1:A:836:ASN:ND2	1:A:837:GLU:H	1.55	1.02
1:C:771:ARG:CD	1:C:957:PRO:HG3	1.90	1.01
1:A:836:ASN:HD22	1:A:837:GLU:H	1.09	0.96

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:825:GLN:NE2	1:C:970:LYS:NZ[1_554]	2.00	0.20
1:C:804:TYR:OH	1:C:825:GLN:OE1[1_556]	2.11	0.09
1:A:902:GLU:OE2	1:A:962:GLU:OE2[3_764]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/255 (89%)	206 (91%)	16 (7%)	5 (2%)	8	5
1	B	224/255 (88%)	218 (97%)	5 (2%)	1 (0%)	39	45
1	C	224/255 (88%)	209 (93%)	15 (7%)	0	100	100
All	All	675/765 (88%)	633 (94%)	36 (5%)	6 (1%)	21	21

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	749	ILE
1	A	749	ILE
1	A	907	VAL
1	A	906	MET
1	A	972	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/234 (88%)	193 (93%)	14 (7%)	20	22
1	B	204/234 (87%)	187 (92%)	17 (8%)	14	14
1	C	208/234 (89%)	181 (87%)	27 (13%)	5	4
All	All	619/702 (88%)	561 (91%)	58 (9%)	11	10

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	945	THR
1	C	771	ARG
1	C	945	THR
1	B	959	MET
1	B	970	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	744	ASN
1	B	776	GLN
1	C	825	GLN
1	A	975	ASN
1	C	836	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	STR	A	1001	-	26,26,26	4.21	14 (53%)	42,42,42	1.72	12 (28%)
2	STR	B	2001	-	26,26,26	5.00	15 (57%)	42,42,42	1.94	9 (21%)
2	STR	C	3001	-	26,26,26	4.70	20 (76%)	42,42,42	1.77	9 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	STR	A	1001	-	-	0/4/62/62	0/4/4/4
2	STR	B	2001	-	-	0/4/62/62	0/4/4/4
2	STR	C	3001	-	-	0/4/62/62	0/4/4/4

The worst 5 of 49 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	STR	C13-C17	-3.01	1.50	1.56
2	C	3001	STR	C16-C15	2.06	1.59	1.54
2	A	1001	STR	C16-C17	2.15	1.59	1.54
2	C	3001	STR	C2-C3	2.15	1.54	1.49
2	B	2001	STR	C2-C3	2.21	1.54	1.49

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2001	STR	C17-C13-C14	-5.04	94.49	99.74
2	B	2001	STR	C5-C4-C3	-4.78	117.16	123.75
2	B	2001	STR	O3-C3-C2	-4.02	115.55	121.60
2	C	3001	STR	O3-C3-C2	-3.80	115.88	121.60
2	C	3001	STR	C19-C10-C5	-3.77	102.52	108.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	STR	1	0
2	B	2001	STR	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3001	STR	12	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	233/255 (91%)	0.36	17 (7%)	18 27	25, 45, 67, 75	0
1	B	230/255 (90%)	0.33	9 (3%)	43 55	32, 50, 66, 71	0
1	C	232/255 (90%)	0.30	9 (3%)	43 55	30, 50, 64, 73	0
All	All	695/765 (90%)	0.33	35 (5%)	32 45	25, 48, 66, 75	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	749	ILE	4.3
1	C	950	HIS	4.0
1	B	950	HIS	3.9
1	C	908	THR	3.7
1	C	952	LEU	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	STR	B	2001	23/23	0.94	0.27	2.22	36,37,43,47	0
2	STR	C	3001	23/23	0.90	0.20	1.88	37,40,42,45	0
2	STR	A	1001	23/23	0.94	0.21	1.79	30,32,34,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.