



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:47 AM GMT

PDB ID : 2YBT
Title : CRYSTAL STRUCTURE OF HUMAN ACIDIC CHITINASE IN COMPLEX WITH BISDIONIN C
Authors : Sutherland, T.E.; Andersen, O.A.; Betou, M.; Eggleston, I.M.; Maizels, R.M.; Van Aalten, D.; Allen, J.E.
Deposited on : 2011-03-10
Resolution : 2.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	<div> <div>2%</div> <div>86%</div> <div>12%</div> <div>..</div> </div>
1	B	381	<div> <div>%</div> <div>86%</div> <div>13%</div> <div>..</div> </div>
1	C	381	<div> <div>%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>
1	D	381	<div> <div>3%</div> <div>81%</div> <div>14%</div> <div>..</div> </div>
1	E	381	<div> <div>2%</div> <div>84%</div> <div>14%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	381	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	C	1398	-	-	-	X
2	GOL	E	1398	-	-	-	X
3	DW0	B	1399	-	-	-	X
3	DW0	C	1400	-	-	-	X
3	DW0	E	1400	-	-	-	X
3	DW0	F	1400	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 19339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACIDIC MAMMALIAN CHITINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	0	0	0
			2983	1924	485	560	14			
1	B	377	Total	C	N	O	S	0	0	0
			2983	1924	485	560	14			
1	C	377	Total	C	N	O	S	0	0	0
			2983	1924	485	560	14			
1	D	377	Total	C	N	O	S	0	0	0
			2983	1924	485	560	14			
1	E	377	Total	C	N	O	S	0	0	0
			2983	1924	485	560	14			
1	F	377	Total	C	N	O	S	0	0	0
			2983	1924	485	560	14			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	GLU	-	EXPRESSION TAG	UNP Q9BZP6
A	19	ALA	-	EXPRESSION TAG	UNP Q9BZP6
A	20	GLU	-	EXPRESSION TAG	UNP Q9BZP6
A	45	ASP	ASN	VARIANT	UNP Q9BZP6
A	47	ASN	ASP	VARIANT	UNP Q9BZP6
A	61	MET	ARG	VARIANT	UNP Q9BZP6
B	18	GLU	-	EXPRESSION TAG	UNP Q9BZP6
B	19	ALA	-	EXPRESSION TAG	UNP Q9BZP6
B	20	GLU	-	EXPRESSION TAG	UNP Q9BZP6
B	45	ASP	ASN	VARIANT	UNP Q9BZP6
B	47	ASN	ASP	VARIANT	UNP Q9BZP6
B	61	MET	ARG	VARIANT	UNP Q9BZP6
C	18	GLU	-	EXPRESSION TAG	UNP Q9BZP6
C	19	ALA	-	EXPRESSION TAG	UNP Q9BZP6
C	20	GLU	-	EXPRESSION TAG	UNP Q9BZP6
C	45	ASP	ASN	VARIANT	UNP Q9BZP6
C	47	ASN	ASP	VARIANT	UNP Q9BZP6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	61	MET	ARG	VARIANT	UNP Q9BZP6
D	18	GLU	-	EXPRESSION TAG	UNP Q9BZP6
D	19	ALA	-	EXPRESSION TAG	UNP Q9BZP6
D	20	GLU	-	EXPRESSION TAG	UNP Q9BZP6
D	45	ASP	ASN	VARIANT	UNP Q9BZP6
D	47	ASN	ASP	VARIANT	UNP Q9BZP6
D	61	MET	ARG	VARIANT	UNP Q9BZP6
E	18	GLU	-	EXPRESSION TAG	UNP Q9BZP6
E	19	ALA	-	EXPRESSION TAG	UNP Q9BZP6
E	20	GLU	-	EXPRESSION TAG	UNP Q9BZP6
E	45	ASP	ASN	VARIANT	UNP Q9BZP6
E	47	ASN	ASP	VARIANT	UNP Q9BZP6
E	61	MET	ARG	VARIANT	UNP Q9BZP6
F	18	GLU	-	EXPRESSION TAG	UNP Q9BZP6
F	19	ALA	-	EXPRESSION TAG	UNP Q9BZP6
F	20	GLU	-	EXPRESSION TAG	UNP Q9BZP6
F	45	ASP	ASN	VARIANT	UNP Q9BZP6
F	47	ASN	ASP	VARIANT	UNP Q9BZP6
F	61	MET	ARG	VARIANT	UNP Q9BZP6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 1,1'-PROPANE-1,3-DIYLBIS(3,7-DIMETHYL-3,7-DIHYDRO-1H-PURINE-2,6-DIONE) (three-letter code: DW0) (formula: C₁₇H₂₀N₈O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			29	17	8	4		
3	A	1	Total	C	N	O	0	0
			29	17	8	4		
3	B	1	Total	C	N	O	0	0
			29	17	8	4		
3	B	1	Total	C	N	O	0	0
			29	17	8	4		
3	C	1	Total	C	N	O	0	0
			29	17	8	4		
3	C	1	Total	C	N	O	0	0
			29	17	8	4		
3	D	1	Total	C	N	O	0	0
			29	17	8	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	N	O	0	0
			29	17	8	4		
3	E	1	Total	C	N	O	0	0
			29	17	8	4		
3	E	1	Total	C	N	O	0	0
			29	17	8	4		
3	F	1	Total	C	N	O	0	0
			29	17	8	4		
3	F	1	Total	C	N	O	0	0
			29	17	8	4		

- Molecule 4 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	202	Total	O	0	0
			202	202		
4	B	193	Total	O	0	0
			193	193		
4	C	197	Total	O	0	0
			197	197		
4	D	146	Total	O	0	0
			146	146		
4	E	187	Total	O	0	0
			187	187		
4	F	132	Total	O	0	0
			132	132		

- Molecule 1: ACIDIC MAMMALIAN CHITINASE







• Molecule 1: ACIDIC MAMMALIAN CHITINASE



• Molecule 1: ACIDIC MAMMALIAN CHITINASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	145.21Å 149.07Å 152.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.07 – 2.22 19.99 – 2.22	Depositor EDS
% Data completeness (in resolution range)	99.2 (20.07-2.22) 99.2 (19.99-2.22)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.21Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.181 , 0.228 0.178 , 0.224	Depositor DCC
R_{free} test set	1623 reflections (1.01%)	DCC
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.315	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.6	EDS
Estimated twinning fraction	0.016 for -h,l,k 0.043 for -l,-k,-h 0.015 for k,h,-l 0.005 for k,l,h 0.005 for l,h,k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 161979 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19339	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DW0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	4/3076 (0.1%)	0.92	2/4194 (0.0%)
1	B	1.10	2/3076 (0.1%)	0.91	7/4194 (0.2%)
1	C	1.11	4/3076 (0.1%)	0.92	3/4194 (0.1%)
1	D	1.03	2/3076 (0.1%)	0.88	4/4194 (0.1%)
1	E	1.06	1/3076 (0.0%)	0.88	3/4194 (0.1%)
1	F	0.98	1/3076 (0.0%)	0.84	2/4194 (0.0%)
All	All	1.06	14/18456 (0.1%)	0.89	21/25164 (0.1%)

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	278	ASN	CB-CG	7.91	1.69	1.51
1	D	221	TYR	CD2-CE2	6.12	1.48	1.39
1	C	320	ALA	CA-CB	6.09	1.65	1.52
1	A	278	ASN	CB-CG	6.00	1.64	1.51
1	A	258	GLU	CG-CD	5.84	1.60	1.51

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35	ARG	NE-CZ-NH2	-7.41	116.59	120.30
1	B	136	ASP	CB-CG-OD1	7.07	124.66	118.30
1	D	273	LEU	CB-CG-CD1	5.79	120.85	111.00
1	A	349	LEU	CA-CB-CG	5.73	128.48	115.30
1	E	379	LEU	CB-CG-CD1	5.71	120.71	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2983	0	2828	27	0
1	B	2983	0	2828	24	0
1	C	2983	0	2828	27	0
1	D	2983	0	2828	42	0
1	E	2983	0	2828	31	0
1	F	2983	0	2828	39	0
2	A	6	0	8	0	0
2	B	6	0	8	0	0
2	C	6	0	8	0	0
2	D	6	0	8	0	0
2	E	6	0	8	0	0
2	F	6	0	8	0	0
3	A	58	0	40	4	0
3	B	58	0	40	1	0
3	C	58	0	40	3	0
3	D	58	0	40	5	0
3	E	58	0	40	3	0
3	F	58	0	40	3	0
4	A	202	0	0	1	0
4	B	193	0	0	3	0
4	C	197	0	0	3	0
4	D	146	0	0	1	0
4	E	187	0	0	2	0
4	F	132	0	0	2	0
All	All	19339	0	17256	186	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 186 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:TYR:HD2	1:A:233:THR:HG22	1.26	0.97
1:A:33:GLN:HE22	1:A:72:ASN:HD21	1.20	0.90
1:E:33:GLN:HE22	1:E:72:ASN:HD21	1.27	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:33:GLN:HE22	1:C:72:ASN:HD21	1.30	0.77
1:C:196:GLU:HB3	1:C:199:GLN:HE21	1.49	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/381 (98%)	361 (96%)	14 (4%)	0	100	100
1	B	375/381 (98%)	366 (98%)	9 (2%)	0	100	100
1	C	375/381 (98%)	368 (98%)	7 (2%)	0	100	100
1	D	375/381 (98%)	363 (97%)	11 (3%)	1 (0%)	46	50
1	E	375/381 (98%)	365 (97%)	9 (2%)	1 (0%)	46	50
1	F	375/381 (98%)	364 (97%)	10 (3%)	1 (0%)	46	50
All	All	2250/2286 (98%)	2187 (97%)	60 (3%)	3 (0%)	56	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	174	ASN
1	F	131	GLU
1	D	131	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/314 (99%)	300 (96%)	12 (4%)	40	48
1	B	312/314 (99%)	301 (96%)	11 (4%)	43	52
1	C	312/314 (99%)	301 (96%)	11 (4%)	43	52
1	D	312/314 (99%)	292 (94%)	20 (6%)	22	23
1	E	312/314 (99%)	297 (95%)	15 (5%)	31	36
1	F	312/314 (99%)	293 (94%)	19 (6%)	23	25
All	All	1872/1884 (99%)	1784 (95%)	88 (5%)	32	38

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	171	LYS
1	D	342	PHE
1	F	273	LEU
1	D	172	GLN
1	D	230	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 64 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	117	GLN
1	D	115	ASN
1	F	192	GLN
1	C	129	GLN
1	D	62	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GOL	A	1398	-	5,5,5	0.46	0	5,5,5	1.24	0
3	DW0	A	1399	-	18,32,32	1.99	4 (22%)	17,49,49	2.54	6 (35%)
3	DW0	A	1400	-	18,32,32	1.53	3 (16%)	17,49,49	1.71	4 (23%)
2	GOL	B	1398	-	5,5,5	0.37	0	5,5,5	0.94	0
3	DW0	B	1399	-	18,32,32	1.91	4 (22%)	17,49,49	1.76	4 (23%)
3	DW0	B	1400	-	18,32,32	1.51	2 (11%)	17,49,49	1.85	4 (23%)
2	GOL	C	1398	-	5,5,5	0.28	0	5,5,5	1.05	0
3	DW0	C	1399	-	18,32,32	1.40	3 (16%)	17,49,49	2.08	4 (23%)
3	DW0	C	1400	-	18,32,32	1.73	2 (11%)	17,49,49	1.78	4 (23%)
2	GOL	D	1398	-	5,5,5	0.40	0	5,5,5	0.82	0
3	DW0	D	1399	-	18,32,32	1.90	3 (16%)	17,49,49	2.80	5 (29%)
3	DW0	D	1400	-	18,32,32	1.81	3 (16%)	17,49,49	2.76	5 (29%)
2	GOL	E	1398	-	5,5,5	0.55	0	5,5,5	1.14	1 (20%)
3	DW0	E	1399	-	18,32,32	1.28	3 (16%)	17,49,49	1.98	6 (35%)
3	DW0	E	1400	-	18,32,32	1.19	2 (11%)	17,49,49	1.40	4 (23%)
2	GOL	F	1398	-	5,5,5	0.44	0	5,5,5	0.42	0
3	DW0	F	1399	-	18,32,32	1.90	3 (16%)	17,49,49	2.87	6 (35%)
3	DW0	F	1400	-	18,32,32	1.45	3 (16%)	17,49,49	1.82	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1398	-	-	0/4/4/4	0/0/0/0
3	DW0	A	1399	-	-	0/6/6/6	0/4/4/4
3	DW0	A	1400	-	-	0/6/6/6	0/4/4/4
2	GOL	B	1398	-	-	0/4/4/4	0/0/0/0
3	DW0	B	1399	-	-	0/6/6/6	0/4/4/4
3	DW0	B	1400	-	-	0/6/6/6	0/4/4/4
2	GOL	C	1398	-	-	0/4/4/4	0/0/0/0
3	DW0	C	1399	-	-	0/6/6/6	0/4/4/4
3	DW0	C	1400	-	-	0/6/6/6	0/4/4/4
2	GOL	D	1398	-	-	0/4/4/4	0/0/0/0
3	DW0	D	1399	-	-	0/6/6/6	0/4/4/4
3	DW0	D	1400	-	-	0/6/6/6	0/4/4/4
2	GOL	E	1398	-	-	0/4/4/4	0/0/0/0
3	DW0	E	1399	-	-	0/6/6/6	0/4/4/4
3	DW0	E	1400	-	-	0/6/6/6	0/4/4/4
2	GOL	F	1398	-	-	0/4/4/4	0/0/0/0
3	DW0	F	1399	-	-	0/6/6/6	0/4/4/4
3	DW0	F	1400	-	-	0/6/6/6	0/4/4/4

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1400	DW0	CAC-N3	2.01	1.51	1.47
3	A	1399	DW0	CAD-NAZ	2.03	1.51	1.47
3	E	1399	DW0	C4-N3	2.05	1.41	1.39
3	A	1400	DW0	C6-N1	2.12	1.41	1.38
3	F	1400	DW0	C6-N1	2.14	1.41	1.38

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1399	DW0	CAV-CAP-N10	-8.95	115.25	120.52
3	F	1399	DW0	CAV-CAP-N10	-8.23	115.67	120.52
3	A	1399	DW0	CAV-CAP-N10	-6.55	116.66	120.52
3	D	1400	DW0	CAL-CAK-N1	-5.95	107.27	112.29
3	A	1399	DW0	C6-C5-C4	-4.89	116.43	119.93

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

10 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1399	DW0	1	0
3	A	1400	DW0	3	0
3	B	1399	DW0	1	0
3	C	1399	DW0	2	0
3	C	1400	DW0	1	0
3	D	1399	DW0	2	0
3	D	1400	DW0	3	0
3	E	1399	DW0	3	0
3	F	1399	DW0	2	0
3	F	1400	DW0	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	377/381 (98%)	-0.48	7 (1%) 70 68	16, 25, 44, 63	0
1	B	377/381 (98%)	-0.52	3 (0%) 87 87	17, 26, 42, 63	0
1	C	377/381 (98%)	-0.48	5 (1%) 79 78	15, 27, 45, 66	0
1	D	377/381 (98%)	-0.30	11 (2%) 55 54	17, 34, 59, 75	0
1	E	377/381 (98%)	-0.41	9 (2%) 62 61	15, 28, 49, 70	0
1	F	377/381 (98%)	-0.23	9 (2%) 62 61	19, 37, 61, 73	0
All	All	2262/2286 (98%)	-0.40	44 (1%) 70 68	15, 29, 52, 75	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	397	PRO	5.8
1	D	174	ASN	4.7
1	A	391	SER	4.3
1	B	397	PRO	4.2
1	B	391	SER	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	DW0	C	1400	29/29	0.84	0.19	4.51	30,41,49,49	29
3	DW0	B	1399	29/29	0.92	0.13	3.24	29,38,48,49	0
2	GOL	C	1398	6/6	0.88	0.16	2.81	39,49,56,63	0
3	DW0	E	1400	29/29	0.86	0.16	2.42	33,42,48,50	29
3	DW0	F	1400	29/29	0.82	0.22	2.34	51,55,58,59	29
2	GOL	E	1398	6/6	0.90	0.14	2.23	29,41,47,49	0
3	DW0	A	1399	29/29	0.95	0.11	1.99	25,34,38,45	0
3	DW0	D	1399	29/29	0.92	0.13	1.84	38,44,49,51	0
2	GOL	D	1398	6/6	0.94	0.13	1.79	28,35,42,47	0
3	DW0	F	1399	29/29	0.91	0.13	1.67	44,48,52,54	0
3	DW0	D	1400	29/29	0.89	0.18	1.56	42,45,50,51	29
3	DW0	B	1400	29/29	0.87	0.20	1.42	41,45,49,50	29
2	GOL	B	1398	6/6	0.96	0.10	1.41	34,41,45,46	0
3	DW0	A	1400	29/29	0.83	0.21	1.35	41,50,59,59	29
3	DW0	C	1399	29/29	0.93	0.12	1.19	28,39,56,57	0
2	GOL	F	1398	6/6	0.96	0.10	0.74	36,43,46,50	0
3	DW0	E	1399	29/29	0.93	0.11	0.56	29,36,49,51	0
2	GOL	A	1398	6/6	0.95	0.09	0.36	26,34,40,47	0

6.5 Other polymers ⓘ

There are no such residues in this entry.