



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:47 AM GMT

PDB ID : 2YC4
Title : INTRAFLAGELLAR TRANSPORT COMPLEX 25-27 FROM CHLAMY-
DOMONAS
Authors : Bhogaraju, S.; Taschner, M.; Lorentzen, E.
Deposited on : 2011-03-11
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	<div> <div>56%</div> <div>33%</div> <div>5%</div> <div>6%</div> </div>
1	B	139	<div> <div>58%</div> <div>32%</div> <div>•</div> <div>6%</div> </div>
2	C	208	<div> <div>51%</div> <div>27%</div> <div>6%</div> <div>15%</div> </div>
2	D	208	<div> <div>2%</div> <div>49%</div> <div>29%</div> <div>6%</div> <div>16%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	D	1204	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTRAFLAGELLAR TRANSPORT PROTEIN 25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	0	0
			1001	632	172	191	6			
1	B	131	Total	C	N	O	S	0	0	0
			975	617	167	186	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP B8LIX8
A	-2	ALA	-	EXPRESSION TAG	UNP B8LIX8
A	-1	ALA	-	EXPRESSION TAG	UNP B8LIX8
A	0	SER	-	EXPRESSION TAG	UNP B8LIX8
B	-3	GLY	-	EXPRESSION TAG	UNP B8LIX8
B	-2	ALA	-	EXPRESSION TAG	UNP B8LIX8
B	-1	ALA	-	EXPRESSION TAG	UNP B8LIX8
B	0	SER	-	EXPRESSION TAG	UNP B8LIX8

- Molecule 2 is a protein called SMALL RAB-RELATED GTPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	177	Total	C	N	O	S	0	0	0
			1329	855	211	259	4			
2	D	175	Total	C	N	O	S	0	0	0
			1324	845	215	258	6			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	EXPRESSION TAG	UNP A8HN58
C	-2	ALA	-	EXPRESSION TAG	UNP A8HN58
C	-1	ALA	-	EXPRESSION TAG	UNP A8HN58
C	0	SER	-	EXPRESSION TAG	UNP A8HN58

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	EXPRESSION TAG	UNP A8HN58
D	-2	ALA	-	EXPRESSION TAG	UNP A8HN58
D	-1	ALA	-	EXPRESSION TAG	UNP A8HN58
D	0	SER	-	EXPRESSION TAG	UNP A8HN58

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Ca 1 1	0	0
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Mg 1 1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Cl 1 1	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	4	Total O 4 4	0	0
6	B	1	Total O 1 1	0	0
6	C	8	Total O 8 8	0	0
6	D	13	Total O 13 13	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: INTRAFLAGELLAR TRANSPORT PROTEIN 25

Chain A: 




• Molecule 1: INTRAFLAGELLAR TRANSPORT PROTEIN 25

Chain B: 




• Molecule 2: SMALL RAB-RELATED GTPASE


Chain C: 



• Molecule 2: SMALL RAB-RELATED GTPASE

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.02Å 68.30Å 93.78Å 90.00° 92.42° 90.00°	Depositor
Resolution (Å)	46.85 – 2.80 46.85 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.85-2.80) 99.3 (46.85-2.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.69Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.225 , 0.316 0.222 , 0.309	Depositor DCC
R_{free} test set	787 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	46.3	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 51.6	EDS
Estimated twinning fraction	0.039 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 17450 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4659	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/1017	0.66	0/1378
1	B	0.52	0/991	0.70	0/1347
2	C	0.49	0/1358	0.71	0/1858
2	D	0.51	0/1350	0.70	1/1840 (0.1%)
All	All	0.51	0/4716	0.69	1/6423 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	8	PRO	N-CA-CB	5.87	110.34	103.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	149	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1001	0	954	37	0
1	B	975	0	905	36	0
2	C	1329	0	1251	65	0
2	D	1324	0	1254	60	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	C	1	0	0	0	0
5	D	1	0	0	2	0
6	A	4	0	0	0	0
6	B	1	0	0	0	0
6	C	8	0	0	2	0
6	D	13	0	0	0	0
All	All	4659	0	4364	184	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 184 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:149:LEU:HD12	2:C:149:LEU:H	1.01	1.16
2:C:36:PHE:CE2	2:C:58:ALA:HB1	1.82	1.14
2:C:36:PHE:HE2	2:C:58:ALA:HB1	0.98	1.11
2:C:77:ALA:HB1	2:D:106:GLU:HG3	1.36	1.02
2:D:36:PHE:CE2	2:D:58:ALA:HB1	1.99	0.96

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/139 (91%)	119 (94%)	7 (6%)	1 (1%)	24	58
1	B	127/139 (91%)	115 (91%)	9 (7%)	3 (2%)	7	25
2	C	169/208 (81%)	154 (91%)	11 (6%)	4 (2%)	7	25
2	D	167/208 (80%)	149 (89%)	13 (8%)	5 (3%)	5	18
All	All	590/694 (85%)	537 (91%)	40 (7%)	13 (2%)	8	28

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	93	ASN
1	B	94	ARG
2	C	8	PRO
2	D	150	ASP
2	D	169	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/121 (88%)	96 (91%)	10 (9%)	11	31
1	B	99/121 (82%)	93 (94%)	6 (6%)	23	55
2	C	138/177 (78%)	122 (88%)	16 (12%)	7	20
2	D	138/177 (78%)	123 (89%)	15 (11%)	8	23
All	All	481/596 (81%)	434 (90%)	47 (10%)	10	28

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	61	THR
2	C	103	SER
2	D	103	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	86	ILE
2	C	126	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	103	HIS
2	D	189	ASN
2	C	146	GLN
1	B	27	ASN
1	B	127	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	131/139 (94%)	-0.41	0	100 100	27, 45, 65, 80	0
1	B	131/139 (94%)	-0.27	0	100 100	27, 46, 69, 80	0
2	C	177/208 (85%)	-0.26	1 (0%)	90 86	27, 47, 74, 111	0
2	D	175/208 (84%)	-0.21	4 (2%)	64 52	28, 44, 87, 110	0
All	All	614/694 (88%)	-0.28	5 (0%)	87 81	27, 46, 74, 111	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	172	GLY	3.4
2	C	160	THR	2.7
2	D	39	LYS	2.2
2	D	89	TYR	2.1
2	D	8	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	MG	C	301	1/1	0.87	0.16	-0.32	66,66,66,66	0
5	CL	D	1204	1/1	0.75	0.13	-1.74	79,79,79,79	0
3	CA	A	201	1/1	0.96	0.10	-2.00	63,63,63,63	0
3	CA	B	201	1/1	0.90	0.06	-2.83	67,67,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.