



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:42 PM GMT

PDB ID : 1YDT
Title : STRUCTURE OF CAMP-DEPENDENT PROTEIN KINASE, ALPHA-CATALYTIC SUBUNIT IN COMPLEX WITH H89 PROTEIN KINASE INHIBITOR N-[2-(4-BROMOCINNAMYLAMINO)ETHYL]-5-ISOQUINOLINE
Authors : Engh, R.A.; Girod, A.; Kinzel, V.; Huber, R.; Bossemeyer, D.
Deposited on : 1996-07-24
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	350	
2	I	20	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-AMP-DEPENDENT PROTEIN KINASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	E	336	Total	C	N	O	P	S	0	0	0
			2778	1801	465	501	2	9			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	124	ALA	PRO	CONFLICT	UNP P00517
E	197	TPO	THR	MODIFIED RESIDUE	UNP P00517
E	286	ASP	ASN	CONFLICT	UNP P00517
E	338	SEP	SER	MODIFIED RESIDUE	UNP P00517

- Molecule 2 is a protein called PROTEIN KINASE INHIBITOR PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	20	Total	C	N	O	0	0	0
			157	94	32	31			

- Molecule 3 is N-[2-(4-BROMOCINNAMYLAMINO)ETHYL]-5-ISOQUINOLINE SULFONAMIDE (three-letter code: IQB) (formula: C₂₀H₂₀BrN₃O₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	Br	C	N	O	S		
3	E	1	27	1	20	3	2	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	18	Total	O	0	0
			18	18		
4	I	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: C-AMP-DEPENDENT PROTEIN KINASE



• Molecule 2: PROTEIN KINASE INHIBITOR PEPTIDE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.58Å 76.28Å 80.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.30 19.48 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-2.30) 62.7 (19.48-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.98 (at 1.99Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.194 , (Not available) 0.222 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.360	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.9	EDS
Estimated twinning fraction	0.026 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 20066 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2983	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, IQB, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.67	1/2827 (0.0%)	0.86	6/3807 (0.2%)
2	I	0.65	0/159	0.82	0/212
All	All	0.66	1/2986 (0.0%)	0.86	6/4019 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	199	CYS	CB-SG	-8.22	1.68	1.82

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	59	LEU	N-CA-C	-5.76	95.45	111.00
1	E	277	LEU	CA-CB-CG	5.17	127.19	115.30
1	E	165	ARG	N-CA-C	5.15	124.91	111.00
1	E	114	SER	N-CA-C	5.08	124.71	111.00
1	E	162	LEU	CA-CB-CG	5.07	126.96	115.30
1	E	196	TRP	N-CA-C	5.02	124.56	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2778	0	2757	86	0
2	I	157	0	146	7	0
3	E	27	0	20	4	0
4	E	18	0	0	3	0
4	I	3	0	0	0	0
All	All	2983	0	2923	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:184:ASP:HB2	4:E:413:HOH:O	1.74	0.87
1:E:242:GLN:HG3	1:E:244:ILE:HG12	1.63	0.81
3:E:351:IQB:H3'1	3:E:351:IQB:O2	1.80	0.80
1:E:242:GLN:HG2	1:E:245:GLN:HG3	1.67	0.76
1:E:243:PRO:HA	1:E:246:ILE:HD12	1.70	0.73
1:E:54:PHE:HD2	1:E:82:LEU:HD12	1.54	0.73
1:E:75:ASP:O	1:E:79:VAL:HG23	1.89	0.72
1:E:295:LYS:H	1:E:295:LYS:HD3	1.54	0.71
1:E:104:VAL:HG21	1:E:183:THR:HG22	1.77	0.67
3:E:351:IQB:H3'2	4:E:413:HOH:O	1.94	0.66
1:E:304:ALA:HA	1:E:309:LYS:HE2	1.78	0.65
1:E:319:LYS:HD2	1:E:319:LYS:N	2.11	0.65
1:E:319:LYS:HD2	1:E:319:LYS:H	1.62	0.64
1:E:242:GLN:HG2	1:E:245:GLN:OE1	2.01	0.60
1:E:266:LYS:O	1:E:270:ARG:HG3	2.02	0.60
1:E:345:LYS:H	1:E:345:LYS:HD2	1.66	0.60
1:E:125:GLY:HA3	1:E:174:ILE:O	2.02	0.60
1:E:303:ILE:HG22	1:E:307:GLN:NE2	2.18	0.59
1:E:285:LYS:N	1:E:285:LYS:HD2	2.18	0.58
1:E:78:LYS:HA	1:E:81:LYS:HE3	1.84	0.57
1:E:30:TRP:O	1:E:33:PRO:HD3	2.05	0.57
1:E:277:LEU:O	1:E:283:ASN:ND2	2.38	0.56
1:E:140:GLU:HB2	1:E:141:PRO:HD3	1.88	0.56
1:E:100:PHE:CD1	1:E:101:PRO:HD2	2.41	0.55
1:E:56:ARG:NH2	1:E:331:GLU:O	2.39	0.55
1:E:90:ASN:O	1:E:94:ILE:HG13	2.06	0.55
1:E:189:LYS:HD3	1:E:191:VAL:HG21	1.89	0.54
1:E:209:ILE:HD11	1:E:219:VAL:HG11	1.89	0.54
1:E:304:ALA:CA	1:E:309:LYS:HE2	2.38	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:304:ALA:CB	1:E:309:LYS:HE2	2.38	0.53
1:E:158:HIS:HE1	1:E:220:ASP:OD2	1.90	0.53
1:E:242:GLN:HG2	1:E:245:GLN:CG	2.38	0.53
1:E:69:TYR:HB3	1:E:120:MET:O	2.09	0.53
1:E:45:ARG:HH12	1:E:334:GLU:HA	1.74	0.52
2:I:22:ILE:O	2:I:22:ILE:HG13	2.08	0.52
1:E:301:ASP:OD2	1:E:304:ALA:HB2	2.10	0.51
1:E:295:LYS:H	1:E:295:LYS:CD	2.20	0.51
1:E:294:HIS:CE1	1:E:295:LYS:HE3	2.45	0.51
1:E:333:GLU:HG2	1:E:336:ARG:HH21	1.75	0.51
1:E:73:ILE:HD13	1:E:335:ILE:HD11	1.92	0.50
1:E:244:ILE:HG13	1:E:245:GLN:N	2.26	0.50
1:E:65:THR:HB	1:E:67:ASN:ND2	2.26	0.50
1:E:33:PRO:HA	1:E:96:GLN:OE1	2.12	0.49
1:E:121:GLU:O	3:E:351:IQB:H6	2.12	0.48
1:E:202:PRO:HD3	2:I:22:ILE:HG23	1.94	0.48
3:E:351:IQB:C3'	3:E:351:IQB:O2	2.57	0.48
1:E:74:LEU:O	1:E:115:ASN:HB3	2.14	0.48
2:I:23:HIS:CD2	2:I:24:ASP:H	2.32	0.48
1:E:43:PHE:HB3	1:E:60:VAL:HG13	1.97	0.47
1:E:242:GLN:CG	1:E:245:GLN:HG3	2.42	0.47
1:E:284:LEU:HB3	1:E:285:LYS:HD2	1.97	0.47
1:E:295:LYS:N	1:E:295:LYS:HD3	2.24	0.47
1:E:288:VAL:O	1:E:288:VAL:HG22	2.14	0.46
1:E:285:LYS:H	1:E:285:LYS:HD2	1.79	0.46
1:E:189:LYS:HD3	1:E:191:VAL:CG2	2.45	0.46
1:E:15:VAL:O	1:E:18:PHE:N	2.48	0.46
1:E:62:HIS:NE2	1:E:64:GLU:HG2	2.31	0.46
1:E:91:GLU:O	1:E:95:LEU:HB2	2.16	0.46
1:E:54:PHE:CD2	1:E:82:LEU:HD12	2.42	0.45
1:E:338:SEP:O3P	1:E:339:ILE:HG13	2.16	0.45
1:E:343:CYS:HB3	1:E:346:GLU:OE1	2.16	0.45
1:E:158:HIS:CG	1:E:217:LYS:HD2	2.52	0.45
1:E:103:LEU:HD21	1:E:153:THR:CG2	2.47	0.45
1:E:127:GLU:HA	1:E:173:LEU:HD23	1.99	0.44
1:E:159:SER:OG	1:E:160:LEU:HD13	2.18	0.44
1:E:39:HIS:HB2	1:E:42:GLN:NE2	2.33	0.44
1:E:304:ALA:HB2	1:E:309:LYS:HE2	1.99	0.44
2:I:23:HIS:CD2	2:I:24:ASP:N	2.86	0.43
1:E:148:ALA:HA	1:E:302:TRP:CZ3	2.53	0.43
1:E:222:TRP:O	1:E:226:VAL:HG23	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:11:ILE:HD12	2:I:11:ILE:HG23	1.76	0.43
1:E:233:ALA:HB2	1:E:258:PRO:HG2	2.00	0.43
1:E:310:VAL:CG1	1:E:311:GLU:N	2.82	0.42
1:E:163:ILE:HG22	1:E:165:ARG:HG3	2.01	0.42
1:E:310:VAL:HG12	1:E:311:GLU:N	2.35	0.42
1:E:155:GLU:HG3	1:E:288:VAL:HG11	2.01	0.42
1:E:317:LYS:HA	1:E:317:LYS:HD2	1.72	0.42
1:E:305:ILE:HG12	1:E:310:VAL:HG21	2.01	0.42
1:E:198:LEU:HD23	1:E:209:ILE:HG22	2.02	0.42
1:E:285:LYS:HG2	1:E:286:ASP:OD2	2.20	0.42
1:E:305:ILE:CG1	1:E:310:VAL:HG21	2.50	0.42
1:E:137:ARG:NH1	1:E:232:ALA:O	2.52	0.42
1:E:148:ALA:HA	1:E:302:TRP:HZ3	1.85	0.41
2:I:20:ASN:HD22	2:I:20:ASN:N	2.18	0.41
1:E:345:LYS:HB3	1:E:345:LYS:HE3	1.79	0.41
1:E:289:ASN:O	1:E:293:ASN:HB2	2.20	0.41
1:E:285:LYS:HZ3	1:E:285:LYS:H	1.68	0.41
1:E:133:ARG:NH1	4:E:400:HOH:O	2.38	0.41
1:E:100:PHE:CG	1:E:101:PRO:HD2	2.55	0.41
1:E:17:GLU:O	1:E:20:ALA:HB3	2.21	0.41
1:E:345:LYS:CD	1:E:345:LYS:H	2.30	0.40
1:E:276:ASP:OD1	1:E:279:LYS:HG3	2.20	0.40
1:E:169:PRO:HD2	2:I:19:ARG:NH2	2.37	0.40
1:E:27:LEU:HA	1:E:27:LEU:HD12	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	332/350 (95%)	310 (93%)	19 (6%)	3 (1%)	21 24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	18/20 (90%)	16 (89%)	2 (11%)	0	100	100
All	All	350/370 (95%)	326 (93%)	21 (6%)	3 (1%)	21	24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	41	ASP
1	E	184	ASP
1	E	214	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	294/302 (97%)	260 (88%)	34 (12%)	7	7
2	I	15/15 (100%)	13 (87%)	2 (13%)	5	5
All	All	309/317 (98%)	273 (88%)	36 (12%)	7	7

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	16	LYS
1	E	17	GLU
1	E	27	LEU
1	E	28	LYS
1	E	36	ASN
1	E	37	THR
1	E	40	LEU
1	E	53	SER
1	E	59	LEU
1	E	60	VAL
1	E	92	LYS
1	E	95	LEU
1	E	104	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	160	LEU
1	E	162	LEU
1	E	174	ILE
1	E	198	LEU
1	E	202	PRO
1	E	208	GLU
1	E	210	ILE
1	E	263	SER
1	E	267	ASP
1	E	268	LEU
1	E	269	LEU
1	E	272	LEU
1	E	277	LEU
1	E	278	THR
1	E	285	LYS
1	E	295	LYS
1	E	311	GLU
1	E	318	PHE
1	E	333	GLU
1	E	336	ARG
1	E	345	LYS
2	I	15	ARG
2	I	20	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	42	GLN
1	E	62	HIS
1	E	67	ASN
1	E	77	GLN
1	E	113	ASN
1	E	158	HIS
1	E	307	GLN
2	I	20	ASN
2	I	23	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	E	197	1	8,10,11	1.56	2 (25%)	7,14,16	1.78	3 (42%)
1	SEP	E	338	1	8,9,10	1.03	0	8,12,14	2.29	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	E	197	1	-	0/8/11/13	0/0/0/0
1	SEP	E	338	1	-	0/6/8/10	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	197	TPO	P-O2P	-2.51	1.45	1.54
1	E	197	TPO	P-O1P	-2.48	1.43	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	338	SEP	OG-CB-CA	-4.76	104.22	108.27
1	E	197	TPO	OG1-P-O1P	-2.41	101.09	107.11
1	E	338	SEP	OG-P-O1P	-2.24	101.44	107.14
1	E	338	SEP	O2P-P-O1P	2.06	117.22	110.58
1	E	197	TPO	O3P-P-O2P	2.38	116.44	107.38
1	E	197	TPO	C-CA-N	2.47	114.98	109.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	E	338	SEP	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	IQB	E	351	-	29,29,29	1.24	3 (10%)	38,39,39	1.36	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IQB	E	351	-	-	0/17/17/17	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	351	IQB	C6-N7	2.11	1.36	1.32
3	E	351	IQB	C2-C1	2.76	1.40	1.37
3	E	351	IQB	C9-C8	3.22	1.40	1.36

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	E	351	IQB	C9-C8-N7	-4.20	120.04	123.91
3	E	351	IQB	C1-S-N1'	-2.92	100.60	106.55
3	E	351	IQB	C6-C5-C10	2.07	119.42	117.70
3	E	351	IQB	C1-C10-C5	2.65	119.97	117.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	351	IQB	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	334/350 (95%)	0.05	8 (2%) 62 71	10, 24, 51, 69	0
2	I	20/20 (100%)	-0.09	2 (10%) 9 14	12, 21, 65, 77	0
All	All	354/370 (95%)	0.04	10 (2%) 56 66	10, 24, 52, 77	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	23	HIS	4.0
1	E	318	PHE	3.7
1	E	311	GLU	3.6
2	I	24	ASP	3.5
1	E	253	GLY	3.2
1	E	39	HIS	3.0
1	E	336	ARG	2.9
1	E	334	GLU	2.5
1	E	339	ILE	2.3
1	E	298	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	E	197	11/12	0.96	0.09	-	14,19,21,21	0
1	SEP	E	338	10/11	0.70	0.19	-	49,52,55,55	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	IQB	E	351	27/27	0.95	0.13	-0.03	20,25,39,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.