wwPDB X-ray Structure Validation Summary Report (i)

Jan 31, 2016 – 11:44 PM GMT

PDB ID	:	1YFI
Title	:	Crystal Structure of restriction endonuclease MspI in complex with its cognate DNA in P212121 space group
Authors	:	Xu, Q.S.; Kucera, R.B.; Roberts, R.J.; Guo, H.-C.
Deposited on	:	2005-01-02
Resolution	:	2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	10	40% 30% 20% 10%
1	D	10	60% 30% 10%
1	E	10	70% 10% 10% 10%
1	F	10	60% 40%
2	A	262	77% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	262	% 73% 26%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*CP*CP*CP*CP*GP*GP*GP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	9	180	85	35	52	8	0	0	0
			Total	C	N	O	P			
1	D	9	183	86	37	52	8	0	0	0
			Total	C	N	O	P			
1	E	10	202	95	40	58	9	0	0	0
			Total	C	N	O	P			
1	F	10	202	95	40	58	9	0	0	0

- Molecule 2 is a protein called Type II restriction enzyme MspI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	262	2103	1328	367	404	4	0	0	0
			Total	C	N	O	S			
2	B	262	2112	1333	368	407	4	0	1	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	30	30	30	0	0
			Total	O		
3	B	15	15	15	0	0
			Total	O		
3	C	1	1	1	0	0
			Total	O		
3	D	3	3	3	0	0
			Total	O		
3	E	3	3	3	0	0
			Total	O		
3	F	1	1	1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*CP*CP*CP*CP*CP*GP*GP*GP*GP*G)-3'

Chain C:

- Molecule 1: 5'-D(*CP*CP*CP*CP*CP*GP*GP*GP*GP*G)-3'

Chain D:



- Molecule 1: 5'-D(*CP*CP*CP*CP*CP*GP*GP*GP*GP*G)-3'

Chain E:

- Molecule 1: 5'-D(*CP*CP*CP*CP*CP*GP*GP*GP*GP*G)-3'

Chain F:



- Molecule 2: Type II restriction enzyme MspI

Chain A:



- Molecule 2: Type II restriction enzyme MspI

Chain B:


%
73%
26%
M1 R2 T3 E4 L5 L6 I15 L18 D28 R29 K32 L33 D44 I45 K49 R50 Y51 N52 T53 N54 E59 K60 D61 I62 D73 L74 D75 N76 I77 V80 L87 T101 F104 T105 F106 H107 N108 Q109 R112 L113 V114 P115 K125 V126 S127 I128
V140 K161 E176 P177 Y178 R179 F182 I183 R184 V187 T188 L189 R190 A191 G196 N197 I198 L199 H200 P201 D202 L203 V217 T218 N221 I230 A231 E232 G233 S234 K235 A236 R237 K238 P239 N246 W247 T248 Y249 F260 K261 G262

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	50.30Å 111.27Å 130.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.94 – 2.70 46.94 – 2.58	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.94-2.70) 99.5 (46.94-2.58)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	5.33 (at 2.58Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.226 , 0.274 0.226 , 0.274	Depositor DCC
R_{free} test set	2088 reflections (10.00%)	DCC
Wilson B-factor (Å^2)	31.9	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 27.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	5 of 23728 reflections (0.021%)	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	5035	wwPDB-VP
Average B, all atoms (Å^2)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.43 % of the origin peak, indicating pseudo translational symmetry.The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.0019e-04.The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	C	0.49	0/201	0.77	0/308
1	D	0.45	0/205	0.78	1/315 (0.3%)
1	E	0.44	0/226	0.80	1/347 (0.3%)
1	F	0.40	0/226	0.70	0/347
2	A	0.39	0/2139	0.60	0/2878
2	B	0.39	0/2148	0.60	0/2890
All	All	0.40	0/5145	0.64	2/7085 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2
1	E	0	2
All	All	0	4

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	12	DC	O5'-C5'-C4'	6.04	126.10	111.00
1	E	6	DG	N9-C1'-C2'	-5.12	102.87	112.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	5	DC	Sidechain
1	C	6	DG	Sidechain
1	E	5	DC	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	E	6	DG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	180	0	101	4	0
1	D	183	0	101	4	0
1	E	202	0	112	2	0
1	F	202	0	112	6	0
2	A	2103	0	2133	38	0
2	B	2112	0	2138	54	0
3	A	30	0	0	2	0
3	B	15	0	0	1	0
3	C	1	0	0	0	0
3	D	3	0	0	0	0
3	E	3	0	0	0	0
3	F	1	0	0	0	0
All	All	5035	0	4697	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:73:ASP:HB3	2:B:75:ASP:OD1	1.78	0.83
2:B:18:LEU:HD11	2:B:29:ARG:HH11	1.44	0.81
2:A:246:ASN:HB2	2:A:261:LYS:HB2	1.64	0.79
2:B:18:LEU:HD11	2:B:29:ARG:NH1	1.98	0.77
1:F:14:DC:H2"	1:F:15:DC:H5"	1.70	0.73

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	260/262 (99%)	241 (93%)	18 (7%)	1 (0%)	39	69
2	B	261/262 (100%)	244 (94%)	16 (6%)	1 (0%)	39	69
All	All	521/524 (99%)	485 (93%)	34 (6%)	2 (0%)	39	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	54	ASN
2	A	159	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	235/235 (100%)	232 (99%)	3 (1%)	76	92
2	B	236/235 (100%)	231 (98%)	5 (2%)	61	87
All	All	471/470 (100%)	463 (98%)	8 (2%)	68	90

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	28	ASP
2	B	218	THR
2	B	73	ASP
2	A	237	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	29	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
2	A	158	GLN
2	B	246	ASN
2	B	52	ASN
2	A	52	ASN
2	B	158	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	C	9/10 (90%)	-0.22	0	100	100	10, 18, 35, 45	0
1	D	9/10 (90%)	-0.43	0	100	100	15, 22, 24, 26	0
1	E	10/10 (100%)	-0.14	0	100	100	15, 29, 46, 51	0
1	F	10/10 (100%)	-0.26	0	100	100	22, 26, 32, 38	0
2	A	262/262 (100%)	-0.23	0	100	100	7, 23, 39, 54	0
2	B	262/262 (100%)	-0.08	2 (0%)	87	88	8, 31, 47, 66	0
All	All	562/564 (99%)	-0.16	2 (0%)	93	94	7, 26, 46, 66	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	MET	3.2
2	B	4	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.