



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 08:30 PM GMT

PDB ID : 4YIF  
Title : Crystal structure of Rv0880  
Authors : Gao, Y.R.; Feng, N.; Li, D.F.; Bi, L.J.  
Deposited on : 2015-03-02  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	<div> <div></div> <div>73% 12% • 15%</div> </div>
1	B	164	<div> <div>5%</div> <div>68% 18% • 14%</div> </div>
1	C	164	<div> <div>5%</div> <div>71% 12% •• 15%</div> </div>
1	D	164	<div> <div>5%</div> <div>65% 15% • 20%</div> </div>
1	E	164	<div> <div></div> <div>68% 12% 20%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	164	<div><div></div><div>4%</div><div>62%</div><div>16%</div><div>21%</div></div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6710 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized HTH-type transcriptional regulator Rv0880.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	0	0
			1064	653	205	201	5			
1	B	141	Total	C	N	O	S	0	0	0
			1072	657	206	204	5			
1	C	139	Total	C	N	O	S	0	0	0
			1058	650	204	199	5			
1	D	132	Total	C	N	O	S	0	0	0
			1003	616	192	190	5			
1	E	131	Total	C	N	O	S	0	0	0
			995	611	190	189	5			
1	F	130	Total	C	N	O	S	0	0	0
			989	609	190	185	5			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP P9WMF1
A	-19	GLY	-	expression tag	UNP P9WMF1
A	-18	SER	-	expression tag	UNP P9WMF1
A	-17	SER	-	expression tag	UNP P9WMF1
A	-16	HIS	-	expression tag	UNP P9WMF1
A	-15	HIS	-	expression tag	UNP P9WMF1
A	-14	HIS	-	expression tag	UNP P9WMF1
A	-13	HIS	-	expression tag	UNP P9WMF1
A	-12	HIS	-	expression tag	UNP P9WMF1
A	-11	HIS	-	expression tag	UNP P9WMF1
A	-10	SER	-	expression tag	UNP P9WMF1
A	-9	SER	-	expression tag	UNP P9WMF1
A	-8	GLY	-	expression tag	UNP P9WMF1
A	-7	LEU	-	expression tag	UNP P9WMF1
A	-6	VAL	-	expression tag	UNP P9WMF1
A	-5	PRO	-	expression tag	UNP P9WMF1
A	-4	ARG	-	expression tag	UNP P9WMF1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P9WMF1
A	-2	SER	-	expression tag	UNP P9WMF1
A	-1	HIS	-	expression tag	UNP P9WMF1
A	0	MET	-	expression tag	UNP P9WMF1
A	1	VAL	-	expression tag	UNP P9WMF1
B	-20	MET	-	initiating methionine	UNP P9WMF1
B	-19	GLY	-	expression tag	UNP P9WMF1
B	-18	SER	-	expression tag	UNP P9WMF1
B	-17	SER	-	expression tag	UNP P9WMF1
B	-16	HIS	-	expression tag	UNP P9WMF1
B	-15	HIS	-	expression tag	UNP P9WMF1
B	-14	HIS	-	expression tag	UNP P9WMF1
B	-13	HIS	-	expression tag	UNP P9WMF1
B	-12	HIS	-	expression tag	UNP P9WMF1
B	-11	HIS	-	expression tag	UNP P9WMF1
B	-10	SER	-	expression tag	UNP P9WMF1
B	-9	SER	-	expression tag	UNP P9WMF1
B	-8	GLY	-	expression tag	UNP P9WMF1
B	-7	LEU	-	expression tag	UNP P9WMF1
B	-6	VAL	-	expression tag	UNP P9WMF1
B	-5	PRO	-	expression tag	UNP P9WMF1
B	-4	ARG	-	expression tag	UNP P9WMF1
B	-3	GLY	-	expression tag	UNP P9WMF1
B	-2	SER	-	expression tag	UNP P9WMF1
B	-1	HIS	-	expression tag	UNP P9WMF1
B	0	MET	-	expression tag	UNP P9WMF1
B	1	VAL	-	expression tag	UNP P9WMF1
C	-20	MET	-	initiating methionine	UNP P9WMF1
C	-19	GLY	-	expression tag	UNP P9WMF1
C	-18	SER	-	expression tag	UNP P9WMF1
C	-17	SER	-	expression tag	UNP P9WMF1
C	-16	HIS	-	expression tag	UNP P9WMF1
C	-15	HIS	-	expression tag	UNP P9WMF1
C	-14	HIS	-	expression tag	UNP P9WMF1
C	-13	HIS	-	expression tag	UNP P9WMF1
C	-12	HIS	-	expression tag	UNP P9WMF1
C	-11	HIS	-	expression tag	UNP P9WMF1
C	-10	SER	-	expression tag	UNP P9WMF1
C	-9	SER	-	expression tag	UNP P9WMF1
C	-8	GLY	-	expression tag	UNP P9WMF1
C	-7	LEU	-	expression tag	UNP P9WMF1
C	-6	VAL	-	expression tag	UNP P9WMF1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	PRO	-	expression tag	UNP P9WMF1
C	-4	ARG	-	expression tag	UNP P9WMF1
C	-3	GLY	-	expression tag	UNP P9WMF1
C	-2	SER	-	expression tag	UNP P9WMF1
C	-1	HIS	-	expression tag	UNP P9WMF1
C	0	MET	-	expression tag	UNP P9WMF1
C	1	VAL	-	expression tag	UNP P9WMF1
D	-20	MET	-	initiating methionine	UNP P9WMF1
D	-19	GLY	-	expression tag	UNP P9WMF1
D	-18	SER	-	expression tag	UNP P9WMF1
D	-17	SER	-	expression tag	UNP P9WMF1
D	-16	HIS	-	expression tag	UNP P9WMF1
D	-15	HIS	-	expression tag	UNP P9WMF1
D	-14	HIS	-	expression tag	UNP P9WMF1
D	-13	HIS	-	expression tag	UNP P9WMF1
D	-12	HIS	-	expression tag	UNP P9WMF1
D	-11	HIS	-	expression tag	UNP P9WMF1
D	-10	SER	-	expression tag	UNP P9WMF1
D	-9	SER	-	expression tag	UNP P9WMF1
D	-8	GLY	-	expression tag	UNP P9WMF1
D	-7	LEU	-	expression tag	UNP P9WMF1
D	-6	VAL	-	expression tag	UNP P9WMF1
D	-5	PRO	-	expression tag	UNP P9WMF1
D	-4	ARG	-	expression tag	UNP P9WMF1
D	-3	GLY	-	expression tag	UNP P9WMF1
D	-2	SER	-	expression tag	UNP P9WMF1
D	-1	HIS	-	expression tag	UNP P9WMF1
D	0	MET	-	expression tag	UNP P9WMF1
D	1	VAL	-	expression tag	UNP P9WMF1
E	-20	MET	-	initiating methionine	UNP P9WMF1
E	-19	GLY	-	expression tag	UNP P9WMF1
E	-18	SER	-	expression tag	UNP P9WMF1
E	-17	SER	-	expression tag	UNP P9WMF1
E	-16	HIS	-	expression tag	UNP P9WMF1
E	-15	HIS	-	expression tag	UNP P9WMF1
E	-14	HIS	-	expression tag	UNP P9WMF1
E	-13	HIS	-	expression tag	UNP P9WMF1
E	-12	HIS	-	expression tag	UNP P9WMF1
E	-11	HIS	-	expression tag	UNP P9WMF1
E	-10	SER	-	expression tag	UNP P9WMF1
E	-9	SER	-	expression tag	UNP P9WMF1
E	-8	GLY	-	expression tag	UNP P9WMF1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	-7	LEU	-	expression tag	UNP P9WMF1
E	-6	VAL	-	expression tag	UNP P9WMF1
E	-5	PRO	-	expression tag	UNP P9WMF1
E	-4	ARG	-	expression tag	UNP P9WMF1
E	-3	GLY	-	expression tag	UNP P9WMF1
E	-2	SER	-	expression tag	UNP P9WMF1
E	-1	HIS	-	expression tag	UNP P9WMF1
E	0	MET	-	expression tag	UNP P9WMF1
E	1	VAL	-	expression tag	UNP P9WMF1
F	-20	MET	-	initiating methionine	UNP P9WMF1
F	-19	GLY	-	expression tag	UNP P9WMF1
F	-18	SER	-	expression tag	UNP P9WMF1
F	-17	SER	-	expression tag	UNP P9WMF1
F	-16	HIS	-	expression tag	UNP P9WMF1
F	-15	HIS	-	expression tag	UNP P9WMF1
F	-14	HIS	-	expression tag	UNP P9WMF1
F	-13	HIS	-	expression tag	UNP P9WMF1
F	-12	HIS	-	expression tag	UNP P9WMF1
F	-11	HIS	-	expression tag	UNP P9WMF1
F	-10	SER	-	expression tag	UNP P9WMF1
F	-9	SER	-	expression tag	UNP P9WMF1
F	-8	GLY	-	expression tag	UNP P9WMF1
F	-7	LEU	-	expression tag	UNP P9WMF1
F	-6	VAL	-	expression tag	UNP P9WMF1
F	-5	PRO	-	expression tag	UNP P9WMF1
F	-4	ARG	-	expression tag	UNP P9WMF1
F	-3	GLY	-	expression tag	UNP P9WMF1
F	-2	SER	-	expression tag	UNP P9WMF1
F	-1	HIS	-	expression tag	UNP P9WMF1
F	0	MET	-	expression tag	UNP P9WMF1
F	1	VAL	-	expression tag	UNP P9WMF1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	130	Total O 130 130	0	0
2	B	104	Total O 104 104	0	0
2	C	74	Total O 74 74	0	0
2	D	69	Total O 69 69	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	80	Total	O	0	0
			80	80		
2	F	72	Total	O	0	0
			72	72		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uncharacterized HTH-type transcriptional regulator Rv0880




- Molecule 1: Uncharacterized HTH-type transcriptional regulator Rv0880




- Molecule 1: Uncharacterized HTH-type transcriptional regulator Rv0880




- Molecule 1: Uncharacterized HTH-type transcriptional regulator Rv0880



- Molecule 1: Uncharacterized HTH-type transcriptional regulator Rv0880



● Molecule 1: Uncharacterized HTH-type transcriptional regulator Rv0880



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.97Å 69.60Å 70.32Å 103.71° 111.06° 105.83°	Depositor
Resolution (Å)	37.11 – 2.00 41.15 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.0 (37.11-2.00) 77.8 (41.15-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, $R_{free}$	0.212 , 0.249 0.209 , 0.244	Depositor DCC
$R_{free}$ test set	2640 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.8	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 38.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 55010 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6710	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/1076	0.58	1/1456 (0.1%)
1	B	0.33	0/1084	0.61	0/1467
1	C	0.31	0/1070	0.63	1/1448 (0.1%)
1	D	0.30	0/1011	0.57	0/1365
1	E	0.30	0/1003	0.51	0/1354
1	F	0.37	1/997 (0.1%)	0.57	0/1346
All	All	0.32	1/6241 (0.0%)	0.58	2/8436 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	141	GLU	CG-CD	-5.48	1.43	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	83	HIS	C-N-CD	-6.21	106.93	120.60
1	A	67	ARG	NE-CZ-NH1	-5.25	117.68	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	29	SER	Peptide
1	C	85	ILE	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1064	0	1101	19	0
1	B	1072	0	1105	27	0
1	C	1058	0	1096	19	0
1	D	1003	0	1043	27	0
1	E	995	0	1035	15	0
1	F	989	0	1034	23	0
2	A	130	0	0	5	1
2	B	104	0	0	5	0
2	C	74	0	0	2	1
2	D	69	0	0	9	0
2	E	80	0	0	2	1
2	F	72	0	0	5	0
All	All	6710	0	6414	107	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:ARG:NH1	2:B:300:HOH:O	1.90	1.01
1:B:74:ASP:OD2	2:B:201:HOH:O	1.85	0.94
1:D:106:ARG:NH2	2:D:221:HOH:O	2.04	0.91
1:D:18:ARG:NH1	2:D:260:HOH:O	2.04	0.90
1:D:11:ASP:OD2	2:D:248:HOH:O	1.95	0.84

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:217:HOH:O	2:E:216:HOH:O[1_544]	1.91	0.29
2:A:214:HOH:O	2:A:218:HOH:O[1_655]	2.12	0.08

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/164 (84%)	136 (99%)	1 (1%)	1 (1%)	26	19
1	B	139/164 (85%)	135 (97%)	3 (2%)	1 (1%)	26	19
1	C	137/164 (84%)	130 (95%)	4 (3%)	3 (2%)	8	3
1	D	128/164 (78%)	125 (98%)	2 (2%)	1 (1%)	24	15
1	E	127/164 (77%)	126 (99%)	1 (1%)	0	100	100
1	F	126/164 (77%)	124 (98%)	1 (1%)	1 (1%)	24	15
All	All	795/984 (81%)	776 (98%)	12 (2%)	7 (1%)	21	13

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	88	ARG
1	C	86	ASP
1	D	9	ALA
1	C	84	PRO
1	B	85	ILE

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/136 (85%)	115 (100%)	0	100	100
1	B	116/136 (85%)	114 (98%)	2 (2%)	68	71
1	C	114/136 (84%)	114 (100%)	0	100	100
1	D	108/136 (79%)	107 (99%)	1 (1%)	84	88
1	E	108/136 (79%)	107 (99%)	1 (1%)	84	88
1	F	106/136 (78%)	105 (99%)	1 (1%)	84	88
All	All	667/816 (82%)	662 (99%)	5 (1%)	88	91

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	90	VAL
1	B	110	GLN
1	D	110	GLN
1	E	129	ARG
1	F	30	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	27	ASN
1	B	110	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	140/164 (85%)	-0.33	1 (0%) 89 89	21, 32, 55, 81	0
1	B	141/164 (85%)	-0.12	8 (5%) 27 29	22, 35, 74, 178	0
1	C	139/164 (84%)	0.12	9 (6%) 22 23	23, 36, 101, 163	0
1	D	132/164 (80%)	0.23	9 (6%) 20 22	21, 42, 83, 111	0
1	E	131/164 (79%)	-0.16	2 (1%) 76 77	25, 37, 60, 85	0
1	F	130/164 (79%)	0.03	7 (5%) 29 31	23, 36, 78, 147	0
All	All	813/984 (82%)	-0.04	36 (4%) 38 39	21, 36, 73, 178	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	86	ASP	10.0
1	C	85	ILE	10.0
1	C	83	HIS	9.1
1	C	84	PRO	8.1
1	E	90	VAL	8.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.