



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:46 PM GMT

PDB ID : 4YOY
Title : Crystal structure of a trimeric exonuclease PhoExo I from *Pyrococcus horikoshii* OT3 in complex with poly-dT and Mg²⁺ ion
Authors : Miyazono, K.; Ito, T.; Tanokura, M.
Deposited on : 2015-03-12
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div> <div>6%</div> <div>91%</div> <div>5% . .</div> </div>
1	C	233	<div> <div>6%</div> <div>92%</div> <div>5% .</div> </div>
1	E	233	<div> <div>20%</div> <div>91%</div> <div>5% .</div> </div>
2	B	7	<div> <div>43%</div> <div>14%</div> <div>43%</div> </div>
2	D	7	<div> <div>43%</div> <div>14%</div> <div>43%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	7	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	C	301	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5806 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-5 exonuclease PhoExo I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1781	1149	296	333	3			
1	C	226	Total	C	N	O	S	0	0	0
			1781	1149	296	333	3			
1	E	224	Total	C	N	O	S	0	0	0
			1766	1140	293	330	3			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	80	ASN	ASP	engineered mutation	UNP A0A060P168
A	230	HIS	-	expression tag	UNP A0A060P168
A	231	HIS	-	expression tag	UNP A0A060P168
A	232	HIS	-	expression tag	UNP A0A060P168
A	233	HIS	-	expression tag	UNP A0A060P168
C	80	ASN	ASP	engineered mutation	UNP A0A060P168
C	230	HIS	-	expression tag	UNP A0A060P168
C	231	HIS	-	expression tag	UNP A0A060P168
C	232	HIS	-	expression tag	UNP A0A060P168
C	233	HIS	-	expression tag	UNP A0A060P168
E	80	ASN	ASP	engineered mutation	UNP A0A060P168
E	230	HIS	-	expression tag	UNP A0A060P168
E	231	HIS	-	expression tag	UNP A0A060P168
E	232	HIS	-	expression tag	UNP A0A060P168
E	233	HIS	-	expression tag	UNP A0A060P168

- Molecule 2 is a DNA chain called 5'-D(*TP*TP*TP*TP*TP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	4	Total	C	N	O	P	0	0	0
			80	40	8	28	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	4	Total	C	N	O	P	0	0	0
			80	40	8	28	4			
2	F	4	Total	C	N	O	P	0	0	0
			80	40	8	28	4			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	95	Total	O	0	0
			95	95		
4	B	5	Total	O	0	0
			5	5		
4	C	89	Total	O	0	0
			89	89		
4	D	3	Total	O	0	0
			3	3		
4	E	41	Total	O	0	0
			41	41		
4	F	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-5 exonuclease PhoExo I




- Molecule 1: 3-5 exonuclease PhoExo I




- Molecule 1: 3-5 exonuclease PhoExo I



- Molecule 2: 5'-D(*TP*TP*TP*TP*TP*TP*T)-3'




- Molecule 2: 5'-D(*TP*TP*TP*TP*TP*TP*T)-3'



- Molecule 2: 5'-D(*TP*TP*TP*TP*TP*TP*T)-3'

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	94.88Å 94.89Å 106.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.91 – 1.95 19.91 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.91-1.95) 100.0 (19.91-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 1.94Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.180 , 0.204 0.187 , 0.208	Depositor DCC
R_{free} test set	3564 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 56.2	EDS
Estimated twinning fraction	0.028 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 70537 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5806	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1813	0.58	0/2450
1	C	0.41	0/1813	0.55	0/2450
1	E	0.31	0/1798	0.50	0/2431
2	B	0.92	0/87	1.39	0/132
2	D	0.78	0/87	1.33	0/132
2	F	0.71	0/87	1.38	0/132
All	All	0.42	0/5685	0.61	0/7727

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1781	0	1842	11	0
1	C	1781	0	1842	8	0
1	E	1766	0	1824	7	0
2	B	80	0	49	2	0
2	D	80	0	49	2	0
2	F	80	0	49	2	0
3	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	1	0	0	0	0
4	A	95	0	0	0	0
4	B	5	0	0	0	0
4	C	89	0	0	0	0
4	D	3	0	0	0	0
4	E	41	0	0	0	0
4	F	3	0	0	0	0
All	All	5806	0	5655	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:55:ARG:HH21	2:D:4:DT:H3	1.13	0.93
1:A:55:ARG:HH22	2:B:4:DT:H3	1.29	0.80
1:C:55:ARG:NH2	2:D:4:DT:H3	1.85	0.75
1:E:177:GLU:OE2	1:E:179:LYS:NZ	2.28	0.66
1:A:182:LYS:HE2	1:A:198:GLU:OE1	2.05	0.56
1:A:218:ARG:NH2	1:C:192:GLU:OE1	2.39	0.55
1:A:93:GLU:HG3	1:A:110:LYS:HD3	1.89	0.54
1:A:55:ARG:NH2	2:B:4:DT:H3	2.03	0.53
1:E:55:ARG:HH11	2:F:4:DT:H3	1.58	0.51
1:C:106:LYS:O	1:C:110:LYS:HG3	2.11	0.51
1:A:39:ARG:NH2	1:A:71:GLU:OE2	2.42	0.49
1:C:177:GLU:HG2	1:C:184:ILE:HB	1.95	0.48
1:E:55:ARG:HE	2:F:4:DT:H3	1.62	0.47
1:E:107:GLU:HG3	1:E:108:ILE:N	2.29	0.47
1:A:104:ARG:O	1:A:107:GLU:HG2	2.15	0.46
1:E:26:VAL:HG11	1:E:147:TYR:HB3	1.97	0.46
1:A:104:ARG:HD3	1:A:107:GLU:OE2	2.16	0.45
1:A:181:ASP:N	1:A:181:ASP:OD1	2.49	0.44
1:E:91:LEU:O	1:E:117:GLN:HG3	2.18	0.44
1:C:117:GLN:HB3	1:C:118:PRO:HD3	1.99	0.44
1:E:58:ILE:HD11	1:E:79:LEU:HG	2.00	0.44
1:A:117:GLN:HB3	1:A:118:PRO:HD3	2.01	0.43
1:C:122:LYS:HA	1:C:122:LYS:HD2	1.85	0.43
1:A:106:LYS:O	1:A:110:LYS:HG3	2.18	0.42
1:C:180:LYS:HB3	1:C:180:LYS:HE2	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/233 (95%)	218 (98%)	4 (2%)	0	100	100
1	C	222/233 (95%)	219 (99%)	3 (1%)	0	100	100
1	E	220/233 (94%)	217 (99%)	3 (1%)	0	100	100
All	All	664/699 (95%)	654 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/200 (96%)	191 (99%)	2 (1%)	82	80
1	C	193/200 (96%)	191 (99%)	2 (1%)	82	80
1	E	191/200 (96%)	188 (98%)	3 (2%)	70	66
All	All	577/600 (96%)	570 (99%)	7 (1%)	78	75

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	ARG
1	A	182	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	52	LEU
1	C	104	ARG
1	E	93	GLU
1	E	104	ARG
1	E	107	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/233 (96%)	0.33	13 (5%) 26 37	24, 41, 84, 112	0
1	C	226/233 (96%)	0.39	13 (5%) 26 37	27, 45, 93, 122	0
1	E	224/233 (96%)	1.04	46 (20%) 1 1	33, 68, 123, 148	0
2	B	4/7 (57%)	-0.14	0 100 100	47, 56, 61, 86	0
2	D	4/7 (57%)	0.09	0 100 100	52, 71, 88, 111	0
2	F	4/7 (57%)	-0.02	0 100 100	63, 75, 85, 112	0
All	All	688/720 (95%)	0.58	72 (10%) 8 13	24, 51, 108, 148	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	53	SER	9.7
1	E	52	LEU	7.3
1	E	53	SER	5.4
1	C	229	SER	5.0
1	E	124	TRP	4.9
1	E	103	ASP	4.9
1	C	52	LEU	4.9
1	E	93	GLU	4.7
1	C	100	GLN	4.6
1	E	227	SER	4.6
1	E	101	ILE	4.5
1	E	205	GLN	4.4
1	E	100	GLN	4.4
1	E	206	GLY	4.1
1	E	122	LYS	3.9
1	C	51	ASP	3.9
1	E	28	VAL	3.9
1	E	54	GLY	3.8
1	A	100	GLN	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	97	ASP	3.7
1	A	205	GLN	3.6
1	C	49	ASN	3.5
1	A	53	SER	3.4
1	E	51	ASP	3.4
1	E	94	SER	3.4
1	E	109	TRP	3.4
1	E	168	VAL	3.4
1	E	126	GLU	3.3
1	E	50	TYR	3.2
1	E	202	GLU	3.2
1	E	125	GLU	3.1
1	E	4	VAL	3.1
1	C	93	GLU	3.1
1	E	208	LYS	3.0
1	E	153	VAL	3.0
1	E	49	ASN	2.9
1	E	115	ASP	2.9
1	E	107	GLU	2.8
1	E	56	GLN	2.8
1	A	54	GLY	2.8
1	E	114	LYS	2.8
1	E	128	GLY	2.7
1	A	52	LEU	2.6
1	E	79	LEU	2.6
1	C	94	SER	2.6
1	A	15	GLU	2.6
1	A	103	ASP	2.5
1	E	121	LYS	2.5
1	A	229	SER	2.5
1	A	168	VAL	2.4
1	E	98	ALA	2.4
1	C	109	TRP	2.4
1	A	109	TRP	2.3
1	E	207	ILE	2.3
1	C	164	GLY	2.3
1	E	222	VAL	2.3
1	C	104	ARG	2.3
1	C	227	SER	2.3
1	A	104	ARG	2.3
1	A	153	VAL	2.2
1	E	88	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	165	GLY	2.2
1	A	131	ILE	2.2
1	E	204	PRO	2.2
1	E	104	ARG	2.1
1	E	5	ALA	2.1
1	E	102	SER	2.1
1	C	50	TYR	2.1
1	E	41	LEU	2.1
1	E	200	LYS	2.1
1	E	91	LEU	2.0
1	E	15	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	C	301	1/1	0.92	0.16	2.61	62,62,62,62	0
3	MG	E	301	1/1	0.94	0.08	-3.98	95,95,95,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.