



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 06:25 PM BST

PDB ID : 1YR1  
Title : Structure of the major extracytoplasmic domain of the trans isomer of the bacterial cell division protein divib from geobacillus stearothermophilus  
Authors : Robson, S.A.; King, G.F.  
Deposited on : 2005-02-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	119	

## 2 Ensemble composition and analysis ⓘ

This entry contains 25 models. Model 21 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:118-A:225 (108)	0.12	21

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 4, 11, 13, 17, 18, 19, 21, 22, 24
2	2, 3, 6, 7, 8, 9, 12, 14
3	5, 15, 16
4	10, 20
Single-model clusters	23; 25

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1869 atoms, of which 927 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called cell-division initiation protein.


Mol	Chain	Residues	Atoms						Trace
1	A	119	Total	C	H	N	O	S	0
			1869	598	927	162	179	3	

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: cell-division initiation protein




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1


- Molecule 1: cell-division initiation protein



#### 4.2.2 Score per residue for model 2

- Molecule 1: cell-division initiation protein






### 4.2.3 Score per residue for model 3

- Molecule 1: cell-division initiation protein


Chain A: 54% 33% 9%



### 4.2.4 Score per residue for model 4

- Molecule 1: cell-division initiation protein


Chain A: 55% 32% 9%



### 4.2.5 Score per residue for model 5


- Molecule 1: cell-division initiation protein

Chain A: 55% 30% 5% 9%




### 4.2.6 Score per residue for model 6

- Molecule 1: cell-division initiation protein




#### 4.2.7 Score per residue for model 7

- Molecule 1: cell-division initiation protein




#### 4.2.8 Score per residue for model 8

- Molecule 1: cell-division initiation protein




#### 4.2.9 Score per residue for model 9

- Molecule 1: cell-division initiation protein




### 4.2.10 Score per residue for model 10

- Molecule 1: cell-division initiation protein




### 4.2.11 Score per residue for model 11

- Molecule 1: cell-division initiation protein




### 4.2.12 Score per residue for model 12

- Molecule 1: cell-division initiation protein



### 4.2.13 Score per residue for model 13


- Molecule 1: cell-division initiation protein






### 4.2.14 Score per residue for model 14

- Molecule 1: cell-division initiation protein




### 4.2.15 Score per residue for model 15

- Molecule 1: cell-division initiation protein




### 4.2.16 Score per residue for model 16

- Molecule 1: cell-division initiation protein




### 4.2.17 Score per residue for model 17

- Molecule 1: cell-division initiation protein




### 4.2.18 Score per residue for model 18

- Molecule 1: cell-division initiation protein




### 4.2.19 Score per residue for model 19

- Molecule 1: cell-division initiation protein




### 4.2.20 Score per residue for model 20

- Molecule 1: cell-division initiation protein




### 4.2.21 Score per residue for model 21 (medoid)

- Molecule 1: cell-division initiation protein




#### 4.2.22 Score per residue for model 22

- Molecule 1: cell-division initiation protein




#### 4.2.23 Score per residue for model 23

- Molecule 1: cell-division initiation protein




#### 4.2.24 Score per residue for model 24

- Molecule 1: cell-division initiation protein



#### 4.2.25 Score per residue for model 25

- Molecule 1: cell-division initiation protein



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics (CANDID/CYANA) followed by simulated annealing (XPLOR)*.

Of the 60 calculated structures, 25 were deposited, based on the following criterion: *lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CANDID	structure solution	1.1
CYANA	structure solution	1.1
XPLOR	refinement	3.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6395
Number of chemical shift lists	2
Total number of shifts	2907
Number of shifts mapped to atoms	2907
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality ⓘ

### 6.1 Standard geometry ⓘ

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	861	845	845	36±3
All	All	21525	21125	21125	893

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:121:ILE:HD11	1:A:168:THR:HG22	0.97	1.34	8	25
1:A:133:LEU:HD11	1:A:164:ILE:HG23	0.88	1.46	9	25
1:A:121:ILE:CD1	1:A:168:THR:HG22	0.82	2.05	12	25
1:A:133:LEU:HD11	1:A:164:ILE:CG2	0.81	2.06	4	25
1:A:161:GLY:HA2	1:A:164:ILE:HD12	0.75	1.59	14	25
1:A:181:ALA:HB3	1:A:224:ILE:HG23	0.72	1.59	7	18
1:A:174:LEU:CD2	1:A:179:LEU:HD23	0.68	2.18	23	10
1:A:196:ARG:NH2	1:A:210:THR:HG23	0.67	2.03	24	1
1:A:174:LEU:HD11	1:A:179:LEU:HB3	0.65	1.66	25	15
1:A:178:VAL:HA	1:A:224:ILE:HG22	0.65	1.68	7	23
1:A:122:ALA:HB3	1:A:168:THR:HG23	0.64	1.69	12	25
1:A:118:TRP:HB2	1:A:134:LEU:HD13	0.63	1.69	7	25
1:A:129:THR:HG23	1:A:145:THR:O	0.62	1.94	1	25
1:A:196:ARG:NH1	1:A:210:THR:HG23	0.62	2.09	12	1

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:171:LEU:HD11	1:A:214:PHE:CE1	0.60	2.31	17	2
1:A:171:LEU:HA	1:A:174:LEU:HD22	0.60	1.74	23	10
1:A:174:LEU:HD12	1:A:175:PRO:O	0.59	1.97	1	15
1:A:131:PHE:HB2	1:A:139:LEU:HD21	0.59	1.73	22	25
1:A:122:ALA:HB2	1:A:171:LEU:HD23	0.59	1.75	9	1
1:A:129:THR:HG23	1:A:145:THR:C	0.57	2.19	8	25
1:A:121:ILE:HD11	1:A:168:THR:CG2	0.57	2.23	5	25
1:A:124:VAL:HG21	1:A:139:LEU:HG	0.56	1.77	7	25
1:A:122:ALA:HB2	1:A:171:LEU:CD2	0.56	2.31	9	25
1:A:133:LEU:CD1	1:A:155:LEU:HD12	0.55	2.31	16	25
1:A:174:LEU:HD21	1:A:179:LEU:HB3	0.55	1.78	23	10
1:A:134:LEU:HD12	1:A:138:ARG:CB	0.55	2.32	7	14
1:A:131:PHE:CB	1:A:139:LEU:HD21	0.54	2.33	2	25
1:A:218:LEU:HD12	1:A:224:ILE:CD1	0.54	2.33	7	10
1:A:124:VAL:HG21	1:A:139:LEU:CD1	0.53	2.32	22	25
1:A:124:VAL:HG21	1:A:139:LEU:CG	0.53	2.33	5	25
1:A:181:ALA:HB1	1:A:202:ASN:ND2	0.52	2.18	25	4
1:A:179:LEU:HD12	1:A:180:GLY:N	0.52	2.19	20	7
1:A:125:TYR:CE2	1:A:130:PHE:CE2	0.51	2.99	9	22
1:A:158:TRP:CZ3	1:A:187:TYR:CE2	0.50	2.99	22	5
1:A:181:ALA:CB	1:A:224:ILE:HG23	0.50	2.35	2	3
1:A:197:VAL:HG22	1:A:209:ALA:O	0.50	2.06	18	7
1:A:201:MET:SD	1:A:224:ILE:HD11	0.50	2.47	5	1
1:A:123:TYR:CE1	1:A:152:ALA:HB3	0.49	2.42	9	2
1:A:171:LEU:HB3	1:A:179:LEU:HD22	0.49	1.85	1	12
1:A:139:LEU:HD23	1:A:140:LEU:N	0.49	2.22	6	3
1:A:158:TRP:CE3	1:A:187:TYR:CE2	0.49	3.01	22	2
1:A:134:LEU:HD12	1:A:138:ARG:HB3	0.49	1.84	14	14
1:A:174:LEU:HD23	1:A:174:LEU:O	0.49	2.08	15	5
1:A:122:ALA:HB3	1:A:168:THR:CG2	0.48	2.38	12	14
1:A:174:LEU:O	1:A:174:LEU:HD23	0.48	2.07	23	5
1:A:179:LEU:C	1:A:179:LEU:HD12	0.48	2.28	1	13
1:A:175:PRO:HB2	1:A:178:VAL:HG23	0.48	1.84	22	13
1:A:179:LEU:HD12	1:A:179:LEU:C	0.48	2.29	10	9
1:A:158:TRP:CH2	1:A:211:ILE:HD11	0.47	2.44	7	12
1:A:124:VAL:HG21	1:A:139:LEU:HD11	0.47	1.86	22	9
1:A:140:LEU:N	1:A:140:LEU:HD22	0.46	2.25	24	4
1:A:174:LEU:HD11	1:A:179:LEU:CB	0.46	2.40	25	1
1:A:218:LEU:HD12	1:A:224:ILE:HD13	0.46	1.86	3	4
1:A:174:LEU:C	1:A:174:LEU:HD23	0.46	2.31	23	5
1:A:140:LEU:HD22	1:A:140:LEU:N	0.46	2.26	25	4

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:118:TRP:CG	1:A:136:ASN:ND2	0.46	2.83	25	3
1:A:124:VAL:HG11	1:A:139:LEU:HD11	0.46	1.87	9	6
1:A:133:LEU:HD22	1:A:168:THR:HG21	0.45	1.88	10	6
1:A:152:ALA:HB1	1:A:153:PRO:HD2	0.45	1.89	18	9
1:A:193:TYR:CD2	1:A:196:ARG:NH1	0.45	2.85	19	2
1:A:158:TRP:CE3	1:A:187:TYR:CD2	0.45	3.05	6	1
1:A:118:TRP:CE2	1:A:136:ASN:ND2	0.45	2.85	17	3
1:A:123:TYR:HE1	1:A:152:ALA:HB3	0.45	1.72	9	2
1:A:174:LEU:HD23	1:A:174:LEU:C	0.45	2.33	15	5
1:A:119:ARG:NH2	1:A:121:ILE:HG21	0.44	2.27	7	1
1:A:123:TYR:CD1	1:A:149:PRO:CG	0.44	3.00	9	2
1:A:131:PHE:CB	1:A:139:LEU:CD2	0.44	2.95	7	20
1:A:171:LEU:CD1	1:A:214:PHE:CE1	0.44	3.01	17	1
1:A:158:TRP:CD2	1:A:187:TYR:CD2	0.44	3.06	6	1
1:A:121:ILE:HD12	1:A:135:GLU:HA	0.43	1.91	2	4
1:A:186:HIS:CD2	1:A:200:TYR:CE1	0.43	3.07	10	1
1:A:145:THR:OG1	1:A:147:THR:HG22	0.42	2.14	6	3
1:A:207:VAL:HG12	1:A:208:SER:N	0.42	2.30	18	18
1:A:139:LEU:HD12	1:A:164:ILE:HD13	0.42	1.90	9	6
1:A:155:LEU:HD22	1:A:158:TRP:CZ2	0.42	2.48	9	1
1:A:186:HIS:CG	1:A:200:TYR:CE1	0.42	3.08	10	1
1:A:198:ILE:HD12	1:A:208:SER:HB3	0.42	1.92	14	1
1:A:129:THR:CG2	1:A:145:THR:N	0.42	2.83	18	23
1:A:140:LEU:CD2	1:A:140:LEU:N	0.42	2.83	25	4
1:A:217:LYS:O	1:A:220:HIS:CE1	0.42	2.73	22	2
1:A:140:LEU:N	1:A:140:LEU:CD2	0.42	2.83	18	3
1:A:199:VAL:HG22	1:A:207:VAL:HB	0.42	1.91	23	1
1:A:221:TYR:CB	1:A:224:ILE:HD12	0.42	2.45	14	2
1:A:181:ALA:HB1	1:A:202:ASN:OD1	0.42	2.15	17	1
1:A:158:TRP:CZ3	1:A:211:ILE:HD11	0.41	2.50	7	2
1:A:181:ALA:HB3	1:A:224:ILE:CG2	0.41	2.39	7	1
1:A:210:THR:O	1:A:214:PHE:N	0.41	2.54	1	11
1:A:201:MET:HE3	1:A:207:VAL:HG21	0.41	1.93	18	2
1:A:201:MET:HE1	1:A:224:ILE:CD1	0.41	2.46	20	1
1:A:187:TYR:CE2	1:A:189:PRO:CG	0.41	3.04	24	1
1:A:145:THR:OG1	1:A:147:THR:HG23	0.41	2.16	25	1
1:A:123:TYR:CD1	1:A:149:PRO:HG3	0.40	2.51	9	1
1:A:185:ILE:HG12	1:A:199:VAL:HG12	0.40	1.93	18	1
1:A:192:GLU:O	1:A:193:TYR:CD1	0.40	2.74	7	1
1:A:203:ASP:CB	1:A:205:TYR:CE2	0.40	3.05	5	1
1:A:218:LEU:N	1:A:218:LEU:HD22	0.40	2.32	25	1

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:218:LEU:HD22	1:A:218:LEU:N	0.40	2.31	21	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	108/119 (91%)	104±1 (96±1%)	3±1 (3±1%)	1±0 (1±0%)	19	64
All	All	2700/2975 (91%)	2594 (96%)	69 (3%)	37 (1%)	19	64

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	157	GLY	25
1	A	138	ARG	12

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/96 (92%)	71±2 (81±2%)	17±2 (19±2%)	5	38
All	All	2200/2400 (92%)	1784 (81%)	416 (19%)	5	38

All 36 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	121	ILE	25
1	A	221	TYR	25

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	125	TYR	25
1	A	146	LYS	25
1	A	168	THR	25
1	A	147	THR	25
1	A	162	ASP	25
1	A	167	MET	24
1	A	212	ARG	22
1	A	139	LEU	19
1	A	210	THR	17
1	A	138	ARG	17
1	A	201	MET	16
1	A	127	ARG	13
1	A	133	LEU	11
1	A	119	ARG	11
1	A	174	LEU	10
1	A	159	LYS	9
1	A	128	GLN	7
1	A	191	ARG	7
1	A	187	TYR	6
1	A	183	SER	6
1	A	184	GLU	6
1	A	196	ARG	6
1	A	182	MET	6
1	A	217	LYS	5
1	A	194	GLU	5
1	A	206	GLU	4
1	A	220	HIS	4
1	A	151	ASP	3
1	A	160	ASP	2
1	A	192	GLU	1
1	A	216	ASP	1
1	A	188	LYS	1
1	A	195	ASP	1
1	A	203	ASP	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 88% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 6395

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1454
Number of shifts mapped to atoms	1454
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	119	$-0.24 \pm 0.09$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	111	$0.16 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	118	$-0.01 \pm 0.12$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	110	$0.59 \pm 0.42$	None needed (imprecise)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 1183 atoms were assigned a chemical shift out of a possible 1337. 13 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	526/528 (100%)	209/210 (100%)	216/216 (100%)	101/102 (99%)
Sidechain	596/688 (87%)	364/403 (90%)	224/253 (89%)	8/32 (25%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	61/121 (50%)	59/63 (94%)	0/54 (0%)	2/4 (50%)
Overall	1183/1337 (88%)	632/676 (93%)	440/523 (84%)	111/138 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 1290 atoms were assigned a chemical shift out of a possible 1463. 14 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	576/583 (99%)	229/232 (99%)	237/238 (100%)	110/113 (97%)
Sidechain	653/759 (86%)	400/444 (90%)	244/278 (88%)	9/37 (24%)
Aromatic	61/121 (50%)	59/63 (94%)	0/54 (0%)	2/4 (50%)
Overall	1290/1463 (88%)	688/739 (93%)	481/570 (84%)	121/154 (79%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	119	ARG	NE	122.70	92.63 – 76.73	23.9
1	A	120	ARG	NE	122.60	92.63 – 76.73	23.8
1	A	187	TYR	HB3	0.42	4.75 – 0.95	-6.4

#### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



## 7.2 Chemical shift list 2

File name: BMRB entry 6395

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

### 7.2.1 Bookkeeping ⓘ

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1453
Number of shifts mapped to atoms	1453
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

### 7.2.2 Chemical shift referencing ⓘ

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	119	$-0.21 \pm 0.12$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	111	$0.07 \pm 0.12$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	118	$0.07 \pm 0.10$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	110	$0.54 \pm 0.49$	None needed (imprecise)

### 7.2.3 Completeness of resonance assignments ⓘ

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 1182 atoms were assigned a chemical shift out of a possible 1337. 10 out of 16 assigned methyl groups (LEU and VAL) were assigned

stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	526/528 (100%)	209/210 (100%)	216/216 (100%)	101/102 (99%)
Sidechain	595/688 (86%)	364/403 (90%)	223/253 (88%)	8/32 (25%)
Aromatic	61/121 (50%)	59/63 (94%)	0/54 (0%)	2/4 (50%)
Overall	1182/1337 (88%)	632/676 (93%)	439/523 (84%)	111/138 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 1289 atoms were assigned a chemical shift out of a possible 1463. 11 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	576/583 (99%)	229/232 (99%)	237/238 (100%)	110/113 (97%)
Sidechain	652/759 (86%)	400/444 (90%)	243/278 (87%)	9/37 (24%)
Aromatic	61/121 (50%)	59/63 (94%)	0/54 (0%)	2/4 (50%)
Overall	1289/1463 (88%)	688/739 (93%)	480/570 (84%)	121/154 (79%)

#### 7.2.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	119	ARG	NE	122.70	92.63 – 76.73	23.9
1	A	120	ARG	NE	122.60	92.63 – 76.73	23.8
1	A	187	TYR	HB3	0.42	4.75 – 0.95	-6.4

#### 7.2.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

