



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 08:47 PM GMT

PDB ID : 4YT0  
Title : Crystal structure of Mitochondrial rhodoquinol-fumarate reductase from *Ascaris suum* with 2-methyl-N-[3-(1-methylethoxy)phenyl]benzamide.  
Authors : Harada, S.; Shiba, T.; Sato, D.; Yamamoto, A.; Nagahama, M.; Yone, A.; Inaoka, D.K.; Sakamoto, K.; Inoue, M.; Honma, T.; Kita, K.  
Deposited on : 2015-03-17  
Resolution : 3.66 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1010 (3.82-3.50)
Clashscore	102246	1125 (3.82-3.50)
Ramachandran outliers	100387	1079 (3.82-3.50)
Sidechain outliers	100360	1078 (3.82-3.50)
RSRZ outliers	91569	1017 (3.82-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	
1	E	645	
2	B	282	
2	F	282	
3	C	188	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	G	188	<p>2% 66% 12% 20%</p>
4	D	156	<p>63% 19% 17%</p>
4	H	156	<p>68% 13% 17%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	HEM	C	201	-	-	-	X
11	MRN	C	202	-	-	-	X
11	MRN	G	202	-	-	-	X
5	MLI	E	701	-	-	X	-
8	SF4	B	302	-	-	X	-
8	SF4	F	302	-	-	X	-

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 18236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase flavoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	616	Total	C	N	O	S	0	0	0
			4787	3004	855	900	28			
1	E	616	Total	C	N	O	S	0	0	0
			4787	3004	855	900	28			

- Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	250	Total	C	N	O	S	0	0	0
			1985	1263	338	361	23			
2	F	250	Total	C	N	O	S	0	0	0
			1985	1263	338	361	23			


- Molecule 3 is a protein called Cytochrome b-large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	153	Total	C	N	O	S	0	0	0
			1217	813	204	194	6			
3	G	150	Total	C	N	O	S	0	0	0
			1195	798	201	190	6			

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	129	Total	C	N	O	S	0	0	0
			998	659	165	169	5			
4	H	129	Total	C	N	O	S	0	0	0
			998	659	165	169	5			

- Molecule 5 is MALONATE ION (three-letter code: MLI) (formula: C<sub>3</sub>H<sub>2</sub>O<sub>4</sub>).



- 
- The image displays the chemical structure of Flavin Adenine Dinucleotide (FAD). The structure is composed of several key components: an adenine base (a purine ring system with an amino group at position 6), a ribose sugar, a pyrophosphate group (two phosphate groups linked together), a ribitol chain (a five-carbon sugar alcohol), and a flavin mononucleotide (FMN) moiety. The FMN moiety consists of an isoalloxazine ring system, a dimethylbenzoyl side chain, and a phosphate group. The structure is shown in a 3D representation with stereochemistry indicated by wedge and dash bonds. The atoms are color-coded: carbon is grey, hydrogen is white, oxygen is red, nitrogen is blue, and phosphorus is orange. The structure is labeled with various atom numbers and charges.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			4	2	2		
7	F	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 8 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			8	4	4		
8	F	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 9 is FE3-S4 CLUSTER (three-letter code: F3S) (formula:  $\text{Fe}_3\text{S}_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			7	3	4		
9	F	1	Total	Fe	S	0	0
			7	3	4		

- 
- Chemical structure of HEM (Heme) showing a central iron atom coordinated by four nitrogen atoms in a porphyrin-like ring, with various side chains and a central heme group.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	G	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

-




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	C	1	Total	C	N	O	0	0
			20	17	1	2		
11	G	1	Total	C	N	O	0	0
			20	17	1	2		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinate dehydrogenase flavoprotein




- Molecule 1: Succinate dehydrogenase flavoprotein




- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial

Chain B: 




- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial

Chain F: 




- Molecule 3: Cytochrome b-large subunit

Chain C: 




- Molecule 3: Cytochrome b-large subunit

Chain G: 




- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial

Chain D: 



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.80Å 123.39Å 219.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.66 19.94 – 3.66	Depositor EDS
% Data completeness (in resolution range)	89.0 (20.00-3.66) 89.2 (19.94-3.66)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.83 (at 3.62Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.193 , 0.262 0.196 , 0.260	Depositor DCC
$R_{free}$ test set	1643 reflections (5.20%)	DCC
Wilson B-factor (Å <sup>2</sup> )	52.9	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 25.7	EDS
Estimated twinning fraction	0.058 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 33243 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	18236	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MLI, MRN, F3S, FES, HEM, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.34	0/4889	0.56	0/6605
1	E	0.34	0/4889	0.57	0/6605
2	B	0.35	0/2029	0.56	0/2739
2	F	0.35	0/2029	0.54	0/2739
3	C	0.36	0/1255	0.56	0/1709
3	G	0.35	0/1232	0.54	0/1676
4	D	0.38	0/1030	0.54	0/1406
4	H	0.35	0/1030	0.53	0/1406
All	All	0.35	0/18383	0.56	0/24885

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4787	0	4720	75	0
1	E	4787	0	4720	81	0
2	B	1985	0	2002	45	0
2	F	1985	0	2003	44	0
3	C	1217	0	1265	22	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	1195	0	1240	14	0
4	D	998	0	985	17	0
4	H	998	0	985	14	0
5	A	7	0	2	0	0
5	E	7	0	2	2	0
6	A	53	0	31	8	0
6	E	53	0	31	8	0
7	B	4	0	0	0	0
7	F	4	0	0	0	0
8	B	8	0	0	2	0
8	F	8	0	0	3	0
9	B	7	0	0	0	0
9	F	7	0	0	0	0
10	C	43	0	30	6	0
10	G	43	0	30	4	0
11	C	20	0	19	1	0
11	G	20	0	19	2	0
All	All	18236	0	18084	290	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 290 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:79:HIS:NE2	6:E:702:FAD:HM82	1.75	1.02
2:B:182:CYS:HG	8:B:302:SF4:FE3	0.79	1.00
1:A:79:HIS:CE1	6:A:702:FAD:HM82	2.10	0.86
1:A:79:HIS:NE2	6:A:702:FAD:C8M	2.44	0.80
2:F:185:CYS:HG	8:F:302:SF4:FE1	0.93	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	614/645 (95%)	568 (92%)	43 (7%)	3 (0%)	34	77
1	E	614/645 (95%)	563 (92%)	51 (8%)	0	100	100
2	B	248/282 (88%)	215 (87%)	28 (11%)	5 (2%)	9	54
2	F	248/282 (88%)	219 (88%)	22 (9%)	7 (3%)	6	47
3	C	151/188 (80%)	139 (92%)	11 (7%)	1 (1%)	26	72
3	G	148/188 (79%)	134 (90%)	11 (7%)	3 (2%)	9	54
4	D	127/156 (81%)	119 (94%)	5 (4%)	3 (2%)	7	51
4	H	127/156 (81%)	114 (90%)	11 (9%)	2 (2%)	12	58
All	All	2277/2542 (90%)	2071 (91%)	182 (8%)	24 (1%)	17	65

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	THR
2	B	140	ASN
2	F	140	ASN
2	F	243	MET
4	D	54	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	502/527 (95%)	471 (94%)	31 (6%)	23	65
1	E	502/527 (95%)	472 (94%)	30 (6%)	24	66
2	B	220/242 (91%)	206 (94%)	14 (6%)	22	64
2	F	220/242 (91%)	204 (93%)	16 (7%)	17	59
3	C	127/158 (80%)	113 (89%)	14 (11%)	8	40
3	G	124/158 (78%)	111 (90%)	13 (10%)	8	42

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	D	98/119 (82%)	89 (91%)	9 (9%)	11	49
4	H	98/119 (82%)	92 (94%)	6 (6%)	23	66
All	All	1891/2092 (90%)	1758 (93%)	133 (7%)	19	61

5 of 133 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	83	GLU
1	E	205	LEU
3	G	110	LEU
4	D	84	MET
4	D	154	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	165	GLN
1	E	117	ASN
2	F	145	GLN
4	D	105	ASN
4	D	140	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	MLI	A	701	-	0,6,6	0.00	-	0,7,7	0.00	-
6	FAD	A	702	-	52,58,58	1.29	5 (9%)	52,89,89	2.22	13 (25%)
7	FES	B	301	2	0,4,4	0.00	-	0,4,4	0.00	-
8	SF4	B	302	2	0,12,12	0.00	-	0,24,24	0.00	-
9	F3S	B	303	2	0,9,9	0.00	-	0,15,15	0.00	-
10	HEM	C	201	3,4	24,50,50	0.99	2 (8%)	16,82,82	1.90	2 (12%)
11	MRN	C	202	-	21,21,21	1.99	3 (14%)	28,28,28	1.13	2 (7%)
5	MLI	E	701	-	0,6,6	0.00	-	0,7,7	0.00	-
6	FAD	E	702	-	52,58,58	1.36	5 (9%)	52,89,89	2.37	12 (23%)
7	FES	F	301	2	0,4,4	0.00	-	0,4,4	0.00	-
8	SF4	F	302	2	0,12,12	0.00	-	0,24,24	0.00	-
9	F3S	F	303	2	0,9,9	0.00	-	0,15,15	0.00	-
10	HEM	G	201	3,4	24,50,50	0.94	2 (8%)	16,82,82	1.71	2 (12%)
11	MRN	G	202	-	21,21,21	2.02	3 (14%)	28,28,28	1.44	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MLI	A	701	-	-	0/0/4/4	0/0/0/0
6	FAD	A	702	-	-	0/30/50/50	0/6/6/6
7	FES	B	301	2	-	0/0/4/4	0/1/1/1
8	SF4	B	302	2	-	0/0/48/48	0/6/5/5
9	F3S	B	303	2	-	0/0/24/24	0/0/3/3
10	HEM	C	201	3,4	-	0/6/54/54	0/0/8/8
11	MRN	C	202	-	-	0/12/12/12	0/2/2/2
5	MLI	E	701	-	-	0/0/4/4	0/0/0/0
6	FAD	E	702	-	-	0/30/50/50	0/6/6/6
7	FES	F	301	2	-	0/0/4/4	0/1/1/1
8	SF4	F	302	2	-	0/0/48/48	0/6/5/5
9	F3S	F	303	2	-	0/0/24/24	0/0/3/3

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	HEM	G	201	3,4	-	2/6/54/54	0/0/8/8
11	MRN	G	202	-	-	0/12/12/12	0/2/2/2

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	C	202	MRN	C3-C8	-5.60	1.38	1.50
11	G	202	MRN	C9-N	-5.35	1.31	1.41
11	C	202	MRN	C1-C2	-5.19	1.40	1.51
11	G	202	MRN	C3-C8	-5.12	1.39	1.50
11	G	202	MRN	C1-C2	-5.06	1.41	1.51

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	702	FAD	N3A-C2A-N1A	-7.82	122.73	128.87
6	A	702	FAD	N3A-C2A-N1A	-7.70	122.82	128.87
6	E	702	FAD	C4-C4X-C10	-6.49	115.79	119.94
11	G	202	MRN	C9-N-C8	-5.26	113.61	126.78
11	C	202	MRN	C9-N-C8	-4.86	114.61	126.78

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	G	201	HEM	C2D-C3D-CAD-CBD
10	G	201	HEM	C4D-C3D-CAD-CBD

There are no ring outliers.

9 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	702	FAD	8	0
8	B	302	SF4	2	0
10	C	201	HEM	6	0
11	C	202	MRN	1	0
5	E	701	MLI	2	0
6	E	702	FAD	8	0
8	F	302	SF4	3	0
10	G	201	HEM	4	0
11	G	202	MRN	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	616/645 (95%)	-0.68	0 100 100	29, 50, 76, 96	1 (0%)
1	E	616/645 (95%)	-0.64	1 (0%) 95 92	30, 59, 85, 119	1 (0%)
2	B	250/282 (88%)	-0.64	1 (0%) 93 88	30, 53, 87, 115	0
2	F	250/282 (88%)	-0.71	0 100 100	30, 51, 75, 100	0
3	C	153/188 (81%)	-0.54	0 100 100	44, 61, 106, 126	0
3	G	150/188 (79%)	-0.43	4 (2%) 58 40	42, 67, 133, 161	0
4	D	129/156 (82%)	-0.58	0 100 100	44, 64, 90, 125	0
4	H	129/156 (82%)	-0.44	1 (0%) 87 76	45, 70, 112, 135	0
All	All	2293/2542 (90%)	-0.62	7 (0%) 94 90	29, 57, 91, 161	2 (0%)

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	114	TRP	3.4
2	B	33	LYS	3.3
1	E	31	ASN	2.9
3	G	183	THR	2.2
4	H	51	PRO	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
11	MRN	C	202	20/20	0.91	0.32	3.63	62,68,83,83	0
11	MRN	G	202	20/20	0.88	0.33	3.44	83,90,127,129	0
10	HEM	C	201	43/43	0.94	0.22	2.09	66,72,76,78	0
10	HEM	G	201	43/43	0.95	0.20	1.87	69,75,77,78	0
6	FAD	E	702	53/53	0.96	0.15	0.30	33,35,39,40	0
6	FAD	A	702	53/53	0.97	0.13	-0.37	24,26,29,30	0
9	F3S	B	303	7/7	0.99	0.13	-0.69	39,41,45,46	0
5	MLI	A	701	7/7	0.98	0.12	-0.90	41,43,46,46	0
9	F3S	F	303	7/7	0.99	0.12	-0.96	38,41,45,47	0
5	MLI	E	701	7/7	0.97	0.12	-1.43	48,49,50,51	0
7	FES	B	301	4/4	0.99	0.09	-1.51	22,22,23,23	0
8	SF4	F	302	8/8	1.00	0.10	-1.51	28,28,29,29	0
8	SF4	B	302	8/8	1.00	0.10	-1.83	26,27,27,28	0
7	FES	F	301	4/4	1.00	0.07	-2.82	29,29,30,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.